

# Equilibrium Statistical Mechanics of Fermion Lattice Systems

Huzihiro Araki <sup>\*</sup> and Hajime Moriya <sup>†</sup>

## Abstract

We study equilibrium statistical mechanics of Fermion lattice systems which require a different treatment compared with spin lattice systems due to the non-commutativity of local algebras for disjoint regions.

Our major result is the equivalence of the KMS condition and the variational principle with a minimal assumption for the dynamics and without any explicit assumption on the potential. Its proof applies to spin lattice systems as well, yielding a vast improvement over known results.

All formulations are in terms of a  $\mathbf{C}^*$ -dynamical systems for the Fermion (CAR) algebra  $\mathcal{A}$  with all or a part of the following assumptions :

- (I) The interaction is even, namely, the dynamics  $\alpha_t$  commutes with the even-oddness automorphism  $\Theta$ . (Automatically satisfied when (IV) is assumed.)
- (II) The domain of the generator  $\delta_\alpha$  of  $\alpha_t$  contains the set  $\mathcal{A}_o$  of all strictly local elements of  $\mathcal{A}$ .
- (III) The set  $\mathcal{A}_o$  is the core of  $\delta_\alpha$ .
- (IV) The dynamics  $\alpha_t$  commutes with lattice translation automorphism group  $\tau$  of  $\mathcal{A}$ .

A major technical tool is the conditional expectation from  $\mathcal{A}$  onto its  $\mathbf{C}^*$ -subalgebras  $\mathcal{A}(I)$  for any subset  $I$  of the lattice, which induces a system of commuting squares. This technique overcomes the lack of tensor product structures for Fermion systems and even simplifies many known arguments for spin lattice systems.

In particular, this tool is used for obtaining the isomorphism between the real vector space of all  $*$ -derivations with their domain  $\mathcal{A}_o$ , commuting with  $\Theta$ , and that of all  $\Theta$ -even standard potentials which satisfy a specific norm convergence condition for the one point interaction energy. This makes it possible to associate a unique standard potential to every dynamics satisfying (I) and (II). The convergence condition for the potential is a consequence of its definition in terms of the  $*$ -derivation and not an additional assumption.

If translation invariance is imposed on  $*$ -derivations and potentials, then the isomorphism is kept and the space of translation covariant standard potentials becomes a separable Banach space with respect to the norm of the one point interaction energy. This is a crucial basis for an application of convex analysis to the equivalence proof in the major result.

Everything goes in parallel for spin lattice systems without the evenness assumption (I).

---

<sup>\*</sup>Mailing address: Research Institute for Mathematical Sciences, Kyoto University, Kitashirakawa-Oiwakecho, Sakyo-ku, Kyoto, 606-8502 Japan

<sup>†</sup>Mailing address: Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization (KEK), 1-1 Oho, Tsukuba, Ibaraki, 305-0801, Japan.

# Contents

|          |                                                         |           |
|----------|---------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                     | <b>3</b>  |
| <b>2</b> | <b>Conditional Expectations</b>                         | <b>9</b>  |
| 2.1      | Basic Properties . . . . .                              | 9         |
| 2.2      | Geometrical Lemma . . . . .                             | 12        |
| 2.3      | Commuting Square . . . . .                              | 13        |
| <b>3</b> | <b>Entropy and Relative Entropy</b>                     | <b>14</b> |
| 3.1      | Definitions . . . . .                                   | 14        |
| 3.2      | Monotone Property . . . . .                             | 16        |
| 3.3      | Strong Subadditivity . . . . .                          | 16        |
| <b>4</b> | <b>Fermion Lattice Systems</b>                          | <b>17</b> |
| 4.1      | Fermion Algebra . . . . .                               | 17        |
| 4.2      | Product Property of the Tracial State . . . . .         | 19        |
| 4.3      | Conditional Expectations for Fermion Algebras . . . . . | 20        |
| 4.4      | Commuting Squares for Fermion Algebras . . . . .        | 25        |
| 4.5      | Commutants of Subalgebras . . . . .                     | 25        |
| <b>5</b> | <b>Dynamics</b>                                         | <b>31</b> |
| 5.1      | Assumptions . . . . .                                   | 31        |
| 5.2      | Local Hamiltonians . . . . .                            | 32        |
| 5.3      | Internal Energy . . . . .                               | 36        |
| 5.4      | Potential . . . . .                                     | 37        |
| 5.5      | General Potential . . . . .                             | 40        |
| <b>6</b> | <b>KMS Condition</b>                                    | <b>41</b> |
| 6.1      | KMS Condition . . . . .                                 | 41        |
| 6.2      | Differential KMS Condition . . . . .                    | 42        |
| <b>7</b> | <b>Gibbs Condition</b>                                  | <b>44</b> |
| 7.1      | Inner Perturbation . . . . .                            | 44        |
| 7.2      | Surface Energy . . . . .                                | 45        |
| 7.3      | Gibbs Condition . . . . .                               | 46        |
| 7.4      | Equivalence to KMS Condition . . . . .                  | 47        |
| 7.5      | Product Form of the Gibbs Condition . . . . .           | 50        |
| <b>8</b> | <b>Translation Invariant Dynamics</b>                   | <b>52</b> |
| 8.1      | Translation Invariance and Covariance . . . . .         | 52        |
| 8.2      | Finite Range Potentials . . . . .                       | 57        |
| <b>9</b> | <b>Thermodynamic Limit</b>                              | <b>60</b> |
| 9.1      | Surface Energy Estimate . . . . .                       | 60        |
| 9.2      | Pressure . . . . .                                      | 63        |
| 9.3      | Mean Energy . . . . .                                   | 66        |

|                                                                      |           |
|----------------------------------------------------------------------|-----------|
| <b>10 Entropy for Fermion Systems</b>                                | <b>67</b> |
| 10.1 SSA for Fermion Systems . . . . .                               | 67        |
| 10.2 Mean Entropy . . . . .                                          | 68        |
| 10.3 Entropy Inequalities for Translation Invariant States . . . . . | 69        |
| <b>11 Variational Principle</b>                                      | <b>69</b> |
| 11.1 Extension of Even States . . . . .                              | 70        |
| 11.2 Variational Inequality . . . . .                                | 72        |
| 11.3 Variational Equality . . . . .                                  | 73        |
| 11.4 Variational Principle . . . . .                                 | 77        |
| <b>12 Equivalence of Variational Principle and KMS Condition</b>     | <b>78</b> |
| 12.1 Variational Principle from Gibbs Condition . . . . .            | 78        |
| 12.2 Some Tools of Convex Analysis . . . . .                         | 79        |
| 12.3 Differential KMS Condition from Variational Principle . . . . . | 84        |
| <b>13 Use of Other Entropy in the Variational Equality</b>           | <b>88</b> |
| 13.1 CNT-Entropy . . . . .                                           | 88        |
| 13.2 Variational Equality in Terms of CNT-Entropy . . . . .          | 89        |
| <b>14 Discussion</b>                                                 | <b>92</b> |
| <b>A Appendix: Van Hove Limit</b>                                    | <b>99</b> |
| A.1 Van Hove Net . . . . .                                           | 99        |
| A.2 Van Hove Limit . . . . .                                         | 102       |

# 1 Introduction

We investigate the equilibrium statistical mechanics of Fermion lattice systems. While equilibrium statistical mechanics of spin lattice systems has been well studied (see e.g. [17], [23] and [40]), there is a crucial difference between spin and Fermion cases. Namely, local algebras for disjoint regions commute elementwise for spin lattice systems, but do not commute for Fermion lattice systems. Due to this difference, the known formulations and proof in the case of spin lattice systems do not necessarily go over to the case of Fermion lattice systems and that is the motivation for this investigation. An example of a Fermion lattice system is the well-studied Hubbard model, to which our results apply.

It turned out that, in the matter of the equivalence of the KMS condition and the variational principle (i.e. the minimum free energy) for translation invariant states, we obtain its proof without any explicit assumption on the potential except for the condition that it is the standard potential corresponding to a translation invariant even dynamics, a minimal condition for a proper formulation of the problem. Without any change in the methods of proof, this strong result holds for spin lattice systems as well – a vast improvement over known results for spin lattice systems and a solution of a problem posed by Bratteli and Robinson (Remark after Theorem 6.2.42. [17]). In addition to this major result, we hope that the present work supplies a general mathematical foundation for equilibrium statistical mechanics of Fermion lattice systems, which was lacking so far.

There are two distinctive features of our approach. One feature is the central role of the time derivative (i.e. the generator of the dynamics). On one hand, this enables us to deal with all types of potentials without any explicit conditions on their long range or many body behavior, as long as the first time derivative of strictly localized operators can be defined. On the other hand, the existence of the dynamics for a given potential is separated from the problems treated here and we can bypass that existence problem via Assumption (III) below.

Another feature is the use of conditional expectations instead of the tensor product structure traditionally used for spin lattice systems. They provide not only a substitute tool (for the tensor product structure), which is applicable for both spin and Fermion lattice systems, but also a method of estimates which does not use the norm of individual potentials, for which we do not impose any explicit condition.

The main subject of our paper is the characterization of equilibrium states in terms of the KMS condition and the variational principle, which have an entirely different appearance but are shown to be equivalent. They refer to canonical ensembles in the infinite volume limit. However, they also refer to grand canonical ensembles if the dynamics is modified by gauge transformations with respect to Fermion numbers [11]. Namely, in the language of potentials, we may add a one-body potential, which consists of the particle number operator(s) times c-number chemical potential(s), and then the canonical ensemble for the so-modified potential is the grand canonical ensemble for the original potential, so that the grand canonical ensemble can be studied as a canonical ensemble for a modified potential, which is in the scope of our theory.

For the sake of notational simplicity, our presentation is for the case of one Fermion at each lattice site. Our results and proofs hold without any essential change for more general case where a finite numbers of Fermions and finite spins coexist at each lattice site. The even-oddness in that case refers to the total Fermion number. For example, for Hubbard model, there are two Fermions at each lattice site, representing the two components of a spin 1/2 Fermion.

Our starting point is a  $\mathbf{C}^*$ -dynamical system  $(\mathcal{A}, \alpha_t)$ , where  $\mathcal{A}$  is the  $\mathbf{C}^*$ -algebra of Fermion creation and annihilation operators on lattice sites of  $\mathbb{Z}^\nu$  with local subalgebras  $\mathcal{A}(I)$  for finite subsets  $I \subset \mathbb{Z}^\nu$  and  $\alpha_t$  is a given strongly continuous one-parameter group of  $*$ -automorphisms of  $\mathcal{A}$ .

Since the normal starting point in statistical mechanics is a potential, a digression on our formulation and strategy starting from a given dynamics may be appropriate at this point. The KMS condition, which is formulated in terms of the dynamics, is one of two main components of our equivalence result. On the other hand, the variational principle, which is formulated in terms of the potential, is the other main component. Therefore both dynamics and potential are indispensable for our main results and their mutual relation is of at most importance.

The key equation for that relation is the following formula. For any operator  $A$  localized in a finite subset  $I$  of the lattice, its time derivative is given by

$$\frac{d}{dt}\alpha_t(A) = \alpha_t(i[H(I), A])$$

where  $H(I)$  is described as a sum of potentials  $\Phi(J)$ , based on a finite subset  $J$  of the lattice, the sum being over all  $J$  except those  $J$  for which  $\Phi(J)$  commutes with any  $A$  localized in  $I$ , thus  $H(I)$  depending on  $I$ .

The problem of construction of  $\alpha_t$  from a given class of potentials is not a straight-forward task and has been studied by many people. As a result, a large number of results are known for quantum spin lattice systems (see e.g. [17]) and most of them can be applied to Fermion lattice systems. There are also some specific analyses for Fermion lattice systems (see e.g. [29]).

In parallel, the equivalence of the KMS condition and the variational principle for translation invariant states has been proved for a wide class of potentials for quantum spin lattice systems. The same proof also works for Fermion lattice systems in most cases; for example this is the case for finite range potentials (see e.g. page 113 of [30]).

While these results cover a wide range of explicit models, it seems difficult to decide exactly which class of potentials determine a dynamics and to show the equivalence in question in most general cases (which is not explicitly known) from the potential point of view.

In the present work, we do not intend to make any contribution to the problem of either construction of a dynamics from a potential, or giving a complete criterion for potentials, which give rise to a unique dynamics. (Thus we do not directly contribute to the study of explicit models.)

On the contrary, we avoid these difficult problems by assuming that the dynamics is already given (since this is needed in any case for the KMS condition) and prove the equivalence result in question under minimal (general) assumptions on the dynamics, explained immediately below.

Note that we do not make any explicit assumptions about the existence of a potential for a given dynamics nor about its property (such as the absolute convergence of the sum defining  $H(I)$  in terms of the potential).

For any given dynamics, for which all finitely localized operators have the time derivative at  $t = 0$  (Assumption (II) below) and which is lattice translation invariant (Assumption (IV) below), we show the existence of a corresponding potential, of which  $H(I)$  is a sum (as in usual formulation) convergent in a well-defined sense.

We now explain our assumptions and interconnection of dynamics with potentials in more detail. The following two assumptions make it possible to associate a potential to any given dynamics satisfying them.

(I) The dynamics is even. In other words,  $\alpha_t \Theta = \Theta \alpha_t$  for any  $t \in \mathbb{R}$ , where  $\Theta$  is an involutive automorphism of  $\mathcal{A}$ , multiplying  $-1$  on all creation and annihilation operators.

(II) The domain  $D(\delta_\alpha)$  of the generator  $\delta_\alpha$  of  $\alpha_t$  includes  $\mathcal{A}_o$ , the union of all  $\mathcal{A}(I)$  for all finite subsets  $I$  of the lattice.

It should be noted that Assumption (I) follows from Assumption (IV) below. (See Proposition 8.1.)

We denote by  $\Delta(\mathcal{A}_o)$  the set of all  $*$ -derivations with  $\mathcal{A}_o$  as their domain and their values in  $\mathcal{A}$ , commuting with  $\Theta$  (on  $\mathcal{A}_o$ ). Then the generator  $\delta_\alpha$  of our  $\alpha_t$ , when restricted to  $\mathcal{A}_o$ , belongs to  $\Delta(\mathcal{A}_o)$ .

It is shown that  $\Delta(\mathcal{A}_o)$  is in one-to-one correspondence with the set  $\mathcal{P}$  of standard even potentials, which are functionals  $\Phi(I)$  of all finite subsets  $I$  of the lattice with values in the self-adjoint  $\Theta$ -even part of the local algebra  $\mathcal{A}(I)$ ,

satisfying our standardness condition and a topological convergence condition (Theorem 5.13).

The topological convergence condition (( $\Phi$ -e) in Definition 5.10) is required in order that the potential is associated with a  $*$ -derivation on  $\mathcal{A}_o$  and refers to the convergence of the interaction energy operator for every finite subset  $I$

$$H(I) = \sum_K \{\Phi(K); K \cap I \neq \emptyset\},$$

where a finite sum is first taken over  $K$  contained in a finite subset  $J$  and the limit of  $J$  tending to the whole lattice is to converge in the norm topology of  $\mathcal{A}$ . (If this condition is satisfied for every one-point set  $I = \{n\}$  ( $n \in \mathbb{Z}^\nu$ ), then it is satisfied for all finite subsets  $I$ .) Note the difference from conventional topological conditions, such as summability of  $\|\Phi(I)\|$  over all  $I$  containing a point  $n$ , which are assumed for the sake of mathematical convenience.

For  $\Phi \in \mathcal{P}$ , internal energy  $U(I)$  and surface energy  $W(I)$  are also given in terms of  $\Phi$  by the conventional formulae for every finite  $I$ .

The connection of the derivation  $\delta$  and the corresponding potential  $\Phi$  is given by

$$\delta A = i[H(I), A] \quad (A \in \mathcal{A}(I)).$$

Due to the  $\Theta$ -evenness assumption (I), the replacement of  $H(I)$  by  $H(K)$  with  $K \supset I$  gives the same  $\delta$  on  $\mathcal{A}(I)$ , a necessary condition for consistency.

The standardness (( $\Phi$ -d) in Definition 5.10) is formulated in terms of conditional expectations and picks up a unique potential for each  $\delta \in \Delta(\mathcal{A}_o)$ . Without the standardness condition, there are many different potentials (called equivalent potentials) which yield exactly the same  $\delta$  through the above formulae. Through the one-to-one correspondence between  $\delta(\in \Delta(\mathcal{A}_o))$  and  $\Phi(\in \mathcal{P})$ , any dynamics  $\alpha_t$  satisfying our standing assumptions (I) and (II) is associated with a unique standard potential  $\Phi \in \mathcal{P}$ . This is a crucial point of our formulation, leading to our major result.

When we want to derive a statement involving  $\alpha_t$  from a condition involving the potential  $\Phi$ , we need the following assumption, guaranteeing the unique determination of  $\alpha_t$  from the given  $\Phi$ :

(III)  $\mathcal{A}_o$  is the core of the generator  $\delta_\alpha$  of the dynamics  $\alpha_t$ .

For the discussion of variational principle, we need the translation invariance assumption for the dynamics:

(IV)  $\alpha_t \tau_k = \tau_k \alpha_t$ , where  $\tau_k$ ,  $k \in \mathbb{Z}^\nu$ , is the automorphism group of  $\mathcal{A}$  representing the lattice translations.

The above Assumptions (I) - (IV) are the only assumptions needed for our theory below. On the other hand, if a potential  $\Phi$  (say, in the class  $\mathcal{P}$ ) is first given for any model, it is a hard problem in general to show that the corresponding derivation  $\delta_\Phi \in \Delta(\mathcal{A}_o)$  is given by some dynamics satisfying Assumptions (II) and (III), or equivalently that the closure of  $\delta_\Phi$  is a generator of a dynamics (i.e. it can be exponentiated to a one-parameter group of automorphisms of  $\mathcal{A}$ ).

We now present our main theorem after the explanation about the variational principle and its ingredients. The set  $\mathcal{P}_\tau$  of all translation covariant potentials in  $\mathcal{P}$  forms a Banach space (Proposition 8.8) with respect to the norm

$$\|\Phi\| \equiv \|H(\{n\})\|,$$

which is independent of the lattice point  $n$ . The finite range potentials are shown to be dense in  $\mathcal{P}_\tau$  with respect to this norm and to imply separability of  $\mathcal{P}_\tau$  (Theorem 8.12 and Corollary 8.13).

In terms of this norm, we obtain the energy estimate

$$\|U(I)\| \leq \|H(I)\| \leq \|\Phi\| \cdot |I|,$$

where  $|I|$  is the cardinality of  $I$  (Lemma 8.6). Then the conventional estimate for  $W(I)$  follows. These estimates are used to show the existence of the thermodynamic functionals, such as pressure  $P(\Phi)$  and mean energy  $e_\Phi(\omega)$ . All these estimates are carried out by the technique of conditional expectations without using the norm of the individual  $\Phi(I)$ .

For any state  $\omega$  of  $\mathcal{A}$ , its local entropy  $S_{\mathcal{A}(I)}(\omega) = S(\omega|_{\mathcal{A}(I)})$  is given as usual by the von Neumann entropy  $S(\cdot)$ . Due to the non-commutativity of local algebras for disjoint regions, not all known properties of entropy for spin lattice systems hold for our Fermion case [33]. However, the strong subadditivity of entropy (SSA) for Fermion systems holds. Then the existence of the mean entropy  $s(\omega)$  for any translation invariant state  $\omega$  for Fermion lattice systems follows by a known method of spin lattice systems.

The variational principle refers to the following equation for a translation invariant state  $\varphi$  of  $\mathcal{A}$  for a given translation covariant potential  $\Phi(\in \mathcal{P}_\tau)$  and  $\beta \in \mathbb{R}$ :

$$P(\beta\Phi) = s(\varphi) - \beta e_\Phi(\varphi) \tag{1.1}$$

Our major result can be formulated as the following two theorems.

**Theorem A.** *Under Assumptions (II) and (IV) for the dynamics  $\alpha_t$ , any translation invariant state, which satisfies the KMS condition for  $\alpha_t$  at the inverse temperature  $\beta$ , is a solution of the equation (1.1), where  $\Phi$  is the unique standard potential corresponding to  $\alpha_t$ .*

**Theorem B.** *Under Assumptions (II), (III) and (IV) for the dynamics  $\alpha_t$ , any solution  $\varphi$  of (1.1) satisfies the KMS condition for  $\alpha_t$  at  $\beta$ .*

*Remark.* These two theorems hold also for spin lattice systems.

We now present an over-all picture of the proof of our main results above. The proof of Theorem A and Theorem B will be carried out through the following steps:

- (1) KMS condition  $\Rightarrow$  Gibbs condition.
- (2) Gibbs condition  $\Rightarrow$  Variational principle.
- (3) Variational principle  $\Rightarrow$  dKMS condition on  $\mathcal{A}_\circ$ .
- (4) dKMS condition on  $\mathcal{A}_\circ \Rightarrow$  dKMS condition on  $D(\delta_\alpha)$ .
- (5) dKMS condition on  $D(\delta_\alpha) \Rightarrow$  KMS condition.

Assumptions (I) and (II) are used throughout (1)-(5). Assumption (IV) is used for the formulation of the variational principle and necessarily for (2) and (3). It is also used to derive Assumption (I), which is not included in the premise of Theorems A and B.

Assumption (III) is used only for (4).

The differential KMS (abbreviated as dKMS) condition in (4) and (5) refers to a known condition, which is entirely described in terms of the generator  $\delta_\alpha$  of  $\alpha_t$  and without use of  $\alpha_t$  (Definition 6.3). This condition on the full domain  $D(\delta_\alpha)$  of the generator  $\delta_\alpha$  of  $\alpha_t$  is known to be equivalent to the KMS condition (which is Step (5)). The differential KMS condition for our purpose is the condition for the restriction of  $\delta_\alpha$  to  $\mathcal{A}_0$ . Thus we need to show Step (4) using the additional assumption (III) on  $\alpha_t$ .

For Steps (1) and (2), we follow the proof for spin lattice systems in principle. However, the Gibbs condition for Fermion lattice systems requires a careful definition. We define the Gibbs condition for a state  $\varphi$  as the requirement that the local algebra  $\mathcal{A}(I)$  is in the centralizer of the perturbed functional  $\varphi^{\beta H(I)}$ , which is obtained from  $\varphi$  by a perturbation  $\beta H(I)$ , for each finite subset  $I$  of the lattice (Definition 7.1 and Lemma 7.2). When  $\mathcal{A}(I)$  and  $\mathcal{A}(I^c)$  commute (as in the case of spin lattice systems), this condition reduces to the product type characterization which was introduced and called the Gibbs condition by Araki and Ion for quantum spin lattice systems [5]. With our definition of the Gibbs condition, we have been able to prove Steps (1) and (2).

The product type characterization mentioned above is the condition that  $\varphi^{\beta H(I)}$  is the product of the tracial state of  $\mathcal{A}(I)$  and its restriction to the complement algebra  $\mathcal{A}(I^c)$ . In the present case of Fermion lattice systems, we show that a Gibbs state satisfies this condition if and only if it is an even state of  $\mathcal{A}$  (Proposition 7.7).

The same kind of formulation and result are valid for a perturbation  $\beta W(I)$ .

For Step (3) as well as for the proof of the variational equality

$$P(\beta\Phi) = \sup_{\omega \in \mathcal{A}_{+,1}^\tau} \left\{ s(\omega) - \beta e_\Phi(\omega) \right\}, \quad (1.2)$$

which is crucial for the variational principle, we need a product state of local Gibbs state. For this purpose, we have a technical result about the existence of a joint extension from states of local algebras for disjoint subsets of the lattice to a state of the algebra for their union, which holds if the individual states are even possibly except one (Theorem 11.2).

As an aside, the converse of Step (1) is shown under Assumptions (I), (II) and (III) (Theorem 7.6).

A major tool of our analysis is the  $\mathbf{C}^*$ -algebra conditional expectation  $E_I : \mathcal{A} \mapsto \mathcal{A}(I)$  with respect to the unique tracial state  $\tau$  of  $\mathcal{A}$ . Its existence is shown not only for finite subsets but for all subsets  $I$  of the lattice (Theorem 4.7). Based on the product property of  $\tau$  for subalgebras  $\mathcal{A}(I)$  and  $\mathcal{A}(J)$  for disjoint  $I$  and  $J$ , we obtain the following commuting square of  $\mathbf{C}^*$ -subalgebras (Theorem 4.13)



for Fermion systems. (It holds trivially for spin systems.)

$$\begin{array}{ccc} \mathcal{A}(I \cup J) & \xrightarrow{E_I} & \mathcal{A}(I) \\ E_J \downarrow & & \downarrow E_{I \cap J} \\ \mathcal{A}(J) & \xrightarrow{E_{I \cap J}} & \mathcal{A}(I \cap J). \end{array}$$

This serves as a replacement for the tensor-product structure in traditional arguments for spin lattice systems.

As by-products, we obtain a few useful results on the CAR algebra: The even-odd automorphism  $\Theta$  is shown to be outer for any infinite CAR algebra (Corollary 4.20) and formulae for commutants of  $\mathcal{A}(I)$  and  $\mathcal{A}(I)_+$  in  $\mathcal{A}$  for finite and infinite  $I$  are obtained (Theorem 4.17 and Theorem 4.19).

Some more results contained in this paper are as follows.

We show the validity of the variational equality (1.2) when the Connes-Narnhofer-Thirring entropy  $h_\omega(\tau)$  with respect to the group of lattice translation automorphisms  $\tau$  is used in place of the mean entropy  $s(\omega)$  (Theorem 13.2). Note that our system  $(\mathcal{A}, \tau)$ , where  $\tau$  denotes the group of lattice translation automorphisms, does not belong to the class of  $\mathbf{C}^*$ -systems considered in [34], being a non-abelian system.

We define general potentials as those which satisfy all conditions for those in  $\mathcal{P}$  except for the standardness. They include all potentials satisfying the following condition:

$$\sum_{I \ni n} \|\Phi(I)\| < \infty \tag{1.3}$$

for every lattice point  $n$ . For each general potential, the corresponding  $H(I)$  and  $\delta$  are defined and there is a unique standard potential in  $\mathcal{P}$  with the same  $\delta$  as a given general potential as described earlier.

Restricting our attention to those general potentials satisfying (1.3) (a condition which is introduced also in some discussion of spin lattice systems), we are able to show by a straightforward argument that the set of solutions of variational principle for a general translation covariant potential satisfying (1.3) coincide with those for the equivalent standard potential (which is automatically translation covariant) (Remark 1 to Proposition 14.1), although the pressure and the mean energy may be different between the two potentials.

## 2 Conditional Expectations

### 2.1 Basic Properties

The following proposition is well-known (see, e.g., Proposition 2.36, Chapter V [43]).

**Proposition 2.1.** *Let  $\mathcal{M}$  be a von Neumann algebra with a faithful normal tracial state  $\tau$  and  $\mathcal{N}$  be its von Neumann subalgebra. Then there exists a unique conditional expectation*

$$E_{\mathcal{N}}^{\mathcal{M}} : a \in \mathcal{M} \rightarrow E_{\mathcal{N}}^{\mathcal{M}}(a) \in \mathcal{N}$$

satisfying

$$\tau(ab) = \tau(E_{\mathcal{N}}^{\mathcal{M}}(a)b) \quad (2.1)$$

for any  $b \in \mathcal{N}$ .

*Remark.* A conditional expectation  $E_{\mathcal{N}}^{\mathcal{M}}$  is linear, positive, unital, and satisfies

$$E_{\mathcal{N}}^{\mathcal{M}}(ab) = E_{\mathcal{N}}^{\mathcal{M}}(a)b, \quad E_{\mathcal{N}}^{\mathcal{M}}(ba) = bE_{\mathcal{N}}^{\mathcal{M}}(a), \quad (2.2)$$

for any  $a \in \mathcal{M}$  and  $b \in \mathcal{N}$ , and

$$\|E_{\mathcal{N}}^{\mathcal{M}}\| = 1. \quad (2.3)$$

We shall obtain a  $\mathbf{C}^*$ -version of this proposition for the Fermion algebra in § 4, where  $\mathcal{M}$  and  $\mathcal{N}$  are  $\mathbf{C}^*$ -algebras with a unique tracial state  $\tau$ . The main step of its proof is the existence of  $E_{\mathcal{N}}^{\mathcal{M}}(a) \in \mathcal{N}$  for every  $a \in \mathcal{M}$  satisfying (2.1). Once it is established, the map  $E_{\mathcal{N}}^{\mathcal{M}}$  is a conditional expectation by standard argument, which we formulate for the sake of completeness as follows.

**Lemma 2.2.** *Let  $\mathcal{M}$  be a unital  $\mathbf{C}^*$ -algebra with a faithful tracial state  $\tau$  and  $\mathcal{N}$  be its subalgebra containing the identity of  $\mathcal{M}$ . Suppose that for every  $a \in \mathcal{M}$  there exists an element  $E_{\mathcal{N}}^{\mathcal{M}}(a)$  of  $\mathcal{N}$  satisfying (2.1). Then the map  $E_{\mathcal{N}}^{\mathcal{M}}$  from  $\mathcal{M}$  to  $\mathcal{N}$  is the unique conditional expectation from  $\mathcal{M}$  to  $\mathcal{N}$  with respect to  $\tau$ , possessing the following properties:*

- (1)  $E_{\mathcal{N}}^{\mathcal{M}}$  is linear, positive and unital map from  $\mathcal{M}$  onto  $\mathcal{N}$ .
- (2) For any  $a \in \mathcal{M}$  and  $b \in \mathcal{N}$ ,

$$E_{\mathcal{N}}^{\mathcal{M}}(ab) = E_{\mathcal{N}}^{\mathcal{M}}(a)b, \quad E_{\mathcal{N}}^{\mathcal{M}}(ba) = bE_{\mathcal{N}}^{\mathcal{M}}(a).$$

- (3)  $E_{\mathcal{N}}^{\mathcal{M}}$  is a projection of norm 1.

*Proof.* First we prove the uniqueness of  $E_{\mathcal{N}}^{\mathcal{M}}(a) \in \mathcal{N}$  satisfying (2.1) for a given  $a \in \mathcal{M}$ . Let  $a'$  and  $a''$  in  $\mathcal{N}$  satisfy (2.1), namely,

$$\tau(ab) = \tau(a'b) = \tau(a''b)$$

for all  $b \in \mathcal{N}$ . Then

$$\tau(b(a' - a'')) = 0.$$

By taking  $b = (a' - a'')^*$  and using the faithfulness of  $\tau$ , we obtain  $a' - a'' = 0$ , hence the uniqueness of  $E_{\mathcal{N}}^{\mathcal{M}}(a) \in \mathcal{N}$  for each  $a \in \mathcal{M}$ .

Except for the positivity, (1) and (2) can be shown in the same pattern as follows. Let  $a = c_1a_1 + c_2a_2$  where  $a_1, a_2 \in \mathcal{M}$  and  $c_1, c_2 \in \mathbb{C}$ . Then for any  $b \in \mathcal{N}$ ,

$$\begin{aligned} \tau(ab) &= c_1\tau(a_1b) + c_2\tau(a_2b) = c_1\tau(E_{\mathcal{N}}^{\mathcal{M}}(a_1)b) + c_2\tau(E_{\mathcal{N}}^{\mathcal{M}}(a_2)b) \\ &= \tau(\{c_1E_{\mathcal{N}}^{\mathcal{M}}(a_1) + c_2E_{\mathcal{N}}^{\mathcal{M}}(a_2)\}b). \end{aligned}$$

Since  $c_1 E_{\mathcal{N}}^{\mathcal{M}}(a_1) + c_2 E_{\mathcal{N}}^{\mathcal{M}}(a_2) \in \mathcal{N}$ , the uniqueness already shown implies

$$c_1 E_{\mathcal{N}}^{\mathcal{M}}(a_1) + c_2 E_{\mathcal{N}}^{\mathcal{M}}(a_2) = E_{\mathcal{N}}^{\mathcal{M}}(a).$$

Therefore,  $E_{\mathcal{N}}^{\mathcal{M}}$  is linear.

In the same way, for any  $a \in \mathcal{M}$  and  $b \in \mathcal{N}$ ,

$$\tau(abb') = \tau(E_{\mathcal{N}}^{\mathcal{M}}(a)bb')$$

holds for all  $b' \in \mathcal{N}$  and hence

$$E_{\mathcal{N}}^{\mathcal{M}}(ab) = E_{\mathcal{N}}^{\mathcal{M}}(a)b.$$

Also

$$\begin{aligned} \tau(bab') &= \tau(ab'b) = \tau(E_{\mathcal{N}}^{\mathcal{M}}(a)b'b) \\ &= \tau(bE_{\mathcal{N}}^{\mathcal{M}}(a)b') \end{aligned}$$

implies

$$E_{\mathcal{N}}^{\mathcal{M}}(ba) = bE_{\mathcal{N}}^{\mathcal{M}}(a).$$

If  $a \in \mathcal{N}$ , then the identity  $\tau(ab) = \tau(E_{\mathcal{N}}^{\mathcal{M}}(a)b)$  with  $b \in \mathcal{N}$  and the uniqueness result imply

$$E_{\mathcal{N}}^{\mathcal{M}}(a) = a.$$

Therefore  $E_{\mathcal{N}}^{\mathcal{M}}$  is a map onto  $\mathcal{N}$ . By taking  $a = \mathbf{1}(\in \mathcal{N})$ , we have

$$E_{\mathcal{N}}^{\mathcal{M}}(\mathbf{1}) = \mathbf{1}.$$

Hence  $E_{\mathcal{N}}^{\mathcal{M}}$  is unital.

Since  $E_{\mathcal{N}}^{\mathcal{M}}(a) \in \mathcal{N}$  for any  $a \in \mathcal{M}$ , we have  $E_{\mathcal{N}}^{\mathcal{M}}(E_{\mathcal{N}}^{\mathcal{M}}(a)) = E_{\mathcal{N}}^{\mathcal{M}}(a)$ . Therefore  $E_{\mathcal{N}}^{\mathcal{M}}$  is a projection.

To show the positivity of the map  $E_{\mathcal{N}}^{\mathcal{M}}$ , we consider the GNS triplet for the tracial state  $\tau_{\mathcal{N}}$  of  $\mathcal{N}$  (which is the restriction of  $\tau$  to  $\mathcal{N}$ ) consisting of a Hilbert space  $\mathcal{H}_{\tau}^{\mathcal{N}}$ , a representation  $\pi_{\tau}^{\mathcal{N}}$  of  $\mathcal{N}$  on  $\mathcal{H}_{\tau}^{\mathcal{N}}$  and a unit vector  $\Omega_{\tau}^{\mathcal{N}} \in \mathcal{H}_{\tau}^{\mathcal{N}}$ , giving rise to the state  $\tau_{\mathcal{N}}(A) = \tau(A) = (\Omega_{\tau}^{\mathcal{N}}, \pi_{\tau}^{\mathcal{N}}(A)\Omega_{\tau}^{\mathcal{N}})$  for  $A \in \mathcal{N}$ .

If  $a \in \mathcal{M}$  and  $a \geq 0$ , then for  $b \in \mathcal{N}$

$$\begin{aligned} (\pi_{\tau}^{\mathcal{N}}(b)\Omega_{\tau}^{\mathcal{N}}, \pi_{\tau}^{\mathcal{N}}(E_{\mathcal{N}}^{\mathcal{M}}(a))\pi_{\tau}^{\mathcal{N}}(b)\Omega_{\tau}^{\mathcal{N}}) &= \tau_{\mathcal{N}}(b^*E_{\mathcal{N}}^{\mathcal{M}}(a)b) \\ &= \tau_{\mathcal{N}}(E_{\mathcal{N}}^{\mathcal{M}}(a)bb^*) = \tau(abb^*) = \tau(b^*ab) \geq 0. \end{aligned}$$

Since  $\pi_{\tau}^{\mathcal{N}}(b)\Omega_{\tau}^{\mathcal{N}}$ ,  $b \in \mathcal{N}$  is dense in  $\mathcal{H}_{\tau}^{\mathcal{N}}$ , we obtain

$$\pi_{\tau}^{\mathcal{N}}(E_{\mathcal{N}}^{\mathcal{M}}(a)) \geq 0.$$

Since  $\pi_{\tau}^{\mathcal{N}}$  is faithful,

$$E_{\mathcal{N}}^{\mathcal{M}}(a) \geq 0,$$

and the positivity of  $E_{\mathcal{N}}^{\mathcal{M}}$  is shown.

For any  $a \in \mathcal{M}$ , the faithfulness of  $\pi_\tau^\mathcal{N}$  implies

$$\begin{aligned}
& \|E_\mathcal{N}^\mathcal{M}(a)\| \\
= & \|\pi_\tau^\mathcal{N}(E_\mathcal{N}^\mathcal{M}(a))\| \\
= & \sup_{b_1, b_2 \in \mathcal{N}} \left\{ |(\pi_\tau^\mathcal{N}(b_1)\Omega_\tau^\mathcal{N}, \{\pi_\tau^\mathcal{N}(E_\mathcal{N}^\mathcal{M}(a))\}\pi_\tau^\mathcal{N}(b_2)\Omega_\tau^\mathcal{N})| ; \|\pi_\tau^\mathcal{N}(b_1)\Omega_\tau^\mathcal{N}\| \leq 1, \|\pi_\tau^\mathcal{N}(b_2)\Omega_\tau^\mathcal{N}\| \leq 1 \right\} \\
= & \sup_{b_1, b_2 \in \mathcal{N}} \left\{ |(\tau(b_1^*E_\mathcal{N}^\mathcal{M}(a)b_2))| ; \tau(b_1^*b_1) \leq 1, \tau(b_2^*b_2) \leq 1 \right\} \\
= & \sup_{b_1, b_2 \in \mathcal{N}} \left\{ |(\tau(b_1^*ab_2))| ; \tau(b_1^*b_1) \leq 1, \tau(b_2^*b_2) \leq 1 \right\} \\
= & \sup_{b_1, b_2 \in \mathcal{N}} \left\{ |(\pi_\tau^\mathcal{M}(b_1)\Omega_\tau^\mathcal{M}, \pi_\tau^\mathcal{M}(a)\pi_\tau^\mathcal{M}(b_2)\Omega_\tau^\mathcal{M})| ; \|\pi_\tau^\mathcal{M}(b_1)\Omega_\tau^\mathcal{M}\| \leq 1, \|\pi_\tau^\mathcal{M}(b_2)\Omega_\tau^\mathcal{M}\| \leq 1 \right\} \\
\leq & \|\pi_\tau^\mathcal{M}(a)\| = \|a\|,
\end{aligned} \tag{2.4}$$

where we have used the cyclicity of  $\pi_\tau^\mathcal{N}(\mathcal{N})$  for  $\mathcal{H}_\tau^\mathcal{N}$  for the second equality,

$$\tau(b_1^*E_\mathcal{N}^\mathcal{M}(a)b_2) = \tau(E_\mathcal{N}^\mathcal{M}(a)b_2b_1^*) = \tau(ab_2b_1^*) = \tau(b_1^*ab_2).$$

for the fourth equality, and the same computation backwards replacing  $\mathcal{N}$  by  $\mathcal{M}$  for the fifth equality. Due to  $E_\mathcal{N}^\mathcal{M}(\mathbf{1}) = \mathbf{1}$  and (2.4), we have

$$\|E_\mathcal{N}^\mathcal{M}\| = 1.$$

We have completed the proof.  $\square$

## 2.2 Geometrical Lemma

Let us consider finite type **I** factors (i.e., full matrix algebras)  $\mathcal{M}$  and  $\mathcal{N}$  such that  $\mathcal{M} \supset \mathcal{N}$ . We have the isomorphisms  $\mathcal{M} \simeq \mathcal{N} \otimes \mathcal{N}_1$ ,  $\mathcal{N} \simeq \mathcal{N} \otimes \mathbf{1}$ , and  $\tau = \tau_\mathcal{N} \otimes \tau_{\mathcal{N}_1}$  where  $\mathcal{N}_1 \equiv \mathcal{M} \cap \mathcal{N}'$  is a finite type **I** factor.

A conditional expectation satisfying (2.1) is given by the slice map:

$$E_\mathcal{N}^\mathcal{M}(bb_1) = \tau(b_1)b \quad (b \in \mathcal{N}, b_1 \in \mathcal{N}_1). \tag{2.5}$$

We give this  $E_\mathcal{N}^\mathcal{M}$  a geometrical picture which we find useful. We introduce the following inner product on  $\mathcal{M}$ :

$$\langle a, b \rangle \equiv \tau(a^*b), \quad (a, b \in \mathcal{M}).$$

$\mathcal{M}$  is then a (finite-dimensional) Hilbert space with this inner product. Let  $P_\mathcal{N}^\mathcal{M}$  be the orthogonal projection onto the subspace  $\mathcal{N}$  of  $\mathcal{M}$ . We show that  $P_\mathcal{N}^\mathcal{M}$  is the same as  $E_\mathcal{N}^\mathcal{M}$  as a map  $\mathcal{M} \mapsto \mathcal{N}$ .

**Lemma 2.3.** *With the notation above,*

$$P_\mathcal{N}^\mathcal{M}a = E_\mathcal{N}^\mathcal{M}(a). \tag{2.6}$$

for any  $a \in \mathcal{M}$ .

*Proof.* Any  $a \in \mathcal{M}$  can be decomposed as  $a = P_{\mathcal{N}}^{\mathcal{M}}a + a'$  where  $a' \in \mathcal{N}^{\perp}$ . For any  $b \in \mathcal{N}$ , we have  $b^* \in \mathcal{N}$  and hence

$$\begin{aligned}\tau(ab) &= \langle b^*, a \rangle = \langle b^*, P_{\mathcal{N}}^{\mathcal{M}}a \rangle + \langle b^*, a' \rangle \\ &= \langle b^*, P_{\mathcal{N}}^{\mathcal{M}}a \rangle = \tau((P_{\mathcal{N}}^{\mathcal{M}}a)b).\end{aligned}$$

Since  $P_{\mathcal{N}}^{\mathcal{M}}a \in \mathcal{N}$ , it follows from Proposition 2.1 that

$$P_{\mathcal{N}}^{\mathcal{M}}a = E_{\mathcal{N}}^{\mathcal{M}}(a).$$

□

### 2.3 Commuting Square

We introduce the following equivalent conditions for a commuting square. (See e.g. [21].)

**Proposition 2.4.** *Let  $\mathcal{M}, \mathcal{N}_1, \mathcal{N}_2$  and  $\mathcal{P}$  be finite type I factors satisfying*

$$\mathcal{M} \supset \mathcal{N}_1 \supset \mathcal{P}, \quad \mathcal{M} \supset \mathcal{N}_2 \supset \mathcal{P}.$$

*Then the following conditions are equivalent:*

- (1)  $E_{\mathcal{N}_1}^{\mathcal{M}}|_{\mathcal{N}_2} = E_{\mathcal{P}}^{\mathcal{N}_2}$
- (2)  $E_{\mathcal{N}_2}^{\mathcal{M}}|_{\mathcal{N}_1} = E_{\mathcal{P}}^{\mathcal{N}_1}$
- (3)  $\mathcal{P} = \mathcal{N}_1 \cap \mathcal{N}_2$  and  $E_{\mathcal{N}_1}^{\mathcal{M}}E_{\mathcal{N}_2}^{\mathcal{M}} = E_{\mathcal{N}_2}^{\mathcal{M}}E_{\mathcal{N}_1}^{\mathcal{M}}$
- (4)  $E_{\mathcal{N}_1}^{\mathcal{M}}E_{\mathcal{N}_2}^{\mathcal{M}} = E_{\mathcal{P}}^{\mathcal{M}}$
- (5)  $E_{\mathcal{N}_2}^{\mathcal{M}}E_{\mathcal{N}_1}^{\mathcal{M}} = E_{\mathcal{P}}^{\mathcal{M}}$ .

*Proof.* (1)  $\iff$  (4):

Assume (1). Let  $a \in \mathcal{M}$  and  $b \in \mathcal{P}$ . By the assumption, we have

$$E_{\mathcal{N}_1}^{\mathcal{M}}(E_{\mathcal{N}_2}^{\mathcal{M}}(a)) = E_{\mathcal{P}}^{\mathcal{N}_2}(E_{\mathcal{N}_2}^{\mathcal{M}}(a)) \in \mathcal{P}.$$

due to  $E_{\mathcal{N}_2}^{\mathcal{M}}(a) \in \mathcal{N}_2$ . On the other hand,

$$\begin{aligned}\tau(E_{\mathcal{N}_1}^{\mathcal{M}}(E_{\mathcal{N}_2}^{\mathcal{M}}(a))b) &= \tau(E_{\mathcal{N}_2}^{\mathcal{M}}(a)b) \quad (\text{due to } b \in (\mathcal{P} \subset \mathcal{N}_1)) \\ &= \tau(ab) \quad (\text{due to } b \in (\mathcal{P} \subset \mathcal{N}_2)).\end{aligned}$$

Hence  $E_{\mathcal{N}_1}^{\mathcal{M}}(E_{\mathcal{N}_2}^{\mathcal{M}}(a)) = E_{\mathcal{P}}^{\mathcal{M}}(a)$  and so  $E_{\mathcal{N}_1}^{\mathcal{M}}E_{\mathcal{N}_2}^{\mathcal{M}} = E_{\mathcal{P}}^{\mathcal{M}}$ .

The converse is obvious: for  $a \in \mathcal{N}_2$ , (4) implies

$$E_{\mathcal{P}}^{\mathcal{N}_2}(a) = E_{\mathcal{P}}^{\mathcal{M}}(a) = E_{\mathcal{N}_1}^{\mathcal{M}}E_{\mathcal{N}_2}^{\mathcal{M}}(a) = E_{\mathcal{N}_1}^{\mathcal{M}}(a)$$

and hence (1).

(2)  $\iff$  (5) :

Exactly the same proof as above, with  $\mathcal{N}_1$  and  $\mathcal{N}_2$  interchanged.

(4)  $\iff$  (3):

Assume (4). By Lemma 2.3, (4) implies

$$P_{\mathcal{N}_1}^{\mathcal{M}}P_{\mathcal{N}_2}^{\mathcal{M}} = P_{\mathcal{P}}^{\mathcal{M}}.$$

Taking adjoints, we obtain

$$P_{\mathcal{N}_2}^{\mathcal{M}} P_{\mathcal{N}_1}^{\mathcal{M}} = P_{\mathcal{P}}^{\mathcal{M}}.$$

This implies

$$E_{\mathcal{N}_2}^{\mathcal{M}} E_{\mathcal{N}_1}^{\mathcal{M}} = E_{\mathcal{P}}^{\mathcal{M}} = E_{\mathcal{N}_1}^{\mathcal{M}} E_{\mathcal{N}_2}^{\mathcal{M}},$$

the last equality being due to (4).

Due to  $\mathcal{N}_1 \supset \mathcal{P}$  and  $\mathcal{N}_2 \supset \mathcal{P}$ , we have  $\mathcal{P} \subset \mathcal{N}_1 \cap \mathcal{N}_2$ . If  $b \in \mathcal{N}_1 \cap \mathcal{N}_2$ , then

$$b = E_{\mathcal{N}_1}^{\mathcal{M}} E_{\mathcal{N}_2}^{\mathcal{M}}(b) = E_{\mathcal{P}}^{\mathcal{M}}(b) \in \mathcal{P}$$

by (4). Hence  $\mathcal{P} = \mathcal{N}_1 \cap \mathcal{N}_2$ . This completes the proof of (4)  $\implies$  (3).

Assume (3). For any  $a \in \mathcal{M}$ , (3) implies  $E_{\mathcal{N}_1}^{\mathcal{M}}(E_{\mathcal{N}_2}^{\mathcal{M}}(a)) = E_{\mathcal{N}_2}^{\mathcal{M}}(E_{\mathcal{N}_1}^{\mathcal{M}}(a)) \in \mathcal{N}_1 \cap \mathcal{N}_2 = \mathcal{P}$  because the range of  $E_{\mathcal{N}_1}^{\mathcal{M}}$  is  $\mathcal{N}_1$  and the range of  $E_{\mathcal{N}_2}^{\mathcal{M}}$  is  $\mathcal{N}_2$ . For any  $b \in \mathcal{P}$  and  $a \in \mathcal{M}$ ,

$$\tau(E_{\mathcal{N}_1}^{\mathcal{M}}(E_{\mathcal{N}_2}^{\mathcal{M}}(a))b) = \tau(E_{\mathcal{N}_2}^{\mathcal{M}}(a)b) = \tau(ab).$$

Hence  $E_{\mathcal{N}_1}^{\mathcal{M}}(E_{\mathcal{N}_2}^{\mathcal{M}}(a)) = E_{\mathcal{P}}^{\mathcal{M}}(a)$ . This implies (4).

(5)  $\iff$  (3):

Exactly the same proof as above, with  $\mathcal{N}_1$  and  $\mathcal{N}_2$  interchanged.  $\square$

### 3 Entropy and Relative Entropy

#### 3.1 Definitions

We introduce some definitions and related lemmas needed for formulation of the main result of this section.

**Lemma 3.1.** *Let  $\mathcal{M}$  be a finite type I factor.*

(i) *Let  $\varphi$  be a positive linear functional on  $\mathcal{M}$ . Then there exists a unique  $\hat{\rho}_{\varphi} \in \mathcal{M}_+$  (called adjusted density matrix) satisfying*

$$\varphi(a) = \tau(\hat{\rho}_{\varphi} a)$$

*for all  $a \in \mathcal{M}$ .*

(ii) *Let  $\mathcal{N}$  be a subfactor of  $\mathcal{M}$  and  $\varphi_{\mathcal{N}}$  be the restriction of  $\varphi$  to  $\mathcal{N}$ . Then*

$$\hat{\rho}_{\varphi_{\mathcal{N}}} = E_{\mathcal{N}}^{\mathcal{M}}(\hat{\rho}_{\varphi})$$

*Proof.* (i) is well-known.

(ii) For  $b \in \mathcal{N}$ ,  $\varphi_{\mathcal{N}}(b) = \varphi(b) = \tau(\hat{\rho}_{\varphi} b) = \tau(E_{\mathcal{N}}^{\mathcal{M}}(\hat{\rho}_{\varphi})b)$ . Since  $E_{\mathcal{N}}^{\mathcal{M}}(\hat{\rho}_{\varphi}) \in \mathcal{N}_+$ , we have

$$\hat{\rho}_{\varphi_{\mathcal{N}}} = E_{\mathcal{N}}^{\mathcal{M}}(\hat{\rho}_{\varphi}).$$

$\square$

*Remark.* The above definition of density matrix is given in terms of the tracial state in contrast to the standard definition using the matrix trace  $\mathbf{Tr}$ . Hence we use the word ‘adjusted’.

**Definition 3.2.** Let  $\hat{\rho}_\varphi$  be the adjusted density matrix of a positive linear functional  $\varphi$  of a finite type  $\mathbf{I}$  factor. Then

$$\widehat{S}(\varphi) \equiv -\varphi(\log \hat{\rho}_\varphi)$$

is called the adjusted entropy of  $\varphi$ .

*Remark.* The adjusted density matrix and the adjusted entropy for a type  $\mathbf{I}_n$  factor  $\mathcal{M}$  with the dimension  $\mathbf{Tr}(\mathbf{1}) = n$  are related to the usual ones by the following relations:

$$\hat{\rho}_\varphi = n\rho_\varphi, \quad \widehat{S}(\varphi) = S(\varphi) - \varphi(\mathbf{1}) \log n. \quad (3.1)$$

The range of the values of entropy is given by the following well-known lemma.

**Lemma 3.3.** If  $\mathcal{M}$  is a type  $\mathbf{I}_n$  factor and  $\varphi$  is a state of  $\mathcal{M}$ , then

$$0 \leq S(\varphi) \leq \log n. \quad (3.2)$$

The equality  $S(\varphi) = 0$  holds if and only if  $\varphi$  is a pure state of  $\mathcal{M}$ . The equality  $S(\varphi) = \log n$  holds if and only if  $\varphi$  is the tracial state  $\tau$  of  $\mathcal{M}$ .

**Definition 3.4.** The relative entropy of  $\varrho$  and  $\sigma$  in  $\mathcal{M}_+$  as well as that of positive linear functionals  $\varphi$  and  $\psi$  are defined by

$$S(\sigma, \varrho) = \tau(\varrho(\log \varrho - \log \sigma)) \quad (3.3)$$

$$S(\psi, \varphi) = \varphi(\log \hat{\rho}_\varphi - \log \hat{\rho}_\psi) (= \tau(\hat{\rho}_\varphi \log \hat{\rho}_\varphi - \hat{\rho}_\varphi \log \hat{\rho}_\psi)). \quad (3.4)$$

*Remark.*  $S(\psi, \varphi)$  remains the same if  $\hat{\rho}_\varphi$  and  $\hat{\rho}_\psi$  are replaced by the density matrices  $\rho_\varphi$  and  $\rho_\psi$  with respect to  $\mathbf{Tr}$ . The right-hand sides of (3.3) and (3.4) are well-defined when  $\varrho, \sigma, \hat{\rho}_\varphi$  and  $\hat{\rho}_\psi$  are regular. Otherwise, one may define them as the limit of regular cases, for example by taking the limit  $\varepsilon \rightarrow 0$  for  $(1-\varepsilon)\varphi + \varepsilon\tau, (1-\varepsilon)\psi + \varepsilon\tau$  for (3.4), and similarly for (3.3). The value of  $S(\psi, \varphi)$  is real or  $+\infty$  for positive linear functionals  $\varphi$  and  $\psi$ .

The following lemma is also well-known.

**Lemma 3.5.** Let  $\varphi$  and  $\psi$  be states. Then  $S(\psi, \varphi)$  is non-negative. It vanishes if and only if  $\varphi = \psi$ .

*Remark.* We note that there are different notations for the relative entropy and that we adopt that of Araki [8] and Kosaki [25]. In comparison with our notation, the order of two states is reversed in that of Umegaki [45], while both the order of states and the sign are reversed in that of Bratteli and Robinson [17].

### 3.2 Monotone Property

Under any conditional expectation  $E$  and under restriction to any subalgebra, the relative entropy is known to be non-increasing:

$$S(\psi \circ E, \varphi \circ E) \leq S(\psi, \varphi), \quad (3.5)$$

$$S(\psi_{\mathcal{N}}, \varphi_{\mathcal{N}}) \leq S(\psi, \varphi). \quad (3.6)$$

(For example, (3.6) is Theorem 4.1(iv) of [25]. (3.5) follows from Theorem 4.1(v) of [25], because  $E$  is a Schwarz map [44].)

When we want to exhibit the dependence of entropy on  $\mathcal{M}$  more explicitly, we use the notation  $S_{\mathcal{M}}$  and  $\hat{S}_{\mathcal{M}}$  instead of  $S$  and  $\hat{S}$ . The relation between the entropy and the relative entropy for a state  $\varphi$  is given by

$$\hat{S}(\varphi) = -S(\tau, \varphi) = S(\varphi) - S(\tau).$$

Note that  $S(\tau) = \log n$  for a type  $\mathbf{I}_n$  factor  $\mathcal{M}$ .

We identify  $\mathcal{M}$  with  $\mathcal{N} \otimes (\mathcal{M} \cap \mathcal{N}')$  and use the notation  $\varphi_{\mathcal{N}} \otimes \tau_{\mathcal{M} \cap \mathcal{N}'}$ . We also identify  $A \in \mathcal{N} \subset \mathcal{M}$  with  $A \otimes \mathbf{1} \in \mathcal{N} \otimes (\mathcal{M} \cap \mathcal{N}')$ .

**Lemma 3.6.** *Let  $\mathcal{M} \supset \mathcal{N}$  be finite type  $\mathbf{I}$  factors, and  $\varphi$  be a state on  $\mathcal{M}$ . Then*

$$\hat{S}_{\mathcal{N}}(\varphi_{\mathcal{N}}) - \hat{S}_{\mathcal{M}}(\varphi) = S_{\mathcal{M}}(\varphi_{\mathcal{N}} \otimes \tau_{\mathcal{M} \cap \mathcal{N}'}, \varphi) = S_{\mathcal{M}}(\varphi \circ E_{\mathcal{N}}^{\mathcal{M}}, \varphi). \quad (3.7)$$

*Proof.* If  $\varphi$  is a faithful state, we show the above identity by a straight-forward calculation. If  $\varphi$  is not faithful, we add  $\varepsilon \cdot \tau$  to  $(1 - \varepsilon)\varphi$  and then take the limit  $\varepsilon \rightarrow 0$ .  $\square$

*Remark.*  $\hat{S}$  in the above Lemma cannot be replaced by  $S$ .

### 3.3 Strong Subadditivity

If the system under consideration enjoys the commuting square property with respect to a tracial state, the strong subadditivity property for the adjusted entropy  $\hat{S}$  holds (see Theorem 12 in [35]).

**Theorem 3.7.** *Let  $\mathcal{M}, \mathcal{N}_1, \mathcal{N}_2$  and  $\mathcal{P}$  be finite type  $\mathbf{I}$  factors satisfying one of the equivalent conditions of Proposition 2.4. Let  $\psi$  be a state on  $\mathcal{M}$ . Then*

$$\hat{S}(\psi) - \hat{S}(\psi_{\mathcal{N}_1}) - \hat{S}(\psi_{\mathcal{N}_2}) + \hat{S}(\psi_{\mathcal{P}}) \leq 0. \quad (3.8)$$

*Proof.* By (3.7) and (3.5)

$$\hat{S}_{\mathcal{N}_2}(\psi_{\mathcal{N}_2}) - \hat{S}_{\mathcal{M}}(\psi) = S_{\mathcal{M}}(\psi \circ E_{\mathcal{N}_2}^{\mathcal{M}}, \psi) \geq S_{\mathcal{M}}(\psi \circ E_{\mathcal{N}_2}^{\mathcal{M}} \circ E_{\mathcal{N}_1}^{\mathcal{M}}, \psi \circ E_{\mathcal{N}_1}^{\mathcal{M}})$$

By the assumption,  $E_{\mathcal{N}_2}^{\mathcal{M}} E_{\mathcal{N}_1}^{\mathcal{M}} = E_{\mathcal{N}_1}^{\mathcal{M}} E_{\mathcal{N}_2}^{\mathcal{M}} = E_{\mathcal{P}}^{\mathcal{M}}$ . Hence,

$$\begin{aligned} S_{\mathcal{M}}(\psi \circ E_{\mathcal{N}_2}^{\mathcal{M}} \circ E_{\mathcal{N}_1}^{\mathcal{M}}, \psi \circ E_{\mathcal{N}_1}^{\mathcal{M}}) &= S_{\mathcal{M}}(\psi_{\mathcal{P}} \otimes \tau_{\mathcal{M} \cap \mathcal{P}'}, \psi_{\mathcal{N}_1} \otimes \tau_{\mathcal{M} \cap \mathcal{N}_1'}) \\ &= S_{\mathcal{N}}(\psi_{\mathcal{P}} \otimes \tau_{\mathcal{N}_1 \cap \mathcal{P}'}, \psi_{\mathcal{N}_1}) \\ &= \hat{S}_{\mathcal{P}}(\psi_{\mathcal{P}}) - \hat{S}_{\mathcal{N}_1}(\psi_{\mathcal{N}_1}), \end{aligned}$$

where the second equality is due to  $\tau_{\mathcal{M} \cap \mathcal{P}'} = \tau_{\mathcal{N}_1 \cap \mathcal{P}'} \otimes \tau_{\mathcal{M} \cap \mathcal{N}_1'}$  and the last equality due to (3.7). Therefore we obtain (3.8).  $\square$



## 4 Fermion Lattice Systems

### 4.1 Fermion Algebra

We introduce Fermion lattice systems where there exists one spinless Fermion at each lattice site and they interact with each other. The restriction to spinless particle (i.e., one degree of freedom for each site) is just a matter of simplification of notation. All results and their proofs in the present work go over to the case of an arbitrary (constant) finite number of degrees of freedom at each lattice site without any essential alteration.

The lattice we consider is  $\nu$ -dimensional lattice  $\mathbb{Z}^\nu$  ( $\nu \in \mathbb{N}$ , an arbitrary positive integer).

**Definition 4.1.** *The Fermion  $\mathbf{C}^*$ -algebra  $\mathcal{A}$  is a unital  $\mathbf{C}^*$  algebra satisfying the following conditions and generated by elements in (1-1):*

- (1-1) *For each lattice site  $i \in \mathbb{Z}^\nu$ , there are elements  $a_i$  and  $a_i^*$  of  $\mathcal{A}$  called annihilation and creation operators, respectively, where  $a_i^*$  is the adjoint of  $a_i$ .*
- (1-2) *The CAR (canonical anticommutation relations) are satisfied for any  $i, j \in \mathbb{Z}^\nu$ :*

$$\begin{aligned} \{a_i^*, a_j\} &= \delta_{i,j} \mathbf{1} \\ \{a_i^*, a_j^*\} &= \{a_i, a_j\} = 0. \end{aligned} \quad (4.1)$$

Here  $\{A, B\} = AB + BA$  (anticommutator),  $\delta_{i,j} = 1$  for  $i = j$ , and  $\delta_{i,j} = 0$  for  $i \neq j$ .

- (1-3) *Let  $\mathcal{A}_0$  be the  $*$ -algebra generated by all  $a_i$  and  $a_i^*$  ( $i \in \mathbb{Z}^\nu$ ), namely the (algebraic) linear span of their monomials  $A_1 \cdots A_n$  where  $A_k$  is  $a_{i_k}$  or  $a_{i_k}^*$ ,  $i_k \in \mathbb{Z}^\nu$ .*

- (2) *For each subset  $I$  of  $\mathbb{Z}^\nu$ , the  $\mathbf{C}^*$ -subalgebra of  $\mathcal{A}$  generated by  $a_i, a_i^*, i \in I$ , is denoted by  $\mathcal{A}(I)$ . If the cardinality  $|I|$  of the set  $I$  is finite, then  $\mathcal{A}(I)$  is referred to as a local algebra or more specifically the local algebra for  $I$ . For the empty set  $\emptyset$ , we define  $\mathcal{A}(\emptyset) = \mathbb{C}\mathbf{1}$ .*

*Remark 1.*  $\mathcal{A}_0$  is dense in  $\mathcal{A}$ .

*Remark 2.* For finite  $I$ ,  $\mathcal{A}(I)$  is known to be isomorphic to the tensor product of  $|I|$  copies of the full  $2 \times 2$  matrix algebra  $M_2(\mathbb{C})$  and hence isomorphic to  $M_{2^{|I|}}(\mathbb{C})$ . Then

$$\mathcal{A}_0 = \bigcup_{|I| < \infty} \mathcal{A}(I)$$

has the unique  $\mathbf{C}^*$ -norm.  $\mathcal{A}$  together with its individual elements  $\{a_i, a_i^* | i \in \mathbb{Z}^\nu\}$  is uniquely defined up to isomorphism and is isomorphic to the UHF-algebra  $\overline{\bigotimes_{i \in \mathbb{Z}^\nu} M_2(\mathbb{C})}$ , where the bar denotes the norm completion.  $\mathcal{A}$  has the unique tracial state  $\tau$  as the extension of the unique tracial state of  $\mathcal{A}(I)$ ,  $|I| < \infty$ .

*Remark 3.* Since  $a_i^*$ 's and  $a_i$ 's anti-commute among different indices,  $a_i^*$  and  $a_i$  with a specific  $i$  can be brought together at any spot in a monomial, with a possible sign change (without changing the ordering among themselves), and this can be done for each  $i$ . Therefore, the monomials of the form

$$A_{i_1} \cdots A_{i_k} \quad (4.2)$$

together with  $\mathbf{1}$  have a dense linear span in  $\mathcal{A}(\mathbf{I})$ , where the indices  $i_1, \dots, i_k \in \mathbf{I}$  are distinct and  $A_{i_\alpha}$  is one of  $a_{i_\alpha}^*$ ,  $a_{i_\alpha}$ ,  $a_{i_\alpha}^* a_{i_\alpha}$ ,  $a_{i_\alpha} a_{i_\alpha}^*$ .

**Definition 4.2.**  $\Theta^{\mathbf{I}}$  denotes a unique automorphism of  $\mathcal{A}$  satisfying

$$\begin{aligned}\Theta^{\mathbf{I}}(a_i) &= -a_i, & \Theta^{\mathbf{I}}(a_i^*) &= -a_i^*, & (i \in \mathbf{I}), \\ \Theta^{\mathbf{I}}(a_i) &= a_i, & \Theta^{\mathbf{I}}(a_i^*) &= a_i^*, & (i \neq \mathbf{I}).\end{aligned}\tag{4.3}$$

In particular, we denote  $\Theta = \Theta^{\mathbb{Z}^\nu}$ .

The even and odd parts of  $\mathcal{A}$  are defined as

$$\mathcal{A}_+ \equiv \{a \in \mathcal{A} \mid \Theta(a) = a\}, \quad \mathcal{A}_- \equiv \{a \in \mathcal{A} \mid \Theta(a) = -a\}.\tag{4.4}$$

*Remark 1.* Such  $\Theta$  exists and is unique because (4.3) preserves CAR. It obviously satisfies

$$\Theta^2 = \text{id}.\tag{4.5}$$

*Remark 2.* For any  $a \in \mathcal{A}(\mathbf{I})$ ,

$$a = a_+ + a_-, \quad a_\pm \equiv \frac{1}{2}(a \pm \Theta(a))\tag{4.6}$$

gives the (unique) splitting of  $a$  into a sum of  $a_+ \in \mathcal{A}(\mathbf{I})_+$  and  $a_- \in \mathcal{A}(\mathbf{I})_-$ , where the even and odd parts of  $\mathcal{A}(\mathbf{I})$  are denoted by  $\mathcal{A}(\mathbf{I})_+$  and  $\mathcal{A}(\mathbf{I})_-$ .

*Remark 3.* For any  $a \in \mathcal{A}_-$ , we have

$$\tau(a) = \tau(\Theta(a)) = -\tau(a) = 0.\tag{4.7}$$

**Lemma 4.3.** Let  $\mathbf{I}$  and  $\mathbf{J}$  be mutually disjoint and  $a_\sigma \in \mathcal{A}(\mathbf{I})_\sigma$ ,  $b_\sigma \in \mathcal{A}(\mathbf{J})_\sigma$  where  $\sigma = +$  or  $-$ . Then

$$a_\sigma b_{\sigma'} = \epsilon(\sigma, \sigma') b_{\sigma'} a_\sigma,\tag{4.8}$$

where

$$\begin{aligned}\epsilon(\sigma, \sigma') &= -1 \quad \text{if } \sigma = \sigma' = - \\ &= +1 \quad \text{otherwise.}\end{aligned}$$

*Proof.* Since  $\mathcal{A}(\mathbf{I})$  is generated by  $a_i$  and  $a_i^*$ ,  $i \in \mathbf{I}$ , polynomials  $p$  of  $a_i$  and  $a_i^*$ ,  $i \in \mathbf{I}$ , are dense in  $\mathcal{A}(\mathbf{I})$ . For any  $\varepsilon > 0$  and a given  $a_\sigma$ ,  $\sigma = +$  or  $-$ , there exists a polynomial  $p$ , i.e. a linear combination  $p$  of monomials of  $a_i$  and  $a_i^*$ ,  $i \in \mathbf{I}$ , satisfying  $\|a_\sigma - p\| < \varepsilon$ . Since  $E_\sigma \equiv (1/2)(\text{id} + \sigma\Theta)$  satisfies  $E_\sigma a_\sigma = a_\sigma$  and  $\|E_\sigma\| = 1$ , we have

$$\|E_\sigma(a_\sigma - p)\| = \|a_\sigma - p_\sigma\| < \varepsilon$$

where  $p_\sigma = E_\sigma p$ . Since  $E_\sigma$  selects even or odd monomials (annihilating others) according as  $\sigma$  is  $+$  or  $-$ ,  $p_\sigma$  is a linear combination of even or odd monomials of  $a_i$  and  $a_i^*$ ,  $i \in \mathbf{I}$ . Similarly there exists a linear combination  $q_{\sigma'}$  of even or odd monomials of  $a_j$  and  $a_j^*$ ,  $j \in \mathbf{J}$ , satisfying  $\|b_{\sigma'} - q_{\sigma'}\| < \varepsilon$ . Since the graded commutation relation (4.8) holds for  $p_\sigma$  and  $q_{\sigma'}$ , it holds for  $a_\sigma$  and  $b_{\sigma'}$ .  $\square$

**Definition 4.4.** (1) For each  $k \in \mathbb{Z}^\nu$ ,  $\tau_k$  denotes a unique automorphism of  $\mathcal{A}$  satisfying

$$\tau_k(a_i^*) = a_{i+k}^*, \quad \tau_k(a_i) = a_{i+k}, \quad (i \in \mathbb{Z}^\nu). \quad (4.9)$$

(2) For a state  $\varphi$  of  $\mathcal{A}$ , the adjoint action of  $\tau_k$  is defined by

$$\tau_k^* \varphi(A) = \varphi(\tau_k(A)), \quad (A \in \mathcal{A}). \quad (4.10)$$

*Remark.* The automorphism  $\tau_k$  represents the lattice translation by the amount  $k \in \mathbb{Z}^\nu$ . The map  $k \in \mathbb{Z}^\nu \mapsto \tau_k$  is a group of automorphisms:

$$\tau_k \tau_l = \tau_{k+l}, \quad (k, l \in \mathbb{Z}^\nu).$$

The subalgebras transform covariantly under this group:

$$\tau_k(\mathcal{A}(I)) = \mathcal{A}(I + k), \quad (4.11)$$

where  $I + k = \{i + k; i \in I\}$  for any subset  $I$  of  $\mathbb{Z}^\nu$  and any  $k \in \mathbb{Z}^\nu$ .

**Definition 4.5.** The sets of all states and all positive linear functionals of  $\mathcal{A}$  are denoted by  $\mathcal{A}_{+,1}^*$  and  $\mathcal{A}_+^*$ ; the sets of all  $\Theta$  invariant and all  $\tau$  invariant ones by  $\mathcal{A}_{+,1}^{*\Theta}$ ,  $\mathcal{A}_+^{*\Theta}$  and  $\mathcal{A}_{+,1}^{*\tau}$ ,  $\mathcal{A}_+^{*\tau}$ , respectively. For any subset  $I$  of  $\mathbb{Z}^\nu$ , the set of all states of  $\mathcal{A}(I)$  is denoted by  $\mathcal{A}(I)_{+,1}^*$ ; the set of all  $\Theta$  invariant ones by  $\mathcal{A}(I)_{+,1}^{*\Theta}$ .

*Remark 1.* Any translation invariant state is automatically even (see, e.g., Example 5.2.21 of [17]):

$$\mathcal{A}_{+,1}^{*\tau} \subset \mathcal{A}_{+,1}^{*\Theta}. \quad (4.12)$$

*Remark 2.* For each subset  $I$  of  $\mathbb{Z}^\nu$ , we can consider the set of all states  $\{\mathcal{A}(I)_+\}^*_{+,1}$  on the even subalgebra  $\mathcal{A}(I)_+$ . There exists an obvious one-to-one correspondence between  $\mathcal{A}(I)_{+,1}^{*\Theta}$  and  $\{\mathcal{A}(I)_+\}^*_{+,1}$  due to (4.7) by the restriction and the unique  $\Theta$  invariant extension.

## 4.2 Product Property of the Tracial State

The following proposition provides a basis for the present section.

**Proposition 4.6.** If  $J_1$  and  $J_2$  are disjoint, then

$$\tau(ab) = \tau(a)\tau(b) \quad (4.13)$$

for arbitrary  $a \in \mathcal{A}(J_1)$  and  $b \in \mathcal{A}(J_2)$ ,

*Proof.* It is enough to prove the formula when  $a$  and  $b$  are monomials of the form (4.2). Let  $a = A_i a'$ , where  $i \in J_1$ ,  $a' \in \mathcal{A}(J_1 \setminus \{i\})$  is a monomial of the form (4.2) and  $A_i$  is one of  $a_i^*$ ,  $a_i$ ,  $a_i^* a_i$ ,  $a_i a_i^*$ . We will now show

$$\tau(ab) = \tau(A_i)\tau(a'b). \quad (4.14)$$

If  $a'b$  is a  $\Theta$ -odd monomial, then  $\tau(a'b) = 0$  by (4.7). If  $A_i$  is  $\Theta$ -even, then  $ab$  is odd and  $\tau(ab) = 0$ , implying (4.14). If  $A_i$  is odd, then  $A_i(a'b) = -(a'b)A_i$ . Hence

$$\begin{aligned}\tau(ab) &= \tau(A_i(a'b)) = -\tau((a'b)A_i) = -\tau(A_i(a'b)) \\ &= 0,\end{aligned}$$

where the third equality is due to the tracial property of  $\tau$ . So (4.14) holds in either case.

If  $a'b$  is even and  $A_i$  is odd, then  $\tau(A_i) = 0$  because  $A_i$  is odd and  $\tau(ab) = 0$  because  $ab = A_i(a'b)$  is odd. Again (4.14) holds.

Finally, if  $a'b$  is even and  $A_i = a_i^*a_i$ , then  $a_i^*$  commutes with  $a'b$  due to CAR and hence

$$\begin{aligned}\tau(ab) &= \tau((a_i^*a_i)(a'b)) = \tau(a_i(a'b)a_i^*) \\ &= \tau(a_i a_i^*(a'b)) \quad (\text{due to } [a_i^*, a'b] = 0) \\ &= \frac{1}{2}\tau((a_i^*a_i + a_i a_i^*)(a'b)) = \frac{1}{2}\tau(a'b).\end{aligned}$$

The same formula for  $a'b = 1$  yields  $\tau(A_i) = \frac{1}{2}$  and hence

$$\tau(ab) = \tau(A_i)\tau(a'b).$$

If  $a'b$  is even and  $A_i = a_i a_i^*$ , the above formula holds in the same way. We have now proved (4.14) for all cases.

Let  $a$  be now given by (4.2). By using (4.14) for  $i_1, i_2, \dots, i_k$  successively, we obtain

$$\tau(ab) = \tau(A_{i_1}) \cdots \tau(A_{i_k})\tau(b).$$

The same equality for  $b = 1$  yields

$$\tau(a) = \tau(A_{i_1}) \cdots \tau(A_{i_k}).$$

Hence we have

$$\tau(ab) = \tau(a)\tau(b).$$

This completes the proof.  $\square$

We may say that the tracial state  $\tau$  is a ‘product’ state although  $\mathcal{A}(J_1)$  and  $\mathcal{A}(J_2)$  do not commute. We will show in the next subsections that this product property of the tracial state implies the commuting square property for the conditional expectations.

### 4.3 Conditional Expectations for Fermion Algebras

We prove the  $\mathbf{C}^*$ -algebraic version of Proposition 2.1 for the Fermion algebra  $\mathcal{A}$  and its subalgebras. We note that  $\mathcal{A}(I)$  is not a von Neumann algebra unless  $I$  is a finite subset of  $\mathbb{Z}^\nu$ . Hence Proposition 2.1 is not directly applicable to the Fermion algebra.

**Theorem 4.7.** *For any subset  $I$  of  $\mathbb{Z}^\nu$ , there exists a conditional expectation*

$$E_I : a \in \mathcal{A} \mapsto E_I(a) \in \mathcal{A}(I) \quad (4.15)$$

*uniquely determined by  $E_I(a) \in \mathcal{A}(I)$  and*

$$\tau(ab) = \tau(E_I(a)b) \quad (b \in \mathcal{A}(I)). \quad (4.16)$$

*For any second subset  $J$  of  $\mathbb{Z}^\nu$ ,*

$$E_I(a) \in \mathcal{A}(I \cap J) \quad (4.17)$$

*for any  $a \in \mathcal{A}(J)$ , and*

$$E_I E_J = E_J E_I = E_{I \cap J}. \quad (4.18)$$

*Proof.* The  $\mathbf{C}^*$ -subalgebra of  $\mathcal{A}$  generated by  $\mathcal{A}(I)$  and  $\mathcal{A}(I^c)_+$  is isomorphic to their tensor product and will be denoted as  $\mathcal{A}(I) \otimes \mathcal{A}(I^c)_+$ . Let

$$E_I^{(1)} \equiv \frac{1}{2}(\text{id} + \Theta^{I^c}). \quad (4.19)$$

It maps  $\mathcal{A}$  onto  $\mathcal{A}(I) \otimes \mathcal{A}(I^c)_+$ . Since

$$\tau(\Theta^{I^c}(a)b) = \tau(\Theta^{I^c}(ab)) = \tau(ab)$$

for all  $a \in \mathcal{A}$  and  $b \in \mathcal{A}(I) \otimes \mathcal{A}(I^c)_+$ ,  $E_I^{(1)}$  satisfies (4.16).

Since  $\tau$  is a product state for the tensor product  $\mathcal{A}(I) \otimes \mathcal{A}(I^c)_+$ , there exists a conditional expectation  $E_I^{(2)}$  from  $\mathcal{A}(I) \otimes \mathcal{A}(I^c)_+$  onto  $\mathcal{A}(I)$  satisfying (4.16), characterized by  $E_I^{(2)}(cd) = \tau(d)c$  for  $c \in \mathcal{A}(I)$  and  $d \in \mathcal{A}(I^c)_+$  and called a slice map. Therefore

$$E_I = E_I^{(2)} E_I^{(1)} \quad (4.20)$$

is a map from  $\mathcal{A}$  onto  $\mathcal{A}(I)$  satisfying (4.16). By Lemma 2.2, it is a unique conditional expectation from  $\mathcal{A}$  onto  $\mathcal{A}(I)$  satisfying (4.16).

To show (4.17), note that  $\mathcal{A}(J)$  is generated by  $\mathcal{A}(J \cap I)$  and  $\mathcal{A}(J \cap I^c)$ , namely, the linear span of products  $ab$  with  $a \in \mathcal{A}(J \cap I)$  and  $b \in \mathcal{A}(J \cap I^c)$  is dense in  $\mathcal{A}(J)$ . Due to the linearity of  $E_I$  and  $\|E_I\| = 1$ , it is enough to show (4.17) for such products. We have  $E_I^{(1)}(b) \in \mathcal{A}(I^c)_+$  and hence

$$E_I(ab) = E_I^{(2)}(aE_I^{(1)}(b)) = a\tau(E_I^{(1)}(b)) = a\tau(b) \in \mathcal{A}(J \cap I),$$

which proves (4.17).

For any  $a \in \mathcal{A}$ ,  $E_J(a) \in \mathcal{A}(J)$  and hence  $E_I(E_J(a)) \in \mathcal{A}(I \cap J)$ . For  $b \in \mathcal{A}(I \cap J)$ , (4.16) implies

$$\tau(E_I(E_J(a))b) = \tau(E_J(a)b) = \tau(ab),$$

where the first equality is due to  $b \in \mathcal{A}(I)$ , while the second equality is due to  $b \in \mathcal{A}(J)$ . This equality and  $E_I(E_J(a)) \in \mathcal{A}(I \cap J)$  imply

$$E_{I \cap J}(a) = E_I(E_J(a))$$

by the uniqueness result. By interchanging I and J, we obtain

$$E_I E_J = E_J E_I = E_{I \cap J},$$

which proves the last statement (4.18).  $\square$

*Remark 1.* For spin lattice systems, the conditional expectation  $E_I$  can be obtained simply as a slice map with respect to the tracial state  $\tau$ . When spins and Fermions coexist at each lattice site,  $E_I$  can be obtained in exactly the same way as Theorem 4.7 (by including spin operators in the even part  $\mathcal{A}(I)_+$ ), provided that the degree of freedom at each lattice site is finite (i.e.  $\mathcal{A}(I)$  is a finite factor of type I for any finite I). In all these cases, the results of our paper are valid as they are proved by the use of conditional expectations  $E_I$ .

*Remark 2.* Theorem 4.7 can be shown by a more elementary (lengthy) method by giving  $E_I$  explicitly for a finite I and then giving  $E_J$  for an infinite J as a limit of  $E_{I_n}$  for an increasing sequence of finite subsets  $I_n$  of  $\mathbb{Z}^\nu$  tending to J. Proof presented above is by a suggestion of a referee.

**Corollary 4.8.** *For each subset I of  $\mathbb{Z}^\nu$ ,*

$$E_I \Theta = \Theta E_I. \quad (4.21)$$

*Proof.* For any  $a \in \mathcal{A}$  and  $b \in \mathcal{A}(I)$ ,

$$\begin{aligned} \tau(E_I(\Theta(a))b) &= \tau(\Theta(a)b) \\ &= \tau(\Theta\{\Theta(a)b\}) = \tau(a\Theta(b)) \\ &= \tau(E_I(a)\Theta(b)) = \tau(\Theta\{E_I(a)\Theta(b)\}) \\ &= \tau(\Theta(E_I(a))b). \end{aligned}$$

Since  $\mathcal{A}(I)$  is invariant under  $\Theta$  as a set, we have  $\Theta(E_I(a)) = E_I(\Theta(a))$  due to the uniqueness of  $E_I$  in the preceding theorem.  $\square$

We now show a continuous dependence of  $E_I$  on the subsets I of  $\mathbb{Z}^\nu$ . We use the following notation for various limits of subsets of  $\mathbb{Z}^\nu$ . If  $\{I_\alpha\}$  is a monotone (not necessarily strictly) increasing or decreasing net of subsets converging to a subset I of  $\mathbb{Z}^\nu$ , we write  $I_\alpha \nearrow I$  or  $I_\alpha \searrow I$ . For these cases,  $I = \cup_\alpha I_\alpha$  or  $I = \cap_\alpha I_\alpha$ , respectively. We use  $I_\alpha \rightarrow I$  for the standard convergence of a net  $I_\alpha$  to I (i.e.,  $\limsup_\alpha I_\alpha = \liminf_\alpha I_\alpha = I$ ). By  $J \nearrow \mathbb{Z}^\nu$  (which is written without any index), we mean a net of all finite subsets tending to  $\mathbb{Z}^\nu$  with the set inclusion as its partial ordering. (In the same way, we use  $J \nearrow I$ .) In this case, J itself serves as the net index and it is a monotone increasing net. Later in § 9 and § 10, we use a more restrictive notion of a van Hove net  $\{I_\alpha\}$  tending to  $\mathbb{Z}^\nu$  or to ‘ $\infty$ ’ (see Appendix for detailed explanation).

**Lemma 4.9.** *Let  $\{I_\alpha\}$  be an increasing net of (finite or infinite) subsets of I such that their union is I. For any  $a \in \mathcal{A}$ ,*

$$\lim_\alpha E_{I_\alpha}(a) = E_I(a). \quad (4.22)$$

*As a special case  $I = \mathbb{Z}^\nu$ ,*

$$\lim_{I_\alpha \nearrow \mathbb{Z}^\nu} E_{I_\alpha}(a) = a. \quad (4.23)$$

*Proof.* Since polynomials of  $a_i$  and  $a_i^*$ ,  $i \in I$ , are dense in  $\mathcal{A}(I)$ , there exists a finite subset  $J_n$  of  $I$  and  $a_n \in \mathcal{A}(J_n)$  such that

$$\|E_I(a) - a_n\| < \frac{1}{n}.$$

Because  $J_n$  is a finite subset of  $I$  and  $\cup_\alpha I_\alpha = I$ , there exists a finite number of  $I_\alpha$ , say,  $I_{\alpha(1)}, \dots, I_{\alpha(k)}$ , such that  $\cup_{l=1}^k I_{\alpha(l)} \supset J_n$ . Since  $I_\alpha$  is a net, there exists an index  $\alpha_n > \alpha(1), \dots, \alpha(k)$ . Since  $I_\alpha$  is increasing,  $I_{\alpha_n} \supset I_{\alpha(1)} \cup \dots \cup I_{\alpha(k)} \supset J_n$ .

For any  $\alpha \geq \alpha_n$ ,  $I_\alpha \supset J_n$  and so  $E_{I_\alpha}(a_n) = a_n$ . Hence by  $I \supset I_\alpha$ , we have

$$\|E_{I_\alpha}(a) - a_n\| = \|E_{I_\alpha}(E_I(a) - a_n)\| \leq \|E_I(a) - a_n\| < \frac{1}{n}$$

due to  $\|E_{I_\alpha}\| \leq 1$ . Thus

$$\|E_{I_\alpha}(a) - E_I(a)\| \leq \|E_{I_\alpha}(a) - a_n\| + \|E_I(a) - a_n\| < \frac{2}{n},$$

for all  $\alpha \geq \alpha_n$ , which proves the assertion (4.22).  $\square$

**Lemma 4.10.** *Let  $\{I_\alpha\}$  be a decreasing net of (finite or infinite) subsets of  $\mathbb{Z}^\nu$  such that their intersection is  $I$ . For any  $a \in \mathcal{A}$ ,*

$$\lim_\alpha E_{I_\alpha}(a) = E_I(a). \quad (4.24)$$

*Proof.* Let  $L_k$  be a monotone increasing sequence of finite subsets of  $\mathbb{Z}^\nu$  such that their union is  $\mathbb{Z}^\nu$ . For any  $\varepsilon > 0$ , there exists  $k_\varepsilon$  such that

$$\|a - E_{L_k}(a)\| < \varepsilon$$

for all  $k \geq k_\varepsilon$  by Lemma 4.9. Hence

$$\|E_I(a) - E_{I \cap L_k}(a)\| = \|E_I(a - E_{L_k}(a))\| < \varepsilon, \quad (4.25)$$

$$\|E_{I_\alpha}(a) - E_{I_\alpha \cap L_k}(a)\| = \|E_{I_\alpha}(a - E_{L_k}(a))\| < \varepsilon \quad (4.26)$$

for all  $k \geq k_\varepsilon$  and all  $\alpha$  due to  $\|E_I\| \leq 1$  and  $\|E_{I_\alpha}\| \leq 1$ .

Since  $I_\alpha \searrow I$ , we have  $(I_\alpha \cap L_k) \searrow (I \cap L_k)$ . Since  $L_{k_\varepsilon}$  is a finite set, there exists  $\alpha_\varepsilon$  such that  $I_\alpha \cap L_{k_\varepsilon} = I \cap L_{k_\varepsilon}$  and hence  $E_{I_\alpha \cap L_{k_\varepsilon}} = E_{I \cap L_{k_\varepsilon}}$  for all  $\alpha \geq \alpha_\varepsilon$ . Therefore, we obtain

$$\begin{aligned} \|E_{I_\alpha}(a) - E_I(a)\| &\leq \|E_{I_\alpha}(a) - E_{I_\alpha \cap L_{k_\varepsilon}}(a)\| + \|E_{I_\alpha \cap L_{k_\varepsilon}}(a) - E_I(a)\| \\ &= \|E_{I_\alpha}(a) - E_{I_\alpha \cap L_{k_\varepsilon}}(a)\| + \|E_{I \cap L_{k_\varepsilon}}(a) - E_I(a)\| < 2\varepsilon \end{aligned}$$

for all  $\alpha \geq \alpha_\varepsilon$ , where the first term is estimated by (4.26), and the second by (4.25). Hence we obtain

$$\lim_\alpha E_{I_\alpha}(a) = E_I(a).$$

$\square$

**Theorem 4.11.** *If a net  $\{I_\alpha\}$  converges to  $I$ , then*

$$\lim_{\alpha} E_{I_\alpha}(a) = E_I(a). \quad (4.27)$$

for all  $a \in \mathcal{A}$ .

*Proof.* By definition,  $I_\alpha \rightarrow I$  means

$$I = \cap_{\beta} (\cup_{\alpha \geq \beta} I_\alpha) = \cup_{\beta} (\cap_{\alpha \geq \beta} I_\alpha).$$

Set

$$J^\beta \equiv \cup_{\alpha \geq \beta} I_\alpha, \quad J_\beta \equiv \cap_{\alpha \geq \beta} I_\alpha.$$

Then  $J^\beta \searrow I$  and  $J_\beta \nearrow I$ . By Lemmas 4.10 and 4.9, there exists a  $\beta_\varepsilon$  for any given  $\varepsilon > 0$  such that for all  $\beta \geq \beta_\varepsilon$

$$\|E_{J^\beta}(a) - E_I(a)\| < \varepsilon, \quad \|E_{J_\beta}(a) - E_I(a)\| < \varepsilon.$$

Hence

$$\|E_{J^\beta}(a) - E_{J_\beta}(a)\| < 2\varepsilon.$$

Since  $J^\beta \supset I_\beta \supset J_\beta$ , we have  $E_{I_\beta} E_{J^\beta} = E_{I_\beta}$ ,  $E_{I_\beta} E_{J_\beta} = E_{J_\beta}$  and

$$\|E_{I_\beta}(a) - E_{J_\beta}(a)\| = \|E_{I_\beta}(E_{J^\beta}(a) - E_{J_\beta}(a))\| < 2\varepsilon.$$

Therefore

$$\|E_{I_\beta}(a) - E_I(a)\| < 3\varepsilon$$

for all  $\beta \geq \beta_\varepsilon$ . This proves (4.27).  $\square$

The following corollary follows immediately from the results obtained in this subsection.

**Corollary 4.12.** *For any countable family  $\{I_n\}$  of subsets of  $\mathbb{Z}^\nu$ ,*

$$\cap_{n=1}^{\infty} \mathcal{A}(I_n) = \mathcal{A}(\cap_{n=1}^{\infty} I_n). \quad (4.28)$$

*Proof.* Let  $J_n \equiv \cap_{k=1}^n I_k$  and  $I \equiv \cap_{n=1}^{\infty} I_n$ . Then  $J_n \searrow I$ . By (4.18),  $E_{J_{n-1}} E_{I_n} = E_{J_n}$  and hence  $E_{J_n} = \prod_{k=1}^n E_{I_k}$ . On one hand,  $J_n \subset I_k$  for  $k = 1, \dots, n$ , and hence  $\mathcal{A}(J_n) \subset \cap_{k=1}^n \mathcal{A}(I_k)$ . On the other hand,  $a \in \cap_{k=1}^n \mathcal{A}(I_k)$  satisfies  $E_{I_k}(a) = a$  for all  $k = 1, \dots, n$  and hence  $E_{J_n}(a) = a \in \mathcal{A}(J_n)$ . Therefore

$$\mathcal{A}(J_n) = \cap_{k=1}^n \mathcal{A}(I_k).$$

Since  $J_n \supset I$ , we have  $\mathcal{A}(J_n) \supset \mathcal{A}(I)$  and hence

$$\cap_{n=1}^{\infty} \mathcal{A}(I_n) = \cap_{n=1}^{\infty} \mathcal{A}(J_n) \supset \mathcal{A}(I).$$

For  $a \in \cap_{n=1}^{\infty} \mathcal{A}(J_n)$ ,  $E_{J_n}(a) = a$  for any  $n$ . Since  $\lim_n E_{J_n}(a) = E_I(a)$  by Lemma 4.10, we have  $a = E_I(a) \in \mathcal{A}(I)$ . Now we obtain the desired conclusion

$$\cap_{n=1}^{\infty} \mathcal{A}(I_n) = \mathcal{A}(I).$$


$\square$



#### 4.4 Commuting Squares for Fermion Algebras

In the following theorem, we show that any two subsets  $I$  and  $J$  of  $\mathbb{Z}^\nu$  are associated with a commuting square of the conditional expectations with respect to the tracial state  $\tau$ . For  $K \subset L \subset \mathbb{Z}^\nu$ , denote the restriction of  $E_K$  to  $\mathcal{A}(L)$  by  $E_K^L$ . Then it is a conditional expectation from  $\mathcal{A}(L)$  to  $\mathcal{A}(K)$  with respect to the tracial state.

**Theorem 4.13.** *For any subsets  $I$  and  $J$  of  $\mathbb{Z}^\nu$ , the following subalgebras of  $\mathcal{A}$  form a commuting square:*



Here the arrow from  $\mathcal{A}(L)$  to  $\mathcal{A}(K)$  represents the conditional expectation  $E_K^L$ .

*Proof.* It follows from (4.18) that

$$E_{I \cap J}^I E_I^{I \cup J} = E_{I \cap J}^{I \cup J} = E_{I \cap J}^J E_J^{I \cup J},$$

which shows the assertion.  $\square$

#### 4.5 Commutants of Subalgebras

We are going to determine the commutants of subalgebras of  $\mathcal{A}$ .

**Lemma 4.14.** *For a finite  $I$ ,*

$$(\mathcal{A}(I)_+)' \cap \mathcal{A} = \mathcal{A}(I^c) + v_I \mathcal{A}(I^c), \quad (4.29)$$

where  $v_I$  is a self-adjoint unitary in  $\mathcal{A}(I)_+$  given by

$$v_I \equiv \prod_{i \in I} v_i, \quad v_i \equiv a_i^* a_i - a_i a_i^*. \quad (4.30)$$

and implementing  $\Theta^I$  on  $\mathcal{A}$ .

*Proof.* By CAR,

$$a_i^* v_i = -a_i^*, \quad a_i v_i = a_i, \quad v_i a_i^* = a_i^*, \quad v_i a_i = -a_i.$$

Thus  $v_i$  anticommutes with  $a_i$  and  $a_i^*$ . If  $j \neq i$ ,  $v_i$  commutes with  $a_j$  and  $a_j^*$  due to  $v_i \in \mathcal{A}(\{i\})_+$ . Therefore for any  $a \in \mathcal{A}(I)$ , we have

$$(\text{Ad } v_I) a \equiv v_I a v_I^* = \Theta(a), \quad (4.31)$$

or equivalently,

$$v_I a = \Theta(a) v_I. \quad (4.32)$$

For any  $a \in \mathcal{A}(\mathbf{I}^c)$ ,

$$v_{\mathbf{I}}a = av_{\mathbf{I}}. \quad (4.33)$$

Due to  $v_{\mathbf{I}}^* = v_{\mathbf{I}} = v_{\mathbf{I}}^2$ ,  $v_{\mathbf{I}}$  is a self-adjoint unitary implementing  $\Theta^{\mathbf{I}}$  on  $\mathcal{A}$ .

Since  $v_{\mathbf{I}} \in \mathcal{A}(\mathbf{I})_+$  implements  $\Theta^{\mathbf{I}}$ ,  $(\mathcal{A}(\mathbf{I})_+)'$  is contained in the fixed point subalgebra  $\mathcal{A}^{\Theta^{\mathbf{I}}}$ . In terms of  $E_{\mathbf{I}^c}^{(1)} = \frac{1}{2}(\text{id} + \Theta^{\mathbf{I}})$ , we have

$$(\mathcal{A}(\mathbf{I})_+)' \subset \mathcal{A}^{\Theta^{\mathbf{I}}} = E_{\mathbf{I}^c}^{(1)}(\mathcal{A}) = \mathcal{A}(\mathbf{I})_+ \otimes \mathcal{A}(\mathbf{I}^c).$$

Since  $\mathcal{A}(\mathbf{I}^c)$  is in  $(\mathcal{A}(\mathbf{I})_+)'$ , we have

$$(\mathcal{A}(\mathbf{I})_+)' = \mathcal{Z}(\mathcal{A}(\mathbf{I})_+) \otimes \mathcal{A}(\mathbf{I}^c) \quad (4.34)$$

where  $\mathcal{Z}(\mathcal{A}(\mathbf{I})_+)$  is the center of  $\mathcal{A}(\mathbf{I})_+$ . Since  $\mathcal{A}(\mathbf{I})_+ = \{v_{\mathbf{I}}\}' \cap \mathcal{A}(\mathbf{I})$ ,  $v_{\mathbf{I}}$  is a self-adjoint unitary in  $\mathcal{A}(\mathbf{I})$  and  $\mathcal{A}(\mathbf{I})$  is a full matrix algebra for a finite  $\mathbf{I}$ , we have

$$\mathcal{Z}(\mathcal{A}(\mathbf{I})_+) = \mathbb{C}\mathbf{1} + \mathbb{C}v_{\mathbf{I}}. \quad (4.35)$$

By (4.34) and (4.35), we obtain (4.29).  $\square$

**Lemma 4.15.** *For a finite  $\mathbf{I}$ ,*

$$\mathcal{A}(\mathbf{I})' \cap \mathcal{A} = \mathcal{A}(\mathbf{I}^c)_+ + v_{\mathbf{I}}\mathcal{A}(\mathbf{I}^c)_-, \quad (4.36)$$

*Proof.* By Lemma 4.14 and  $\mathcal{A}(\mathbf{I})' \subset (\mathcal{A}(\mathbf{I})_+)'$ , any element  $a \in \mathcal{A}(\mathbf{I})'$  is of the form

$$a = a_1 + v_{\mathbf{I}}a_2, \quad a_1, a_2 \in \mathcal{A}(\mathbf{I}^c).$$

Take any unitary  $u \in \mathcal{A}(\mathbf{I})_-$  (e.g.,  $u = a_i + a_i^*$ ,  $i \in \mathbf{I}$ ). Then we have

$$\begin{aligned} a &= \frac{1}{2}(a + uau^*) = \frac{1}{2}(a_1 + ua_1u^*) + \frac{1}{2}v_{\mathbf{I}}(a_2 - ua_2u^*) \\ &= (a_1)_+ + v_{\mathbf{I}}(a_2)_- \end{aligned}$$

due to  $uv_{\mathbf{I}} = -v_{\mathbf{I}}u$ , where

$$(a_1)_+ = \frac{1}{2}(a_1 + \Theta(a_1)) \in \mathcal{A}(\mathbf{I}^c)_+, \quad (a_2)_- = \frac{1}{2}(a_2 - \Theta(a_2)) \in \mathcal{A}(\mathbf{I}^c)_-.$$

Hence

$$\mathcal{A}(\mathbf{I})' \subset \mathcal{A}(\mathbf{I}^c)_+ + v_{\mathbf{I}}\mathcal{A}(\mathbf{I}^c)_-.$$

The inverse inclusion follows from (4.32) and Lemma 4.3. Hence (4.36) holds.  $\square$

**Lemma 4.16.** *For an infinite  $\mathbf{I}$ ,*

$$\mathcal{A}(\mathbf{I})' \cap \mathcal{A} = \mathcal{A}(\mathbf{I}^c)_+. \quad (4.37)$$

*Proof.* It is clear that elements of  $\mathcal{A}(\mathcal{I}^c)_+$  and  $\mathcal{A}(\mathcal{I})$  commute. Hence it is enough to prove  $\mathcal{A}(\mathcal{I})' \cap \mathcal{A} \subset \mathcal{A}(\mathcal{I}^c)_+$ .

Let  $a \in \mathcal{A}(\mathcal{I})' \cap \mathcal{A}$ . Then

$$a_{\pm} = \frac{1}{2}(a \pm \Theta(a)) \in \mathcal{A}(\mathcal{I})' \cap \mathcal{A}$$

because  $\Theta(\mathcal{A}(\mathcal{I})) = \mathcal{A}(\mathcal{I})$ . For any finite subset  $K$  of  $\mathcal{I}$ ,  $a_{\pm} \in (\mathcal{A}(K))'_{\pm}$ . Hence by Lemma 4.15,

$$a_+ \in \mathcal{A}(K^c)_+.$$

Consider an increasing sequence of finite subsets  $K_n \nearrow \mathcal{I}$ . We apply Corollary 4.12 to  $(K_n)^c \searrow \mathcal{I}^c$ , and obtain

$$a_+ \in \bigcap_{n=1}^{\infty} \mathcal{A}((K_n)^c)_+ = \mathcal{A}(\mathcal{I}^c)_+. \quad (4.38)$$

We now prove  $a_- = 0$ , which yields the desired conclusion due to  $a = a_+ + a_-$  and (4.38). For a monotone increasing sequence of finite subsets  $L_n$  of  $\mathbb{Z}^\nu$  such that  $L_n \nearrow \mathbb{Z}^\nu$ , we have  $\lim_n E_{L_n}(a_-) = a_-$  and hence there exists  $n_\varepsilon$  for any given  $\varepsilon > 0$  such that

$$\|E_{L_n}(a_-) - a_-\| < \varepsilon \quad (4.39)$$

for  $n \geq n_\varepsilon$ . For any  $k$ , we set  $K_k \equiv \mathcal{I} \cap L_k (\subset \mathcal{I})$ . Then  $a_- \in \mathcal{A}(K_k)'$  and by Lemma 4.15 we have

$$a_- = v_{K_k} b_k$$

for some  $b_k \in \mathcal{A}((K_k)^c)_-$ . For any  $i \in K_k$ ,

$$E_{\{i\}^c}(a_-) = \tau(v_i) v_{(K_k \setminus \{i\})} b_k = 0. \quad (4.40)$$

Now take an  $n_0 \geq n_\varepsilon$ . Since  $K_k \nearrow \mathcal{I}$  and  $\mathcal{I}$  is an infinite set while any  $L_{n_0}$  is a finite set, there exists a number  $k$  such that  $K_k$  contains a point  $i$  of  $\mathbb{Z}^\nu$  such that  $i \notin L_{n_0}$ . Then  $L_{n_0} \subset \{i\}^c$ . It follows from (4.40) that

$$E_{L_{n_0}}(a_-) = E_{L_{n_0}} E_{\{i\}^c}(a_-) = 0.$$

This and (4.39) imply

$$\|a_-\| < \varepsilon.$$

Since  $\varepsilon$  is arbitrary, we obtain  $a_- = 0$ . □

Combining Lemma 4.15 and Lemma 4.16, we obtain

**Theorem 4.17.** (1) For a finite  $\mathcal{I}$ ,

$$\mathcal{A}(\mathcal{I})' \cap \mathcal{A} = \mathcal{A}(\mathcal{I}^c)_+ + v_{\mathcal{I}} \mathcal{A}(\mathcal{I}^c)_-,$$

where  $v_{\mathcal{I}}$  is given by (4.30).

(2) For an infinite  $\mathcal{I}$ ,

$$\mathcal{A}(\mathcal{I})' \cap \mathcal{A} = \mathcal{A}(\mathcal{I}^c)_+.$$

As a preparation for the remaining case (the commutant of  $\mathcal{A}(\mathbf{I})_+$  for infinite  $\mathbf{I}$ ), we present the following technical Lemma for the sake of completeness. We define

$$u_{11}^{(i)} \equiv a_i^* a_i, \quad u_{12}^{(i)} \equiv a_i^*, \quad u_{21}^{(i)} \equiv a_i, \quad u_{22}^{(i)} \equiv a_i a_i^*. \quad (4.41)$$

**Lemma 4.18.** *Let  $\mathbf{I} = (i_1, \dots, i_{|\mathbf{I}|})$  be a finite subset of  $\mathbb{Z}^\nu$ . Put*

$$u_{\alpha\alpha}^{(i_j)} \equiv u_{\alpha\alpha}^{(i_j)} \text{ for } \alpha = 1, 2, \quad u_{\alpha\beta}^{(i_j)} \equiv u_{\alpha\beta}^{(i_j)} v_{\{i_1, \dots, i_{j-1}\}} \text{ for } \alpha \neq \beta. \quad (4.42)$$

Define

$$u_{kl} \equiv \prod_{j=1}^{|\mathbf{I}|} u_{k_j l_j}^{(i_j)}, \quad (4.43)$$

where  $k_n$  and  $l_n$  are either 1 or 2, respectively,  $k = (k_1, \dots, k_{|\mathbf{I}|})$  and  $l = (l_1, \dots, l_{|\mathbf{I}|})$ . Then the following holds.

- (1) The set of all  $u_{kl}$  form a self-adjoint system of matrix units of  $\mathcal{A}(\mathbf{I})$ .
- (2) Let  $\sigma(k, l)$  be the number of  $n$  such that  $k_n \neq l_n$ . Then

$$\Theta(u_{kl}) = (-1)^{\sigma(k, l)} u_{kl}. \quad (4.44)$$

- (3) Any  $a \in \mathcal{A}$  has a unique expansion

$$a = \sum_{k, l} u_{kl} a_{kl} \quad (4.45)$$

with  $a_{kl} \in \mathcal{A}(\mathbf{I}^c)$  and  $a_{kl}$  is uniquely given by

$$a_{kl} = 2^{|\mathbf{I}|} E_{\mathbf{I}^c}(u_{lk} a). \quad (4.46)$$

*Proof.* (1) By using (4.1) for the case of  $i = j$ ,  $\{u_{\alpha\beta}^{(i)}\}_{\alpha\beta} (\alpha, \beta = 1, 2)$  satisfies the relations

$$(u_{\alpha\beta}^{(i)})^* = u_{\beta\alpha}^{(i)}, \quad u_{\alpha\beta}^{(i)} u_{\alpha'\beta'}^{(i)} = \delta_{\beta\alpha'} u_{\alpha\beta'}^{(i)}, \quad \sum_{\alpha} u_{\alpha\alpha}^{(i)} = 1, \quad (4.47)$$

for a self-adjoint system of matrix units. Since  $v_{\{i_1, \dots, i_{j-1}\}}$  is a self-adjoint unitary commuting with  $a_{i_j}$  and  $a_{i_j}^*$ , the same computation shows that  $\{u_{\alpha\beta}^{(i_j)}\}_{\alpha\beta} (\alpha, \beta = 1, 2)$  satisfies the same relations.

Since  $v_{\{i_1, \dots, i_{j-1}\}}$  anticommutes with  $a_{i_k}$  and  $a_{i_k}^*$  for  $k < j$  and commutes with them for  $k \geq j$ ,  $\{u_{\alpha\beta}^{(i_j)}\}_{\alpha\beta}$  commutes with each other for different  $j$ .

Since they generate all  $\mathcal{A}(\{i_k\})$  recursively for  $k = 1, \dots, n$ , they form a self-adjoint system of matrix units of  $\mathcal{A}(\mathbf{I})$ .

- (2)  $\Theta(u_{\alpha\alpha}^{(i)}) = u_{\alpha\alpha}^{(i)}$ ,  $\Theta(u_{\alpha\beta}^{(i)}) = -u_{\alpha\beta}^{(i)}$  for  $\alpha \neq \beta$ , and  $\Theta(v_{\{i_1, \dots, i_{j-1}\}}) = v_{\{i_1, \dots, i_{j-1}\}}$  imply (4.44).

(3) For a full matrix algebra  $\mathcal{A}(\mathbf{I})$  contained in a  $\mathbf{C}^*$ -algebra  $\mathcal{A}$ , the following expansion of any  $a \in \mathcal{A}$  in term of a self-adjoint system of a matrix units  $\{u_{kl}\}$  of  $\mathcal{A}(\mathbf{I})$  is well-known.

$$\begin{aligned} a &= \sum_{k, l} u_{kl} b_{kl}, \\ b_{kl} &= \sum_m u_{mk} a u_{lm} \in \mathcal{A}(\mathbf{I})'. \end{aligned} \quad (4.48)$$

By Lemma 4.15, there are  $b_{kl1}$  and  $b_{kl2}$  in  $\mathcal{A}(\Gamma^c)$  satisfying

$$b_{kl} = b_{kl1} + v_I b_{kl2}. \quad (4.49)$$

By direct computation,  $u_{kl}v_I = \pm u_{kl}$  where the sign depends on  $k$  and  $l$ . Thus we have the expansion (4.45) with  $a_{kl} = b_{kl1} \pm b_{kl2} \in \mathcal{A}(\Gamma^c)$ .

The coefficient  $a_{kl} \in \mathcal{A}(\Gamma^c)$  is uniquely determined by the following computation and given by (4.46).

$$\begin{aligned} E_{\Gamma^c}(u_{lk}a) &= E_{\Gamma^c}\left(\sum_{l'} u_{ll'} a_{kl'}\right) \\ &= \sum_{l'} E_{\Gamma^c}(u_{ll'}) a_{kl'} = \sum_{l'} \tau(u_{ll'}) a_{kl'} = 2^{-|\Gamma|} a_{kl}. \end{aligned}$$

Here we have used the following relation:

$$\begin{aligned} \tau(u_{kl}) &= \tau(u_{km}u_{ml}) = \tau(u_{ml}u_{km}) \\ &= \delta_{kl}\tau(u_{mm}) = \delta_{kl}2^{-|\Gamma|}\tau\left(\sum_m u_{mm}\right) = 2^{-|\Gamma|}\delta_{kl}. \end{aligned}$$

□

**Theorem 4.19.** (1) For a finite  $I$ ,

$$(\mathcal{A}(I)_+)' \cap \mathcal{A} = \mathcal{A}(\Gamma^c) + v_I \mathcal{A}(\Gamma^c), \quad (4.50)$$

where  $v_I$  is given by (4.30).

(2) For an infinite  $I$ ,

$$(\mathcal{A}(I)_+)' \cap \mathcal{A} = \mathcal{A}(\Gamma^c). \quad (4.51)$$

*Proof.* (1) is given by Lemma 4.14.

To prove (2), we consider an infinite  $I$ . Clearly  $(\mathcal{A}(I)_+)' \cap \mathcal{A} \supset \mathcal{A}(\Gamma^c)$  due to (4.8). Hence it is enough to prove that any  $b \in (\mathcal{A}(I)_+)' \cap \mathcal{A}$  belongs to  $\mathcal{A}(\Gamma^c)$ .

Let  $\{L_n\}$  be an increasing sequence of finite subsets of  $\mathbb{Z}^\nu$  such that their union is  $\mathbb{Z}^\nu$ . Set  $I_n \equiv L_n \cap I$ . Then  $I_n \nearrow I$ .

For any  $\varepsilon > 0$ , there exist a positive integer  $l_\varepsilon$  and an element  $b_\varepsilon$  of  $\mathcal{A}(L_{l_\varepsilon})$  satisfying

$$\|b - b_\varepsilon\| < \varepsilon.$$

For any  $n$ ,  $b \in (\mathcal{A}(I_n)_+)'$  due to  $I_n \subset I$  and  $b \in (\mathcal{A}(I)_+)'$ . The conclusion of (1) implies

$$b = b_n^0 + v_{I_n} b_n^1, \quad (4.52)$$

where  $b_n^0, b_n^1 \in \mathcal{A}(\{I_n\}^c)$ .

Since  $I_n \nearrow I$  and  $I$  is infinite, there exists an  $n_\varepsilon$  such that  $I_{n_\varepsilon}$  contains a point  $i$  which does not belong to  $L_{l_\varepsilon}$ . Then  $i \in I_n$  for all  $n \geq n_\varepsilon$ . Due to  $b_\varepsilon \in \mathcal{A}(L_{l_\varepsilon})$  and  $\{i\}^c \supset L_{l_\varepsilon}$ ,

$$E_{\{i\}^c}(b_\varepsilon) = b_\varepsilon. \quad (4.53)$$

Since  $b_n^0, b_n^1 \in \mathcal{A}(\{I_n\}^c) \subset \mathcal{A}(\{i\}^c)$  for all  $n \geq n_\varepsilon$ , we have

$$\begin{aligned} E_{\{i\}^c}(b_n^0) &= b_n^0 \\ E_{\{i\}^c}(v_{I_n} b_n^1) &= \tau(v_i) v_{I_n \setminus \{i\}} b_n^1 = 0. \end{aligned} \quad (4.54)$$

This implies

$$E_{\{i\}^c}(b) = E_{\{i\}^c}(b_n^0) + E_{\{i\}^c}(v_{I_n} b_n^1) = b_n^0. \quad (4.55)$$

It follows from (4.53) and (4.55)

$$\|b_\varepsilon - b_n^0\| = \|E_{\{i\}^c}(b_\varepsilon) - E_{\{i\}^c}(b)\| \leq \|b_\varepsilon - b\| < \varepsilon.$$

Therefore,

$$\|b - b_n^0\| \leq \|b - b_\varepsilon\| + \|b_\varepsilon - b_n^0\| < 2\varepsilon \quad (4.56)$$

for all  $n \geq n_\varepsilon$ . Hence

$$b = \lim_n b_n^0.$$

For any fixed  $m \in \mathbb{N}$ ,  $b_n^0 \in \mathcal{A}(\{I_n\}^c) \subset \mathcal{A}(\{I_m\}^c)$  for all  $n \geq m$  due to  $I_n \supset I_m$ . Thus  $b \in \mathcal{A}(\{I_m\}^c)$  for any  $m$ . By Corollary 4.12,

$$b \in \cap_m \mathcal{A}(\{I_m\}^c) = \mathcal{A}(\cap_m (I_m^c)) = \mathcal{A}(\{\cup_m I_m\}^c) = \mathcal{A}(I^c).$$

□

As a by-product, we obtain the following.

**Corollary 4.20.** *For any infinite  $I$ , the restriction of  $\Theta$  to  $\mathcal{A}(I)$  is outer.*

*Proof.* We denote the restriction of  $\Theta$  by the same letter. For any infinite subsets  $I$  and  $J$ ,  $(\mathcal{A}(I), \Theta)$  is isomorphic to  $(\mathcal{A}(J), \Theta)$  as a pair of  $\mathbf{C}^*$ -algebra and its automorphism through any bijective map between  $I$  and  $J$ . Therefore it is enough to show the assertion for a proper infinite subset  $I$  of  $\mathbb{Z}^\nu$ .

Suppose that  $u$  is a unitary element in  $\mathcal{A}(I)$  such that

$$u^* a u = \Theta(a),$$

for all  $a \in \mathcal{A}(I)$ . Substituting  $u$  into  $a$ , we have  $\Theta(u) = u$ . Let  $b \in \mathcal{A}(I^c)_-$  and  $b \neq 0$ . Then  $ub \in \mathcal{A}_-$ . By (4.8)

$$ba = \Theta(a)b.$$

Hence  $ub \in \mathcal{A}(I)'$ . Therefore  $ub \in (\mathcal{A}(I)')_-$ , which implies  $ub = 0$  by Lemma 4.16. This implies

$$b = u^*(ub) = 0,$$

a contradiction. □

## 5 Dynamics

### 5.1 Assumptions

We consider a one-parameter group of  $*$ -automorphisms  $\alpha_t$  of the Fermion algebra  $\mathcal{A}$ . Throughout this work,  $\alpha_t$  is assumed to be strongly continuous, that is,  $t \in \mathbb{R} \mapsto \alpha_t(A) \in \mathcal{A}$  is norm continuous for each  $A \in \mathcal{A}$ . In order to associate a potential to the dynamics  $\alpha_t$  (see § 5.4 for details), we need the following two assumptions on  $\alpha_t$  and its generator  $\delta_\alpha$  with the domain  $D(\delta_\alpha)$  :

(I)  $\alpha_t \Theta = \Theta \alpha_t$  for all  $t \in \mathbb{R}$ .

(II)  $\mathcal{A}_o$  is in the domain of  $\delta_\alpha$ , namely,  $\mathcal{A}_o \subset D(\delta_\alpha)$ .

The assumption (I) of  $\Theta$ -even dynamics comes from two sources. On the physical side, the generator of the time translation  $\alpha_t$  should be  $i = \sqrt{-1}$  times the commutator with the energy operator which is a physical observable and hence  $\Theta$ -even.

On the technical side, the potential to be introduced below has to commute with a fixed local element of  $\mathcal{A}$  when the support region of the potential is far away in order that the expression for the action of the generator on that local element converges and makes sense.

For  $\alpha_t$  to be uniquely specified by the associated potential to be introduced in § 5.4, we need the following assumption:

(III)  $\mathcal{A}_o$  is the core of  $\delta_\alpha$ , namely, if  $\delta$  denotes the restriction of  $\delta_\alpha$  to  $\mathcal{A}_o$ , its closure  $\bar{\delta}$  is  $\delta_\alpha$ .

The assumption (III) will be used to derive a conclusion involving  $\alpha_t$  such as the KMS condition from other conditions involving the associated potential such as the Gibbs condition and the variational principle.

Later, when we discuss translation invariant equilibrium states, we will add the assumption of translation invariance:

(IV)  $\alpha_t \tau_k = \tau_k \alpha_t$  for any  $t \in \mathbb{R}$ ,  $k \in \mathbb{Z}^\nu$ .

Later in Proposition 8.1, it will be shown that Assumption (IV) implies Assumption (I).

By Assumptions (I) and (II), the restriction  $\delta$  of  $\delta_\alpha$  to  $\mathcal{A}_o$  satisfies

$$\delta\Theta(A) = \Theta(\delta A) \tag{5.1}$$

for any  $A \in \mathcal{A}_o$ . In the rest of this section, we deal with an arbitrary  $*$ -derivation  $\delta$  with the domain  $\mathcal{A}_o$  commuting with  $\Theta$  (eq.(5.1)) irrespective of whether it comes from a dynamics  $\alpha_t$  or not. Of course, we can use the results about such a general  $\delta$  for the restriction of  $\delta_\alpha$  to  $\mathcal{A}_o$ .

## 5.2 Local Hamiltonians

Since  $\mathcal{A}(\mathbf{I})$  is a finite type  $\mathbf{I}$  factor for each finite subset  $\mathbf{I}$  of  $\mathbb{Z}^\nu$ , there exists a self-adjoint element  $H_{\mathbf{I}}^0 \in \mathcal{A}$  satisfying

$$\delta A = i[H_{\mathbf{I}}^0, A] \quad (5.2)$$

for any  $A \in \mathcal{A}(\mathbf{I})$  where  $\delta$  is any  $*$ -derivation with its domain  $\mathcal{A}_\circ$  and values in  $\mathcal{A}$  (i.e.,  $\delta$  is a linear map from  $\mathcal{A}_\circ$  into  $\mathcal{A}$  satisfying  $\delta(AB) = (\delta A)B + A(\delta B)$  and  $\delta(A^*) = (\delta A)^*$ ). Although this is well-known (see, e.g., [38]), we include its proof for the sake of completeness.

**Lemma 5.1.** *Let  $\{u_{ij}\}$  be a self-adjoint system of matrix units of  $\mathcal{A}(\mathbf{I})$ . Define*

$$h_{ij} \equiv \sum_l u_{li} \delta u_{jl} - \delta_{ij} 2^{-|\mathbf{I}|} \sum_l \sum_m u_{lm} \delta u_{ml}.$$

*Then  $h_{ij} \in \mathcal{A}(\mathbf{I})'$ . Define*

$$iH \equiv \sum_{i,j} u_{ij} h_{ij}.$$

*It satisfies  $H^* = H$  and*

$$[iH, A] = \delta A$$

*for  $A \in \mathcal{A}(\mathbf{I})$ . Furthermore,*

$$E_{\mathbf{I}^c}(H) = 0. \quad (5.3)$$

*Proof.* (1) We first prove  $h_{ij} \in \mathcal{A}(\mathbf{I})'$ . If  $i \neq j$ ,

$$\begin{aligned} [h_{ij}, u_{\alpha\beta}] &= \sum_l u_{li} (\delta u_{jl}) u_{\alpha\beta} - u_{\alpha i} \delta u_{j\beta} \\ &= \sum_l u_{li} (\delta(u_{jl} u_{\alpha\beta}) - u_{jl} \delta u_{\alpha\beta}) - u_{\alpha i} \delta u_{j\beta} \\ &= u_{\alpha i} \delta u_{j\beta} - u_{\alpha i} \delta u_{j\beta} = 0. \end{aligned}$$

If  $i = j$ ,

$$\begin{aligned} [h_{ii}, u_{\alpha\beta}] &= \sum_l u_{li} (\delta u_{il}) u_{\alpha\beta} - u_{\alpha i} \delta u_{i\beta} \\ &\quad - \sum_l \sum_m 2^{-|\mathbf{I}|} \left\{ u_{lm} (\delta u_{ml}) u_{\alpha\beta} - u_{\alpha\beta} u_{lm} \delta u_{ml} \right\} \\ &= \sum_l u_{li} (\delta(u_{il} u_{\alpha\beta}) - u_{il} \delta u_{\alpha\beta}) - u_{\alpha i} \delta u_{i\beta} \\ &\quad - \sum_l \sum_m 2^{-|\mathbf{I}|} \left\{ u_{lm} (\delta(u_{ml} u_{\alpha\beta}) - u_{ml} \delta u_{\alpha\beta}) - u_{\alpha\beta} u_{lm} \delta u_{ml} \right\} \\ &= u_{\alpha i} \delta u_{i\beta} - \delta u_{\alpha\beta} - u_{\alpha i} \delta u_{i\beta} \\ &\quad - \sum_m 2^{-|\mathbf{I}|} u_{\alpha m} \delta u_{m\beta} + 2^{-|\mathbf{I}|} (2^{|\mathbf{I}|} \mathbf{1}) \delta u_{\alpha\beta} + 2^{-|\mathbf{I}|} \sum_m u_{\alpha m} \delta u_{m\beta} \\ &= 0. \end{aligned}$$



(2) We prove  $[iH, u_{\alpha\beta}] = \delta u_{\alpha\beta}$ , which yields  $[iH, A] = \delta A$  for any  $A \in \mathcal{A}(\mathbf{I})$  by linearity.

$$\begin{aligned}
[iH, u_{\alpha\beta}] &= \sum_{i,j} [u_{ij}, u_{\alpha\beta}] h_{ij} = \sum_i u_{i\beta} h_{i\alpha} - \sum_j u_{\alpha j} h_{\beta j} \\
&= \sum_i u_{ii} \delta u_{\alpha\beta} - \sum_m 2^{-|\mathbf{I}|} u_{\alpha m} \delta u_{m\beta} \\
&\quad - u_{\alpha\beta} \sum_j \delta u_{jj} + \sum_m 2^{-|\mathbf{I}|} u_{\alpha m} \delta u_{m\beta} \\
&= \delta u_{\alpha\beta} - u_{\alpha\beta} \delta \left( \sum_j u_{jj} \right) \\
&= \delta u_{\alpha\beta} - u_{\alpha\beta} \delta \mathbf{1} \\
&= \delta u_{\alpha\beta},
\end{aligned}$$

where we have used  $h_{ij} \in \mathcal{A}(\mathbf{I})'$  for the first equality.

(3) Next we prove  $H^* = H$  or  $iH + (iH)^* = 0$ . By using  $u_{ij}^* = u_{ji}$  and  $(\delta a)^* = \delta a^*$ , we obtain

$$\begin{aligned}
iH + (iH)^* &= \sum u_{ij} (h_{ij} + h_{ji}^*), \\
h_{ij} + h_{ji}^* &= \sum_l \left\{ u_{li} \delta u_{jl} + (\delta u_{li}) u_{jl} \right\} - \delta_{ij} 2^{-|\mathbf{I}|} \sum_l \sum_m \left\{ u_{lm} \delta u_{ml} + (\delta u_{lm}) u_{ml} \right\} \\
&= \sum_l \delta (u_{li} u_{jl}) - \delta_{ij} 2^{-|\mathbf{I}|} \sum_l \sum_m \delta (u_{lm} u_{ml}) \\
&= \delta_{ij} \delta \left( \sum_l u_{ll} \right) - \delta_{ij} \delta \left( \sum_l u_{ll} \right) = 0.
\end{aligned}$$

Hence  $iH + (iH)^* = 0$ .

(4) We prove the last statement. Note that  $\tau(u_{ij}) = 2^{-|\mathbf{I}|} \delta_{ij}$ . Hence

$$\begin{aligned}
iE_{\mathbf{I}^c}(H) &= 2^{-|\mathbf{I}|} \sum_i h_{ii} = \sum_i \left\{ \sum_l u_{li} \delta u_{il} - 2^{-|\mathbf{I}|} \sum_l \sum_m u_{lm} \delta u_{ml} \right\} \\
&= 0.
\end{aligned}$$

□

We denote this  $H$  by  $H_{\mathbf{I}}^0$ .

**Lemma 5.2.** *If  $\delta$  is a  $*$ -derivation with domain  $\mathcal{A}_\circ$  and values in  $\mathcal{A}$  commuting with  $\Theta$ , then there exists a self-adjoint element  $H(\mathbf{I}) \in \mathcal{A}_+$  satisfying*

$$\delta A = i[H(\mathbf{I}), A]$$

for all  $A \in \mathcal{A}(\mathbf{I})$  and

$$E_{\mathbf{I}^c}(H(\mathbf{I})) = 0.$$

*Proof.* Due to commutativity of  $\delta$  and  $\Theta$  and  $\Theta^2 = \mathbf{1}$ , we have

$$\begin{aligned}
\delta A &= \Theta(\delta \Theta(A)) = \Theta(i[H_{\mathbf{I}}^0, \Theta(A)]) \\
&= i[\Theta(H_{\mathbf{I}}^0), A]
\end{aligned}$$

for any  $A \in \mathcal{A}(\mathbf{I})$ . Set

$$H(\mathbf{I}) \equiv (H_1^0)_+ = \frac{1}{2}(H_1^0 + \Theta(H_1^0)) \ (\in \mathcal{A}_+). \quad (5.4)$$

Then we have  $H(\mathbf{I})^* = H(\mathbf{I})$  and

$$\delta A = i[H(\mathbf{I}), A] \quad (A \in \mathcal{A}(\mathbf{I})).$$

Since  $E_{\mathbf{I}^c}(H_1^0) = 0$ , it follows from (5.4) and (4.21) that

$$E_{\mathbf{I}^c}(H(\mathbf{I})) = 0.$$

□

The local Hamiltonian operator  $H(\mathbf{I})$  obtained in the above lemma has the following properties:

- (H-i)  $H(\mathbf{I})^* = H(\mathbf{I}) \in \mathcal{A}$ .
- (H-ii)  $\Theta(H(\mathbf{I})) = H(\mathbf{I})$  ( i.e.  $H(\mathbf{I}) \in \mathcal{A}_+$  ).
- (H-iii)  $\delta A = i[H(\mathbf{I}), A]$  ( $A \in \mathcal{A}(\mathbf{I})$ ).
- (H-iv)  $E_{\mathbf{I}^c}(H(\mathbf{I})) = 0$ .

*Remark.* The property (H-iv) implies

$$\tau(H(\mathbf{I})) = \tau(E_{\mathbf{I}^c}(H(\mathbf{I}))) = 0. \quad (5.5)$$

**Lemma 5.3.**  $H(\mathbf{I})$  satisfying (H-ii)–(H-iv) is uniquely determined by  $\delta$ .

*Proof.* If  $H(\mathbf{I})$  and  $H(\mathbf{I})'$  satisfy (H-ii)–(H-iv), then  $\Delta = H(\mathbf{I}) - H(\mathbf{I})'$  satisfies  $[\Delta, A] = 0$  for all  $A \in \mathcal{A}(\mathbf{I})$  due to (H-iii). By Lemma 4.15 and (H-ii) for  $\Delta$ ,

$$\Delta \in \mathcal{A}(\mathbf{I})' \cap \mathcal{A}_+ = \mathcal{A}(\mathbf{I}^c)_+.$$

Hence (H-iv) implies

$$\Delta = E_{\mathbf{I}^c}(\Delta) = E_{\mathbf{I}^c}(H(\mathbf{I})) - E_{\mathbf{I}^c}(H(\mathbf{I}')) = 0.$$

Therefore  $H(\mathbf{I})$  satisfying (H-ii)–(H-iv) is unique. □

We call  $H(\mathbf{I})$  the standard Hamiltonian for the region  $\mathbf{I}$ .

*Remark.* For the empty set  $\emptyset$ ,  $H(\emptyset) = 0$  by (H-iv).

Under the conditions (H-ii)–(H-iv), the property  $H(\mathbf{I})^* = H(\mathbf{I})$  of (H-i) and the property  $(\delta A)^* = \delta A^*$  ( $A \in \mathcal{A}(\mathbf{I})$ ) for  $\delta$  are equivalent, because of the following reason. If  $H(\mathbf{I})^* = H(\mathbf{I})$ , then  $(\delta A)^* = \delta A^*$  immediately follows from (H-iii). If  $(\delta A)^* = \delta A^*$ , then  $H(\mathbf{I})^*$  satisfies (H-iii) along with (H-ii) and (H-iv). Hence  $H(\mathbf{I})^* = H(\mathbf{I})$  by the uniqueness result Lemma 5.3.

**Lemma 5.4.** If  $\mathbf{I} \subset \mathbf{J}$  is a pair of finite subsets, then

$$H(\mathbf{I}) = H(\mathbf{J}) - E_{\mathbf{I}^c}(H(\mathbf{J})). \quad (5.6)$$

*Proof.*  $H(J)$  satisfies (H-ii) and (H-iii) for the region  $I \subset J$ . Furthermore,  $E_{I^c}(H(J)) \in \mathcal{A}(I^c)_+$  due to (H-ii) for  $H(J)$  and hence it commutes with  $A \in \mathcal{A}(I)$ . Therefore  $H(J) - E_{I^c}(H(J))$  satisfies (H-ii)-(H-iv) for the region  $I$ . By the uniqueness (Lemma 5.3), we obtain  $H(I) = H(J) - E_{I^c}(H(J))$ .  $\square$

We give the number (H-v) to the condition above:

(H-v)  $H(I) = H(J) - E_{I^c}(H(J))$  for any finite subsets  $I \subset J$  of  $\mathbb{Z}^\nu$ .

The proof above has shown that (H-v) is derived from (H-ii)-(H-iv).

So far we have derived the properties (H-i), (H-ii), (H-iv) and (H-v) for the family  $\{H(I)\}$  from its definition in terms of  $\delta$  through the relation (H-iii).

In the converse direction, any family of an element  $H(I) \in \mathcal{A}$  for each finite subset  $I$  of  $\mathbb{Z}^\nu$  defines a derivation  $\delta$  on  $\mathcal{A}_o$  by (H-iii).

This definition requires a consistency: if  $A \in \mathcal{A}(I)$  and  $A \in \mathcal{A}(J)$ , we have a definition of  $\delta(A)$  by  $H(I)$  and  $H(J)$ . The proof that they are the same is given as follows. First we note that  $A \in \mathcal{A}(I) \cap \mathcal{A}(J) = \mathcal{A}(I \cap J)$ . Thus it is enough to show

$$[H(I), A] = [H(K), A] \quad (5.7)$$

for any  $K \subset I$  and  $A \in \mathcal{A}(K)$ , because, using this identity for the pair  $I \supset K = I \cap J$  and  $J \supset K$ ; we obtain  $[H(I), A] = [H(J), A]$  for any  $A \in \mathcal{A}(I \cap J)$ .

Since  $E_{K^c}(H(I))$  is  $\Theta$ -even by (H-ii) and (4.21),  $E_{K^c}(H(I))$  is in  $\mathcal{A}(K^c)_+$  and commutes with  $A \in \mathcal{A}(K)$ . By (H-v),

$$H(K) = H(I) - E_{K^c}(H(I))$$

which leads to the consistency equation (5.7).

$\delta$  defined by (H-iii) is a  $*$ -derivation with domain  $\mathcal{A}_o$  due to (H-i), and commutes with  $\Theta$  by (H-ii).

We have not used (H-iv) in this argument, but have imposed it on  $H(I)$  to obtain the uniqueness of  $H(I)$  for a given  $\delta$ . Namely, by Lemma 5.2 and Lemma 5.3, the correspondence of  $\delta$  and  $H(I)$  is bijective, for which the condition (H-iv) is used.

Summarizing the argument so far, we have obtained Theorem 5.7 stated below after introduction of two definitions.

**Definition 5.5.** *The real vector space of all  $*$ -derivations with their definition domain  $\mathcal{A}_o$  and commuting with  $\Theta$  (on  $\mathcal{A}_o$ ) is denoted by  $\Delta(\mathcal{A}_o)$ .*

*Remark.* Under Assumptions (I) and (II), the restriction  $\delta$  of the generator  $\delta_\alpha$  of  $\alpha_t$  belongs to  $\Delta(\mathcal{A}_o)$

**Definition 5.6.** *The real vector space of functions  $H(I)$  of finite subsets  $I$  satisfying the following four conditions is denoted by  $\mathbf{H}$  and its element  $H$  is called a local Hamiltonian.*

(H-i)  $H(I)^* = H(I) \in \mathcal{A}$ ,

(H-ii)  $\Theta(H(I)) = H(I)$  (i.e.  $H(I) \in \mathcal{A}_+$ )

(H-iv)  $E_{I^c}(H(I)) = 0$ ,

(H-v)  $H(I) = H(J) - E_{I^c}(H(J))$  for any finite subsets  $I \subset J$  of  $\mathbb{Z}^\nu$ .

**Theorem 5.7.** *The following relation between  $H \in \mathbf{H}$  and  $\delta \in \Delta(\mathcal{A}_o)$  gives a bijective, real linear map from  $\mathbf{H}$  to  $\Delta(\mathcal{A}_o)$ .*

(H-iii)  $\delta A = i[H(I), A]$  ( $A \in \mathcal{A}(I)$ ).

*Remark.* The value  $\delta A$  of the derivation  $\delta \in \Delta(\mathcal{A}_o)$  for  $A \in \mathcal{A}_o$  is in general not in  $\mathcal{A}_o$ .

### 5.3 Internal Energy

For a finite subset  $I$  of  $\mathbb{Z}^\nu$ , set

$$U(I) \equiv E_I(H(I)) \quad (\in \mathcal{A}(I)) \quad (5.8)$$

and call it the internal energy for the region  $I$ . Due to  $H(\emptyset) = 0$ ,  $U(\emptyset) = 0$ .

Due to the property (5.5),

$$E_I E_{I^c}((H(J))) = \tau((H(J))) = 0.$$

By (H-v), we obtain for  $I \subset J$

$$\begin{aligned} U(I) &= E_I H(I) = E_I (\{H(J) - E_{I^c}(H(J))\}) \\ &= E_I H(J) = E_I E_J H(J) = E_I U(J). \end{aligned} \quad (5.9)$$

Furthermore, for any finite subset  $I$  and any subset  $J$  of  $\mathbb{Z}^\nu$ , we have

$$E_J(U(I)) = E_J E_I(U(I)) = E_{J \cap I}(U(I)) = U(I \cap J), \quad (5.10)$$

where the last equality is due to (5.9). Due to (5.5),

$$\tau(U(I)) = \tau(E_I(H(I))) = \tau(H(I)) = 0. \quad (5.11)$$

Let us denote

$$H_J(I) \equiv E_J(H(I)). \quad (5.12)$$

**Lemma 5.8.** (1) *For any pair of finite subsets  $I$  and  $J$ ,*

$$H_J(I) = U(J) - U(I^c \cap J). \quad (5.13)$$

(2) *For any finite subset  $I$ ,*

$$H(I) = \lim_{J \nearrow \mathbb{Z}^\nu} (U(J) - U(I^c \cap J)). \quad (5.14)$$

*Proof.* (1): By applying (H-v) for pairs  $I \supset I \cap J$  and  $J \supset I \cap J$ , we obtain

$$\begin{aligned} H(I \cap J) &= H(I) - E_{(I \cap J)^c}(H(I)), \\ H(I \cap J) &= H(J) - E_{(I \cap J)^c}(H(J)). \end{aligned}$$

Therefore

$$H(I) = H(J) - E_{(I \cap J)^c}(H(J) - H(I)).$$

By applying  $E_J$  to this equation, we obtain

$$H_J(I) = U(J) - E_J E_{(I \cap J)^c}(H(J) - H(I)).$$

Since

$$J \cap (I \cap J)^c = J \cap (I^c \cup J^c) = (J \cap I^c) \cup (J \cap J^c) = J \cap I^c,$$

we obtain

$$E_J E_{(I \cap J)^c} = E_{J \cap (I \cap J)^c} = E_{J \cap I^c} = E_J E_{I^c} = E_{I^c} E_J.$$

Since  $E_{I^c}(H(I)) = 0$  by (H-iv), we have

$$E_J E_{(I \cap J)^c}(H(J) - H(I)) = E_{I^c} E_J(H(J)) = E_{I^c}(U(J)).$$

Thus

$$H_J(I) = U(J) - E_{I^c}(U(J)).$$

By this and (5.10), we arrive at (5.13).

(2): By (4.23), we have

$$H(I) = \lim_{J \nearrow \mathbb{Z}^\nu} H_J(I). \quad (5.15)$$

This and (5.13) imply the desired (5.14).  $\square$

## 5.4 Potential

We introduce the potential  $\{\Phi(I)\}$  in terms of  $\{H(I)\}$  and derive its characterizing properties. As a consequence, we establish the one-to-one correspondence between  $\{\Phi(I)\}$  and  $\{H(I)\}$ .

**Lemma 5.9.** *For a given  $\{H(I)\} \in \mathbf{H}$  and the corresponding  $\{U(I)\}$ , there exists one and only one family of  $\{\Phi(I) \in \mathcal{A}; \text{ finite } I \subset \mathbb{Z}^\nu\}$  satisfying the following conditions :*

- (1)  $\Phi(I) \in \mathcal{A}(I)$ .
- (2)  $\Phi(I)^* = \Phi(I)$ ,  $\Theta(\Phi(I)) = \Phi(I)$ ,  $\Phi(\emptyset) = 0$ .
- (3)  $E_J(\Phi(I)) = 0$  if  $J \subset I$  and  $J \neq I$ .
- (4)  $U(I) = \sum_{K \subset I} \Phi(K)$ .
- (5)  $H(I) = \lim_{J \nearrow \mathbb{Z}^\nu} \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \subset J\}$ .

*Proof.* We show this lemma in several steps.

Step 1. Existence of  $\Phi$  satisfying (1) and (4) for all finite  $I$ .

The following expression for  $\Phi(I)$  in terms of  $U(K)$ ,  $K \subset I$  satisfies (1) and (4) for all  $I$  and hence the existence.

$$\Phi(I) = \sum_{K \subset I} (-1)^{|I|-|K|} U(K). \quad (5.16)$$

In fact, substituting this expression into  $\sum_{J \subset I} \Phi(J)$ , we obtain

$$\begin{aligned} \sum_{J \subset I} \sum_{K \subset J} (-1)^{|J|-|K|} U(K) &= \sum_{K \subset I} \alpha(K) U(K), \\ \alpha(K) &= \sum_{J: K \subset J \subset I} (-1)^{|J|-|K|} = \sum_{m=|K|}^{|I|} (-1)^{m-|K|} \beta_m, \end{aligned} \quad (5.17)$$

where  $\beta_m$  is the number of distinct  $J$  satisfying

$$K \subset J \subset I, \quad |J| = m.$$

This is the number of way for choosing  $m - |K|$  elements (for  $J \setminus K$ ) out of  $I \setminus K$ , which is  $\binom{|I|-|K|}{m-|K|}$ . Putting  $l = m - |K|$ ,  $n = |I| - |K|$ , we obtain

$$\alpha(K) = \sum_{l=0}^n (-1)^l \binom{n}{l} = (1-1)^n = 0$$

for all  $K \neq I$  (then  $n \geq 1$ ), while we have  $\alpha(I) = 1$ . Hence (4) is satisfied by  $\Phi(I)$  given as (5.16) for all  $I$ .

Step 2. Uniqueness of  $\Phi$  satisfying (4).

The relation (4) implies

$$\Phi(I) = U(I) - \sum_{K \subset I, K \neq I} \Phi(K) \quad (5.18)$$

which obviously determines  $\Phi(I)$  uniquely for a given  $\{U(I)\}$  by the mathematical induction on  $|I| = m$  starting from  $\Phi(\emptyset) = U(\emptyset) = 0$ .

Step 3. Property (2).

We already obtain  $\Phi(\emptyset) = 0$ . Since  $U(I)^* = U(I)$  and  $\Theta(U(I)) = U(I)$ ,  $\Phi(I)$  defined by (5.16) as a real linear combination of  $U(K)$ ,  $K \subset I$  satisfies (2).

Step 4. Property (3).

We note that (3) is equivalent to the following condition:

$$E_J(\Phi(I)) = 0, \quad \text{for } J \not\supset I, \quad (5.19)$$

because  $E_J(\Phi(I)) = E_J E_I(\Phi(I)) = E_{J \cap I}(\Phi(I))$  by Theorem 4.7,  $J \cap I \subset I$ , and  $J \cap I \neq I$  if and only if  $J \not\supset I$ . On the other hand,  $E_J(\Phi(I)) = \Phi(I)$  if  $J \supset I$  due to  $\Phi(I) \in \mathcal{A}(I) \subset \mathcal{A}(J)$ .

We now prove (3) by the mathematical induction on  $|I| = m$ . For  $m = 1$ , the only  $J$  satisfying  $J \subset I$  and  $J \neq I$  is  $J = \emptyset$  for which  $\Phi(J) = 0$ . Then  $\Phi(I) = U(I)$  and

$$E_J(\Phi(I)) = \tau(\Phi(I)) \mathbf{1} = \tau(U(I)) = 0$$

due to (5.11). Suppose (3) holds for  $|I| < m$ . We consider  $I$  with  $|I| = m$ . We apply  $E_J$  (for  $J \subset I$ ,  $J \neq I$ ) on both sides of (5.18). All  $K$  in the summation on the right-hand side satisfy  $|K| < m$  due to  $K \subset I$  and  $K \neq I$ . Hence the inductive assumption is applicable to  $\Phi(K)$  on the right-hand side. If  $K \not\subset J$ , we have  $E_J(\Phi(K)) = 0$  by (5.19). If  $K \subset J$ , we have  $E_J(\Phi(K)) = \Phi(K)$ . Therefore, by using  $E_J U(I) = U(J)$  (due to  $J \subset I$ ), we obtain

$$\begin{aligned} E_J \Phi(I) &= E_J U(I) - \sum_{K \subset I, K \neq I} E_J \Phi(K) \\ &= U(J) - \sum_{K \subset J} \Phi(K) = 0. \end{aligned}$$

This proves (3).

Step 5. Property (5).

For a finite subset  $J$  and  $I \subset J$ ,  $H_J(I)$  is written in terms of  $\Phi$  by (5.13) and (4) as

$$H_J(I) (= E_J(H(I))) = \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \subset J\}. \quad (5.20)$$

Due to (5.15),  $\Phi$  satisfies (5).  $\square$

We collect useful formulae for  $U$  and  $H$  in terms of  $\Phi$  which have been obtained above:

$$U(I) = \sum_{K \subset I} \Phi(K), \quad (5.21)$$

$$H_J(I) = \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \subset J\}, \quad (5.22)$$

$$H(I) = \lim_{J \nearrow \mathbb{Z}^\nu} \left( \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \subset J\} \right) (= \lim_{J \nearrow \mathbb{Z}^\nu} H_J(I)). \quad (5.23)$$

**Definition 5.10.** A function  $\Phi$  of finite subsets  $I$  of  $\mathbb{Z}^\nu$  with the value  $\Phi(I)$  in  $\mathcal{A}$  is called a standard potential if it satisfies the following conditions :

( $\Phi$ -a)  $\Phi(I) \in \mathcal{A}(I)$ ,  $\Phi(\emptyset) = 0$ .

( $\Phi$ -b)  $\Phi(I)^* = \Phi(I)$ .

( $\Phi$ -c)  $\Theta(\Phi(I)) = \Phi(I)$ .

( $\Phi$ -d)  $E_J(\Phi(I)) = 0$  if  $J \subset I$  and  $J \neq I$ .

( $\Phi$ -e) For each fixed finite subset  $I$  of  $\mathbb{Z}^\nu$ , the net

$$H_J(I) = \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \subset J\},$$

is a Cauchy net in the norm topology of  $\mathcal{A}$  for  $J \nearrow \mathbb{Z}^\nu$ . The index set for the net is the set of all finite subsets  $J$  of  $\mathbb{Z}^\nu$ , partially ordered by the set inclusion.

*Remark.* ( $\Phi$ -d) is equivalent to the following condition:

( $\Phi$ -d)'  $E_J(\Phi(I)) = 0$  unless  $I \subset J$ ,

because  $E_J(\Phi(I)) = E_J E_I(\Phi(I)) = E_{J \cap I}(\Phi(I))$ .

**Definition 5.11.** The real vector space of all standard potentials is denoted by  $\mathcal{P}$ .

*Remark.*  $\mathcal{P}$  is a real vector space as a function space, where the linear operation is defined by

$$(c\Phi + d\Psi)(I) = c\Phi(I) + d\Psi(I), \quad c, d \in \mathbb{R}, \quad \Phi, \Psi \in \mathcal{P}. \quad (5.24)$$

We show the one-to-one correspondence of  $\Phi \in \mathcal{P}$  and  $H \in \mathbf{H}$ .

**Theorem 5.12.** The equations (5.22) and (5.23) for  $\Phi \in \mathcal{P}$  and  $H \in \mathbf{H}$  give a bijective, real linear map from  $\mathcal{P}$  to  $\mathbf{H}$ .

*Proof.* First note that (4) of Lemma 5.9 is satisfied for  $U(I) = E_I(H(I))$  due to ( $\Phi$ -d), if (5.22) and (5.23) are satisfied. By Lemma 5.9, there exists a unique  $\Phi \in \mathcal{P}$  satisfying (5.22) and (5.23) for any given  $H \in \mathbf{H}$ .

The map is evidently linear. The only remaining task is to prove the property (H-i), (H-ii), (H-iv) and (H-v) for the  $H(I)$  given by (5.22) and (5.23), on the basis of  $(\Phi\text{-a})$ - $(\Phi\text{-e})$ . (H-i), (H-ii) and (H-iv) follow from  $(\Phi\text{-b})$ ,  $(\Phi\text{-c})$  and  $(\Phi\text{-d})'$ , respectively.

To show (H-v), let  $L$  be a finite subset containing  $J \supset I$ . Then

$$\begin{aligned} H_L(J) - H_L(I) &= \sum_K \{\Phi(K); K \cap J \neq \emptyset, K \cap I = \emptyset, K \subset L\} \\ &= E_{I^c} \left( \sum_K \{\Phi(K); K \cap J \neq \emptyset, K \subset L\} \right) \\ &= E_{I^c}(H_L(J)) \end{aligned}$$

due to (5.22),  $(\Phi\text{-a})$  and  $(\Phi\text{-d})'$ . By taking limit  $L \nearrow \mathbb{Z}^\nu$ , we obtain

$$H(J) - H(I) = E_{I^c}(H(J)),$$

where the convergence is due to  $(\Phi\text{-e})$  and  $\|E_{I^c}\| = 1$ .  $\square$

*Remark.* We will use later the real linearity of the above map:

$$H_{c\Phi+d\Psi}(I) = cH_\Phi(I) + dH_\Psi(I), \quad c, d \in \mathbb{R}, \quad \Phi, \Psi \in \mathcal{P}, \quad (5.25)$$

$$U_{c\Phi+d\Psi}(I) = cU_\Phi(I) + dU_\Psi(I), \quad c, d \in \mathbb{R}, \quad \Phi, \Psi \in \mathcal{P}, \quad (5.26)$$

where  $H_\Phi(I)$  and  $U_\Phi(I)$  denote  $H(I)$  and  $U(I)$  corresponding to  $\Phi \in \mathcal{P}$ .

**Theorem 5.13.** *The following relation between  $\Phi \in \mathcal{P}$  and  $\delta_\Phi \in \Delta(\mathcal{A}_o)$  gives a bijective, real linear map from  $\mathcal{P}$  to  $\Delta(\mathcal{A}_o)$ .*

$$\delta_\Phi A = i[H(I), A] \quad (A \in \mathcal{A}(I)), \quad (5.27)$$

$$H(I) = \lim_{J \nearrow \mathbb{Z}^\nu} \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \subset J\}. \quad (5.28)$$

*Proof.* This is a consequence of Theorem 5.7 and Theorem 5.12.  $\square$

*Remark 1.* The technique using the conditional expectations for associating a unique standard potential with a given  $*$ -derivation has been developed for quantum spin lattice systems by one of the authors [12]. The corresponding formalism for classical lattice systems is developed in [13]. Also see [23] where  $E_I$  for the quantum spin case is called a partial trace.

*Remark 2.* We note that  $\mathcal{P}$  is a Fréchet space with respect to a countable family of seminorms  $\{\|H(\{i\})\|\}, i \in \mathbb{Z}^\nu$ .

## 5.5 General Potential

If the function

$$\Phi : I \in \{\text{finite subsets of } \mathbb{Z}^\nu\} \longmapsto \Phi(I) \quad (5.29)$$

satisfies  $(\Phi\text{-a})$ ,  $(\Phi\text{-b})$ ,  $(\Phi\text{-c})$  and  $(\Phi\text{-e})$ , we call it a general potential.



By  $(\Phi\text{-e})$ , we define  $H(I)$  by (5.23) and (5.22). Then, for any finite subsets  $K \supset I$ ,

$$H(K) - H(I) = \lim_{J \nearrow \mathbb{Z}^\nu} \sum_L \{\Phi(L); L \cap K \neq \emptyset, L \cap I = \emptyset, L \subset J\} \quad (5.30)$$

due to  $(\Phi\text{-e})$ . Therefore, we can define  $\delta_\Phi$  with the domain  $\mathcal{A}_\circ$  by

$$\delta_\Phi A = i[H(I), A] \quad \text{for } A \in \mathcal{A}(I), \quad (5.31)$$

which is a consistent definition due to (5.30) by essentially the same argument as the one leading to (5.7). The properties  $(\Phi\text{-a})$ ,  $(\Phi\text{-b})$ ,  $(\Phi\text{-c})$ , and  $(\Phi\text{-e})$  imply that  $\delta_\Phi \in \Delta(\mathcal{A}_\circ)$ . Two general potentials  $\Phi$  and  $\Phi'$  are said to be equivalent if  $\delta_\Phi = \delta_{\Phi'}$ . It follows from Theorem 5.13 that there is a unique standard potential which is equivalent to any given general potential defined above. The equivalence is discussed, e.g., in [23] and [40] with the name of physical equivalence. We will consider the consequence of equivalence for a specific class of general potentials in § 14.

## 6 KMS Condition

### 6.1 KMS Condition

We recall the definition of the KMS condition for a given dynamics  $\alpha_t$  of  $\mathcal{A}$  (see e.g. [17]).

**Definition 6.1.** *A state  $\varphi$  of  $\mathcal{A}$  is called an  $\alpha_t$ -KMS state at the inverse temperature  $\beta \in \mathbb{R}$  or  $(\alpha_t, \beta)$ -KMS state (or more simply KMS state) if it satisfies one of the following two equivalent conditions :*

(A) *Let  $D_\beta$  be the strip region*

$$\begin{aligned} D_\beta &= \left\{ z \in \mathbb{C}; 0 \leq \operatorname{Im} z \leq \beta \right\} \quad \text{if } \beta \geq 0, \\ &= \left\{ z \in \mathbb{C}; \beta \leq \operatorname{Im} z \leq 0 \right\} \quad \text{if } \beta < 0, \end{aligned}$$

*in the complex plane  $\mathbb{C}$  and  $\mathring{D}_\beta$  be its interior.*

*For every  $A$  and  $B$  in  $\mathcal{A}$ , there exists a function  $F(z)$  of  $z \in D_\beta$  (depending on  $A$  and  $B$ ) such that*

- (1)  *$F(z)$  is analytic in  $\mathring{D}_\beta$ ,*
- (2)  *$F(z)$  is continuous and bounded on  $D_\beta$ ,*
- (3) *For all real  $t \in \mathbb{R}$ ,*

$$F(t) = \varphi(A\alpha_t(B)), \quad F(t + i\beta) = \varphi(\alpha_t(B)A).$$

(B) *Let  $\mathcal{A}_{ent}$  be the set of all  $B \in \mathcal{A}$  for which  $\alpha_t(B)$  has an analytic extension to  $\mathcal{A}$ -valued entire function  $\alpha_z(B)$  as a function of  $z \in \mathbb{C}$ . For  $A \in \mathcal{A}$  and  $B \in \mathcal{A}_{ent}$ ,*

$$\varphi(A\alpha_{i\beta}(B)) = \varphi(BA).$$

*Remark.* In (A), the condition (1) is empty if  $\beta = 0$ . The boundedness in (2) can be omitted (see, e.g., proposition 5.3.7 in [17]).  $\mathcal{A}_{ent}$  is known to be dense in  $\mathcal{A}$ .

For a state  $\varphi$  on  $\mathcal{A}$ , let  $\{\mathcal{H}_\varphi, \pi_\varphi, \Omega_\varphi\}$  denote its GNS triplet, namely,  $\pi_\varphi$  is a (GNS) representation of  $\mathcal{A}$  on the Hilbert space  $\mathcal{H}_\varphi$ , and  $\Omega_\varphi$  is a cyclic unit vector in  $\mathcal{H}_\varphi$ , representing  $\varphi$  as the vector state. If  $\varphi$  is an  $(\alpha_t, \beta)$ -KMS state, then  $\Omega_\varphi$  is separating for the generated von Neumann algebra  $\mathfrak{M}_\varphi \equiv \pi_\varphi(\mathcal{A})''$ . Let  $\Delta_\varphi$  and  $\sigma_t^\varphi$  be the modular operator and modular automorphisms for  $\Omega_\varphi$  and  $\varphi$ , respectively, [42].

The KMS condition implies that

$$\sigma_t^\varphi(\pi_\varphi(A)) = \pi_\varphi(\alpha_{-\beta t}(A)), \quad A \in \mathcal{A}. \quad (6.1)$$

It is a result of Takesaki [42] that the KMS condition of a one-parameter automorphism group of a von Neumann algebra with respect to a cyclic vector implies the separating property of the vector, and the modular automorphism group of the von Neumann algebra with respect to the cyclic and separating vector is characterized by the KMS condition at  $\beta = -1$  with respect to the state given by that vector.

For the sake of brevity in stating an assumption later, we use the following terminology.

**Definition 6.2.** A state  $\varphi$  is said to be modular if  $\Omega_\varphi$  is separating for  $\pi_\varphi(\mathcal{A})''$ .

## 6.2 Differential KMS Condition

It is convenient to introduce the following condition in terms of the generator  $\delta_\alpha$  of the dynamics  $\alpha_t$ , equivalent to the KMS condition with respect to  $\alpha_t$ .

**Definition 6.3.** Let  $\delta$  be a  $*$ -derivation of  $\mathcal{A}$  with its domain  $D(\delta)$ . A state  $\varphi$  is said to satisfy the differential  $(\delta, \beta)$ -KMS condition (or briefly,  $(\delta, \beta)$ -dKMS condition) if the following two conditions are satisfied

(C-1)  $\varphi(A^*\delta A)$  is pure imaginary for all  $A \in D(\delta)$ .

(C-2)  $-i\beta\varphi(A^*\delta A) \geq S(\varphi(AA^*), \varphi(A^*A))$  for all  $A \in D(\delta)$  where the function  $S(x, y)$  is given for  $x \geq 0, y \geq 0$  by:

$$\begin{aligned} S(x, y) &= y \log y - y \log x \quad \text{if } x > 0, y > 0, \\ S(x, y) &= +\infty \quad \text{if } x = 0, y > 0, \\ S(x, y) &= 0 \quad \text{if } x \geq 0, y = 0. \end{aligned}$$

We use the following known result (see, e.g., Theorem 5.3.15 in [17]).

**Theorem 6.4.** Let  $\delta_\alpha$  be a generator of  $\alpha_t$ , namely,  $e^{t\delta_\alpha} = \alpha_t$ . Then the  $(\delta_\alpha, \beta)$ -dKMS condition and the  $(\alpha_t, \beta)$ -KMS condition are equivalent.

*Remark.* The function  $S(x, y)$  is the relative entropy for linear functionals of one-dimensional  $*$ -algebra. The order of the arguments  $x, y$  in our notation is opposite to that of the definition in [45]. (Both the order of the argument and the sign are opposite to those in [17].) Our definition here is in accordance with our definition of the relative entropy previously given.

**Lemma 6.5.**  $S(x, y)$  is convex and lower semi-continuous in  $x, y$ .

*Proof.* A convenient expression for  $S(x, y)$  is

$$S(x, y) = \sup_n \sup_{s(t)} \left\{ y \log n - \int_{\frac{1}{n}}^{\infty} \left( y s(t)^2 + t^{-1} x \{1 - s(t)\}^2 \right) \frac{dt}{t} \right\}, \quad (6.2)$$

where  $s(t)$  varies over the linear span of characteristic functions of finite intervals in  $[0, +\infty)$ . The equality is immediate for  $x = 0, y > 0$  as well as for  $x \geq 0, y = 0$ . For  $x > 0, y > 0$ , (6.2) follows from identities for  $\lambda = x/y$ .

$$\begin{aligned} y(\log y - \log x) &= \sup_n \left\{ -y \log \left( \frac{x}{y} + \frac{1}{n} \right) \right\} \\ &= \sup_n \left\{ y \log n - \int_{\frac{1}{n}}^{\infty} y \frac{\lambda}{t + \lambda} \frac{dt}{t} \right\}, \\ -y \frac{\lambda}{t + \lambda} &= \sup_{s \in \mathbb{R}} \left\{ - \left( y s^2 + x t^{-1} (1 - s)^2 \right) \right\}. \end{aligned}$$

From the expression above,  $S(x, y)$  is seen to be convex and lower semi-continuous in  $(x, y)$  because it is a supremum of homogeneous linear functions of  $(x, y)$ .

(The variational expression (6.2) for general von Neumann algebras is established by Kosaki [25]. This expression indicates manifestly some basic properties of relative entropy for the general case.)  $\square$

**Lemma 6.6.** *The conditions (C-1) and (C-2) are stable under the simultaneous limit of  $A$  and  $\delta A$  in norm topology and  $\varphi$  in the weak\* topology as well as under the convex combination of states  $\varphi$ .*

*Proof.* Let  $A_n, A \in D(\delta)$ ,  $\|A_n - A\| \rightarrow 0$ ,  $\|\delta A_n - \delta A\| \rightarrow 0$ ,  $|\varphi_n(B) - \varphi(B)| \rightarrow 0$  for every  $B \in \mathcal{A}$ . Then

$$\begin{aligned} &|\varphi_n(A_n^* \delta A_n) - \varphi(A^* \delta A)| \\ &\leq |\varphi_n(A_n^* \delta A_n - A \delta A)| + |\varphi_n(A \delta A) - \varphi(A \delta A)|, \end{aligned}$$

which converges to 0 as  $n \rightarrow \infty$ . Therefore, the condition (C-1) holds for  $\varphi$  and  $A$  if it holds for  $\varphi_n$  and  $A_n$ .

Similarly,

$$\varphi_n(A_n A_n^*) \rightarrow \varphi(A A^*), \quad \varphi_n(A_n^* A_n) \rightarrow \varphi(A^* A),$$

as  $n \rightarrow \infty$ . By the lower semi-continuity of  $S(x, y)$  in  $(x, y)$ , we then obtain

$$S(\varphi(A A^*), \varphi(A^* A)) \leq \liminf_n S(\varphi(A_n A_n^*), \varphi(A_n^* A_n)).$$

Hence we obtain the condition (C-2) for  $\varphi$  and  $A$  if it holds for  $\varphi_n$  and  $A_n$ . Since  $\varphi(A^* \delta A)$  is affine in  $\varphi$  while  $S(\varphi(A A^*), \varphi(A^* A))$  is convex in  $\varphi$ , the conditions (C-1) and (C-2) are stable under the convex combination of  $\varphi$ .  $\square$

**Corollary 6.7.** *Let  $\alpha_t$  be a one-parameter group of \*-automorphisms of  $\mathcal{A}$  satisfying the conditions (II) and (III). Let  $\delta_\alpha$  be the generator of  $\alpha_t$ . Then a state  $\varphi$  is an  $(\alpha_t, \beta)$ -KMS state if and only if it is a  $(\delta, \beta)$ -dKMS state, where  $\delta$  denotes the restriction of  $\delta_\alpha$  to  $\mathcal{A}_o$ .*

*Proof.* The restriction  $\delta$  of  $\delta_\alpha$  to  $\mathcal{A}_\circ$  makes sense due to the assumption (II). By Theorem 6.4, it suffices to prove that the dKMS condition for  $\delta$  implies the same for  $\delta_\alpha$ . By Assumption (III), there exists a sequence  $A_n \in \mathcal{A}_\circ$  for any given  $A \in D(\delta_\alpha)$  such that  $\|A_n - A\| \rightarrow 0$ ,  $\|\delta A_n - \delta_\alpha A\| \rightarrow 0$ . Hence the conditions (C-1) and (C-2) for  $\delta$  imply the same for  $\delta_\alpha$  due to Lemma 6.6.  $\square$

## 7 Gibbs Condition

In this section, we define the Gibbs condition. We first recall the notion of perturbation of dynamics and states.

### 7.1 Inner Perturbation

Consider a given dynamics  $\alpha_t$  of  $\mathcal{A}$  with its generator  $\delta$  on the domain  $D(\delta)$ . For each  $h = h^* \in \mathcal{A}$ , there exists the unique perturbed dynamics  $\alpha_t^h$  of  $\mathcal{A}$  with its generator  $\delta^h$  given by

$$\delta^h(A) \equiv \delta(A) + i[h, A] \quad (A \in D(\delta)) \quad (7.1)$$

on the same domain as the generator  $\delta$  of  $\alpha_t$ . This  $\alpha_t^h(A)$  is explicitly given by

$$\alpha_t^h(A) = u_t^h \alpha_t(A) (u_t^h)^* \quad (7.2)$$

where

$$u_t^h \equiv \mathbf{1} + \sum_{m=1}^{\infty} i^m \int_0^t dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{m-1}} dt_m \alpha_{t_m}(h) \cdots \alpha_{t_1}(h). \quad (7.3)$$

This is unitary and satisfies the following cocycle equation:

$$u_s^h \alpha_s(u_t^h) = u_{s+t}^h$$

The same statements hold for a von Neumann algebra  $\mathfrak{M}$  and its one parameter group of  $*$ -automorphisms  $\alpha_t$ ; the  $t$ -continuity of  $\alpha_t$  for each fixed  $x \in \mathfrak{M}$  in the strong operator topology of  $\mathfrak{M}$  is to be assumed.

Let  $\Omega$  be a cyclic and separating vector for  $\mathfrak{M}$ . Let  $\Delta_\Omega$  be the modular operator for  $\Omega$  and  $\sigma_t^\omega$  be the corresponding modular automorphism group

$$\sigma_t^\omega(x) = \Delta_\Omega^{it} x \Delta_\Omega^{-it},$$

where  $\omega$  indicates the positive linear functional

$$\omega(x) = (\Omega, x\Omega), \quad (x \in \mathfrak{M}).$$

For  $h = h^* \in \mathfrak{M}$ , the perturbed vector  $\Omega^h$  is given by

$$\begin{aligned} & \Omega^h \\ & \equiv \sum_{m=0}^{\infty} \int_0^{\frac{1}{2}} dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{m-1}} dt_m \Delta_\varphi^{t_m} \pi_\varphi(h) \Delta_\varphi^{t_{m-1}-t_m} \pi_\varphi(h) \cdots \Delta_\varphi^{t_1-t_2} \pi_\varphi(h) \Omega \\ & = \text{Exp}_r \left( \int_0^{\frac{1}{2}} ; \Delta_\varphi^t \pi_\varphi(h) \Delta_\varphi^{-t} dt \right) \Omega, \end{aligned} \quad (7.4)$$

where the sum is known to converge absolutely ([2]). The notation  $\text{Exp}_r$  is taken from [3].

The positive linear functional  $\omega^h$  on  $\mathfrak{M}$  is defined by

$$\omega^h(x) \equiv (\Omega^h, x\Omega^h) \quad (x \in \mathfrak{M}). \quad (7.5)$$

The vector  $\Omega^h$  defined above is cyclic and separating for  $\mathfrak{M}$ . Its modular automorphism group  $\sigma_t^{\omega^h}$  of  $\mathfrak{M}$  coincides with  $(\sigma_t^\omega)^h$ , i.e. the perturbed dynamics of  $(\sigma_t^\omega, \mathfrak{M})$  by  $h$ .  $\Omega^h$  is in the natural positive cone of  $(\Omega, \mathfrak{M})$  (see, e.g., [43] and [17]) for any self-adjoint element  $h \in \mathfrak{M}$  and satisfies

$$(\Omega^{h_1})^{h_2} = \Omega^{h_1+h_2} \quad (7.6)$$

for any self-adjoint elements  $h_1, h_2 \in \mathfrak{M}$ . We have

$$(\omega^{h_1})^{h_2} = \omega^{h_1+h_2}, \quad \sigma_t^{\{\omega^{(h_1+h_2)}\}} = (\sigma_t^\omega)^{(h_1+h_2)} = \{(\sigma_t^\omega)^{h_1}\}^{h_2}, \quad (7.7)$$

where  $\{(\sigma_t^\omega)^{h_1}\}^{h_2}$  indicates the dynamics which is given by the successive perturbations first by  $h_1$  and then by  $h_2$ . We denote the normalization of  $\omega^h$  by  $[\omega^h]$ :

$$[\omega^h] = \omega^h(\mathbf{1})^{-1} \omega^h = \omega^{(h - \{\log \omega^h(\mathbf{1})\} \mathbf{1})}. \quad (7.8)$$

We use the following estimates (Theorem 2 of [4]) and a formula (e.g. (3.5) of [7] and Theorem 3.10 of [9]) later.

$$\|\Omega^h\| \leq \exp \frac{1}{2} \|h\|, \quad \log \omega^h(\mathbf{1}) \leq \|h\|. \quad (7.9)$$

$$S(\varphi^h, \varphi) = -\varphi(h). \quad (7.10)$$

## 7.2 Surface Energy

Let us consider  $\Phi \in \mathcal{P}$ . For any finite subset  $I$  of  $\mathbb{Z}^\nu$ , we define

$$W(I) \equiv H(I) - U(I). \quad (7.11)$$

By (5.21), (5.22) and (5.23), the expression for  $W(I)$  in terms of the potential is given as follows.

$$\begin{aligned} W(I) &= \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \cap I^c \neq \emptyset\} \\ &\left( = \lim_{J \nearrow \mathbb{Z}^\nu} \left( \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \cap I^c \neq \emptyset, K \subset J\} \right) \right). \end{aligned} \quad (7.12)$$

$W(I)$  is the sum of all (interaction) potentials between the inside and the outside of  $I$  by definition, and will be called the surface energy.

### 7.3 Gibbs Condition

We are now in a position to introduce our Gibbs condition for a state  $\varphi$  of  $\mathcal{A}$  for a given  $\delta \in \Delta(\mathcal{A}_o)$ . We use the following notation in its definition below. As in § 6.1,  $\{\mathcal{H}_\varphi, \pi_\varphi, \Omega_\varphi\}$  is the GNS triplet for  $\varphi$ . The normal extension of  $\varphi$  to the weak closure  $\mathfrak{M}_\varphi (= \pi_\varphi(\mathcal{A})'')$  is denoted by the same letter  $\varphi$ :

$$\begin{aligned}\varphi(x) &= (\Omega_\varphi, x\Omega_\varphi) \quad (x \in \mathfrak{M}_\varphi), \\ \varphi(\pi_\varphi(a)) &= \varphi(a) \quad (a \in \mathcal{A}).\end{aligned}$$

Let  $\Phi(\mathbf{I})$ ,  $H(\mathbf{I})$ ,  $U(\mathbf{I})$  and  $W(\mathbf{I})$  be those uniquely associated with  $\delta$ . The following operators will be used for perturbations of dynamics and states

$$\hat{h} = \pi_\varphi(\beta H(\mathbf{I})), \quad \hat{u} = \pi_\varphi(\beta U(\mathbf{I})), \quad \hat{w} = \pi_\varphi(\beta W(\mathbf{I})). \quad (7.13)$$

**Definition 7.1.** For  $\delta \in \Delta(\mathcal{A}_o)$ , a state  $\varphi$  of  $\mathcal{A}$  is said to satisfy the  $(\delta, \beta)$ -Gibbs condition, or alternatively the  $(\Phi, \beta)$ -Gibbs condition, if the following two conditions are satisfied.

(D-1)  $\varphi$  is a modular state. (See Definition 6.2.)

(D-2) For each finite subset  $\mathbf{I}$  of  $\mathbb{Z}^\nu$ ,  $\sigma_t^{\varphi^{\hat{w}}}$  satisfies

$$\sigma_t^{\varphi^{\hat{w}}}(\pi_\varphi(A)) = \pi_\varphi\left(e^{-i\beta U(\mathbf{I})t} A e^{i\beta U(\mathbf{I})t}\right)$$

for all  $A \in \mathcal{A}(\mathbf{I})$ .

The condition (D-2) is equivalent to the following condition (D-2)' as shown in the subsequent Lemma and hence we may define the  $(\delta, \beta)$ -Gibbs condition by (D-1) and (D-2)'.

(D-2)' For each finite subset  $\mathbf{I}$  of  $\mathbb{Z}^\nu$  and  $A \in \mathcal{A}(\mathbf{I})$ ,  $\pi_\varphi(A)$  is  $\sigma_t^{\varphi^{\hat{h}}}$ -invariant, namely,  $\pi_\varphi(\mathcal{A}(\mathbf{I}))$  is in the centralizer of the positive linear functional  $\varphi^{\hat{h}}$ .

**Lemma 7.2.** The conditions (D-2) and (D-2)' are equivalent.

*Proof.* First assume (D-2). Since  $\hat{h} = \hat{w} + \hat{u}$ , we have  $\varphi^{\hat{h}} = (\varphi^{\hat{w}})^{\hat{u}}$  and hence

$$\begin{aligned}\sigma_t^{\varphi^{\hat{h}}} &= \left\{(\sigma_t^{\varphi^{\hat{w}}})^{\hat{u}}\right\} \\ &= \left(\sigma_t^{\varphi^{\hat{w}}}\right)^{\hat{u}}\end{aligned}$$

Since  $e^{-i\beta U(\mathbf{I})t} U(\mathbf{I}) e^{i\beta U(\mathbf{I})t} = U(\mathbf{I})$ ,  $\pi_\varphi(U(\mathbf{I}))$  is invariant under  $\sigma_t^{\varphi^{\hat{w}}}$  by (D-2). Then unitary cocycle bridging  $\sigma_t^{\varphi^{\hat{w}}}$  and  $\sigma_t^{\varphi^{\hat{h}}}$  becomes  $e^{i\hat{u}t}$ . Hence

$$\sigma_t^{\varphi^{\hat{h}}} = \text{Ad}(e^{i\hat{u}t}) \circ \sigma_t^{\varphi^{\hat{w}}}.$$

Therefore, for  $\pi_\varphi(A)$ ,  $A \in \mathcal{A}(\mathbf{I})$ , we have

$$\begin{aligned}\sigma_t^{\varphi^{\hat{h}}}(\pi_\varphi(A)) &= e^{i\hat{u}t} \sigma_t^{\varphi^{\hat{w}}}(\pi_\varphi(A)) e^{-i\hat{u}t} \\ &= \pi_\varphi(\text{Ad}(e^{i\beta U(\mathbf{I})t}) \circ \text{Ad}(e^{-i\beta U(\mathbf{I})t}) \circ A) \\ &= \pi_\varphi(A).\end{aligned}$$

Thus (D-2)' is satisfied.

We show the converse. Assume (D-2)'. Since  $\hat{w} = \hat{h} - \hat{u}$ ,  $\sigma_t^{\varphi^{\hat{w}}}$  is the perturbed dynamics of  $\sigma_t^{\varphi^{\hat{h}}}$  by  $-\hat{u}$ . Since  $u \in \mathcal{A}(\mathbf{I})$  is  $\sigma_t^{\varphi^{\hat{h}}}$ -invariant (being in the centralizer), the corresponding unitary cocycle is  $e^{-i\hat{u}t}$ . Hence, for  $\pi_\varphi(A)$ ,  $A \in \mathcal{A}(\mathbf{I})$ , we have

$$\begin{aligned}\sigma_t^{\varphi^{\hat{w}}}(\pi_\varphi(A)) &= e^{-i\hat{u}t} \sigma_t^{\varphi^{\hat{h}}}(\pi_\varphi(A)) e^{+i\hat{u}t} \\ &= e^{-i\beta\pi_\varphi(U(\mathbf{I}))t} \pi_\varphi(A) e^{i\beta\pi_\varphi(U(\mathbf{I}))t} \\ &= \pi_\varphi\left(e^{-i\beta U(\mathbf{I})t} A e^{i\beta U(\mathbf{I})t}\right),\end{aligned}$$

and (D-2) is derived.  $\square$

We introduce the local Gibbs state.

**Definition 7.3.** For finite  $\mathbf{I}$ , the local Gibbs state of  $\mathcal{A}(\mathbf{I})$  (or local Gibbs state for  $\mathbf{I}$ ) with respect to  $(\delta, \beta)$  is given by

$$\varphi_{\mathbf{I}}^c(A) \equiv \frac{\tau(e^{-\beta U(\mathbf{I})} A)}{\tau(e^{-\beta U(\mathbf{I})})}, \quad A \in \mathcal{A}(\mathbf{I}). \quad (7.14)$$

**Corollary 7.4.** If  $\varphi$  satisfies the  $(\delta, \beta)$ -Gibbs condition, then the restriction of  $\varphi^{\hat{h}}$  to  $\mathcal{A}(\mathbf{I})$  is  $\varphi^{\hat{h}}(\mathbf{1})$  times the tracial state  $\tau$  and that of  $\varphi^{\hat{w}}$  is  $\varphi^{\hat{w}}(\mathbf{1})$  times the local Gibbs state  $\varphi_{\mathbf{I}}^c$  given by (7.14).

*Proof.* Since  $\varphi^{\hat{h}}$  has the tracial property for  $\mathcal{A}(\mathbf{I})$  by (D-2)', its restriction to  $\mathcal{A}(\mathbf{I})$  must be  $\varphi^{\hat{h}}(\mathbf{1})$  times the unique tracial state  $\tau$ .

Since the inner automorphism group

$$\alpha_t^{\mathbf{I}} \equiv \text{Ad}(e^{-i\beta U(\mathbf{I})t}) \quad (7.15)$$

leaves  $\mathcal{A}(\mathbf{I})$  invariant and has the same action on  $\mathcal{A}(\mathbf{I})$  as the modular automorphism of  $\varphi^{\hat{w}}|_{\mathcal{A}(\mathbf{I})}$  (the restriction of  $\varphi^{\hat{w}}$  to  $\mathcal{A}(\mathbf{I})$ ),  $\varphi^{\hat{w}}|_{\mathcal{A}(\mathbf{I})}$  satisfies  $(\alpha_t^{\mathbf{I}}, -1)$  KMS condition and hence must be  $\varphi^{\hat{w}}(\mathbf{1})$  times the unique KMS state given by the local Gibbs state  $\varphi_{\mathbf{I}}^c$ .  $\square$

## 7.4 Equivalence to KMS Condition

**Theorem 7.5.** Let  $\alpha_t$  be dynamics of  $\mathcal{A}$  satisfying conditions (I) and (II) and  $\delta$  be the restriction of its generator  $\delta_\alpha$  to  $\mathcal{A}_0$ . Then any  $(\alpha_t, \beta)$ -KMS state  $\varphi$  of  $\mathcal{A}$  satisfies  $(\delta, \beta)$ -Gibbs condition.

*Proof.* As already indicated, it is known that the KMS condition implies (D-1). It remains to show (D-2). We have

$$(d/ds) \left( \sigma_s^{\varphi^{\hat{w}}}(x) - \sigma_s^\varphi(x) \right)_{s=0} = i[\hat{w}, x],$$

for  $x \in \mathfrak{M}_\varphi$ . By the group property of the automorphisms,

$$(d/dt) \sigma_t^{\varphi^{\hat{w}}}(x) = \sigma_t^{\varphi^{\hat{w}}} \left\{ (d/ds) \sigma_s^{\varphi^{\hat{w}}}(x) \Big|_{s=0} \right\}$$

for  $x$  in the domain of the generator of  $\sigma_t^{\varphi\hat{w}}$ . For the same  $x$ , we have

$$(d/dt)\sigma_t^{\varphi\hat{w}}(x) = \sigma_t^{\varphi\hat{w}} \left\{ (d/ds)\sigma_s^{\varphi}(x) \Big|_{s=0} + i[\hat{w}, x] \right\}.$$

The KMS condition implies that

$$\sigma_s^{\varphi}(\pi_{\varphi}(A)) = \pi_{\varphi}(\alpha_{-\beta s}(A)), \quad A \in \mathcal{A}.$$

Therefore, if  $A \in \mathcal{A}$  is in the domain of the generator of  $\alpha_t$ , we have

$$(d/dt)\sigma_t^{\varphi\hat{w}}(\pi_{\varphi}(A)) = \sigma_t^{\varphi\hat{w}} \left\{ (d/ds)(\pi_{\varphi}\{\alpha_{-\beta s}(A)\}) \Big|_{s=0} \right\} + \sigma_t^{\varphi\hat{w}}(\pi_{\varphi}\{[i\beta W(\mathbf{I}), A]\}).$$

Now we take  $A \in \mathcal{A}(\mathbf{I})$ . By (H-iii),

$$\begin{aligned} (d/dt)\sigma_t^{\varphi\hat{w}}(\pi_{\varphi}(A)) &= \sigma_t^{\varphi\hat{w}}(-i\beta\pi_{\varphi}\{[H(\mathbf{I}), A]\}) + \sigma_t^{\varphi\hat{w}}(i\beta\pi_{\varphi}\{[W(\mathbf{I}), A]\}) \\ &= -i\beta\sigma_t^{\varphi\hat{w}}(\pi_{\varphi}\{[U(\mathbf{I}), A]\}). \end{aligned}$$

For  $A \in \mathcal{A}(\mathbf{I})$ ,  $e^{i\beta U(\mathbf{I})t}Ae^{-i\beta U(\mathbf{I})t} \in \mathcal{A}(\mathbf{I})$ , and we have

$$\begin{aligned} &(d/dt)\sigma_t^{\varphi\hat{w}}(\pi_{\varphi}\{e^{i\beta U(\mathbf{I})t}Ae^{-i\beta U(\mathbf{I})t}\}) \\ &= \sigma_t^{\varphi\hat{w}} \left\{ (d/ds)\sigma_s^{\varphi\hat{w}}(\pi_{\varphi}\{e^{i\beta U(\mathbf{I})(t+s)}Ae^{-i\beta U(\mathbf{I})(t+s)}\}) \Big|_{s=0} \right\} \\ &= \sigma_t^{\varphi\hat{w}}(-i\beta\pi_{\varphi}\{[U(\mathbf{I}), e^{i\beta U(\mathbf{I})t}Ae^{-i\beta U(\mathbf{I})t}]\}) + \pi_{\varphi}\left\{ d/ds(e^{i\beta U(\mathbf{I})(t+s)}Ae^{-i\beta U(\mathbf{I})(t+s)}) \Big|_{s=0} \right\} \\ &= 0. \end{aligned}$$

This implies that

$$\sigma_t^{\varphi\hat{w}}(\pi_{\varphi}\{e^{i\beta U(\mathbf{I})t}Ae^{-i\beta U(\mathbf{I})t}\})$$

is a constant function of  $t$  and hence equals to its value at  $t = 0$ , which is  $\pi_{\varphi}(A)$ . Thus

$$\sigma_{-t}^{\varphi\hat{w}}(\pi_{\varphi}(A)) = \pi_{\varphi}(e^{i\beta U(\mathbf{I})t}Ae^{-i\beta U(\mathbf{I})t})$$

and (D-2) is shown.  $\square$

To show the converse, we need the assumption (III) for the dynamics  $\alpha_t$ .

**Theorem 7.6.** *Let  $\alpha_t$  be a dynamics of  $\mathcal{A}$  satisfying the conditions (I), (II) and (III). Let  $\delta$  be the restriction of its generator  $\delta_{\alpha}$  to  $\mathcal{A}_0$ . Then any  $(\delta, \beta)$ -Gibbs state  $\varphi$  of  $\mathcal{A}$  satisfies  $(\alpha_t, \beta)$ -KMS condition.*

*Proof.* We use (D-2)'. It says that

$$(d/dt)\sigma_t^{\varphi\hat{h}}(\pi_{\varphi}(A)) = 0$$

for all  $A \in \mathcal{A}(\mathbf{I})$ . By the group property of the automorphism,

$$(d/dt)\sigma_t^{\varphi}(x) = \sigma_t^{\varphi} \left\{ (d/ds)\sigma_s^{\varphi}(x) \Big|_{s=0} \right\}.$$



For any  $A \in \mathcal{A}_o$ , there exists a finite subset  $I$  such that  $A \in \mathcal{A}(I)$ . Since  $\varphi = (\varphi^{\hat{h}})^{-\hat{h}}$ , we have

$$\begin{aligned} (d/dt)\sigma_t^\varphi(\pi_\varphi(A)) &= \sigma_t^\varphi \left\{ (d/ds)\sigma_s^{\varphi^{\hat{h}}}(\pi_\varphi(A)) \Big|_{s=0} - [i\hat{h}, \pi_\varphi(A)] \right\} \\ &= \sigma_t^\varphi(-i\beta\pi_\varphi([H(I), A])) \\ &= -\beta\sigma_t^\varphi(\pi_\varphi(\delta A)). \end{aligned} \quad (7.16)$$

We note that for any  $A \in \mathcal{A}$

$$\sigma_t^\varphi(\pi_\varphi(A)) = \Delta_\varphi^{it}\pi_\varphi(A)\Delta_\varphi^{-it}, \quad \Delta_\varphi\Omega_\varphi = \Omega_\varphi.$$

By applying (7.16) on  $\Omega_\varphi$  and setting  $t = 0$ , we conclude that  $\pi_\varphi(A)\Omega_\varphi$  is in the domain of  $\log \Delta_\varphi$  and

$$i(\log \Delta_\varphi)\pi_\varphi(A)\Omega_\varphi = -\beta\pi_\varphi(\delta(A))\Omega_\varphi \quad (7.17)$$

for all  $A \in \mathcal{A}_o$ .

By Assumption (III), for every  $A \in D(\delta_\alpha)$ , there exists a sequence  $\{A_n\}$ ,  $A_n \in \mathcal{A}_o$  such that  $\{A_n\}$  and  $\{\delta A_n (= \delta_\alpha A_n)\}$  converge to  $A$  and  $\delta_\alpha A (= \bar{\delta} A)$ , respectively, in the norm topology of  $\mathcal{A}$ . Since  $\log \Delta_\varphi$  is a (self-adjoint) closed operator,  $\pi_\varphi(A)\Omega_\varphi$  must be in the domain of  $\log \Delta_\varphi$  and (7.17) holds for any  $A \in D(\delta_\alpha)$ .

For  $A \in D(\delta_\alpha)$  and  $t \in \mathbb{R}$ , we set

$$\xi_t \equiv \sigma_t^\varphi(\pi_\varphi\{\alpha_{\beta t}(A)\})\Omega_\varphi = \Delta_\varphi^{it}\pi_\varphi(\alpha_{\beta t}(A))\Omega_\varphi.$$

For  $A \in D(\delta_\alpha)$ ,  $\alpha_t(A)$  is in  $D(\delta_\alpha)$  for any  $t \in \mathbb{R}$ . Therefore, we can substitute  $\alpha_{\beta t}(A)$  into  $A$  of (7.17) and obtain

$$\begin{aligned} (d/dt)\xi_t &= \Delta_\varphi^{it} \left\{ (d/ds)\Delta_\varphi^{is}\pi_\varphi\{\alpha_{\beta t}(A)\}\Omega_\varphi \Big|_{s=0} \right\} + \Delta_\varphi^{it} \left\{ (d/dt)\pi_\varphi\{\alpha_{\beta t}(A)\}\Omega_\varphi \right\} \\ &= \Delta_\varphi^{it} \left\{ -\beta\pi_\varphi\{\delta(\alpha_{\beta t}(A))\}\Omega_\varphi + \pi_\varphi\{\beta\delta(\alpha_{\beta t}(A))\}\Omega_\varphi \right\} \\ &= 0. \end{aligned}$$

Therefore, we have  $\xi_t = \xi_0$  and

$$\sigma_t^\varphi(\pi_\varphi\{\alpha_{\beta t}(A)\})\Omega_\varphi = \pi_\varphi(A)\Omega_\varphi.$$

Since  $\Omega_\varphi$  is separating for  $\mathfrak{M}_\varphi$ , we obtain

$$\sigma_t^\varphi(\pi_\varphi\{\alpha_{\beta t}(A)\}) = \pi_\varphi(A).$$

This implies

$$\pi_\varphi\{\alpha_{\beta t}(A)\} = \sigma_{-t}^\varphi(\pi_\varphi(A)).$$

Since  $D(\delta_\alpha)(\supset \mathcal{A}_o)$  is norm dense in  $\mathcal{A}$ , we have

$$\pi_\varphi\{\alpha_{-\beta t}(A)\} = \sigma_t^\varphi(\pi_\varphi(A)),$$

for every  $A \in \mathcal{A}$ .

Since  $\varphi$  satisfies  $(\sigma_t^\varphi, -1)$ -KMS condition as a state of  $\mathfrak{M}_\varphi$ , we obtain the  $(\alpha_t, \beta)$ -KMS condition for  $\varphi$ .  $\square$

## 7.5 Product Form of the Gibbs Condition

In the case of quantum spin lattice systems, for any region  $I \subset \mathbb{Z}^\nu$ ,  $\mathcal{A} = \mathcal{A}(I) \otimes \mathcal{A}(I^c)$ . In this situation, the Gibbs condition implies that  $\varphi^{\hat{w}} (= \varphi^{\pi_\varphi(\beta W(I))})$  is a product of the local Gibbs state of  $\mathcal{A}(I)$  and its restriction to  $\mathcal{A}(I^c)$ , or equivalently  $\varphi^{\hat{h}} (= \varphi^{\pi_\varphi(\beta H(I))})$  is a product of the tracial state of  $\mathcal{A}(I)$  and its restriction to  $\mathcal{A}(I^c)$  for any finite region  $I$  [5].

However, this product property for  $\varphi^{\hat{w}}$  and  $\varphi^{\hat{h}}$  for the present Fermion case does not seem to be automatic in general. We show that such a product property holds if and only if the Gibbs state  $\varphi$  is  $\Theta$ -even, where the product property refers to the validity of the formula

$$\psi(AB) = \psi(A)\psi(B)/\psi(\mathbf{1}), \quad A \in \mathcal{A}(I), \quad B \in \mathcal{A}(I^c) \quad (7.18)$$

for  $\psi = \varphi^{\hat{h}}$  and for  $\psi = \varphi^{\hat{w}}$

**Proposition 7.7.** *Assume the conditions (I) and (II) for the dynamics. Let  $I$  be a non-empty finite subset of  $\mathbb{Z}^\nu$ . If  $\varphi$  satisfies the Gibbs condition, then  $\varphi^{\pi_\varphi(\beta W(I))}$  has the product property (7.18) if and only if  $\varphi$  is  $\Theta$ -even. The same is true for  $\varphi^{\pi_\varphi(\beta H(I))}$ .*

*Proof.* First assume that  $\varphi$  is even. It follows from the Gibbs condition that  $\mathcal{A}(I)$  is in the centralizer of  $\varphi^{\hat{h}}$  and the restriction of  $\varphi^{\hat{h}}$  to  $\mathcal{A}(I)$  is tracial. We will show

$$\varphi^{\hat{h}}([A_1, A_2]B) = 0 \quad (7.19)$$

for any  $A_1, A_2 \in \mathcal{A}(I)$  and any  $B \in \mathcal{A}(I^c)$ . It is enough to show this for all combinations of even and odd  $A_1, A_2$  and  $B$  because the general case follows from these cases by linearity.

Since  $A_1$  and  $A_2$  are in the centralizer of  $\varphi^{\hat{h}}$ , we have

$$\varphi^{\hat{h}}(A_1 A_2 B) = \varphi^{\hat{h}}(A_2 B A_1), \quad \varphi^{\hat{h}}(A_2 A_1 B) = \varphi^{\hat{h}}(A_1 B A_2).$$

If one, or more of  $A_1, A_2, B$  is even, then  $BA_1 = A_1 B$  or  $BA_2 = A_2 B$  holds. Hence (7.19) follows for this case.

The remaining case is when  $A_1, A_2, B$  are all odd. We now show that  $\varphi^{\hat{h}}$  is even so that (7.19) holds in this case.

Since  $\varphi$  is assumed to be even at this part of proof,  $\Theta$  leaves  $\varphi$  invariant and hence there exists an involutive unitary  $\mathcal{U}_\Theta$  on the GNS representation space  $\mathcal{H}_\varphi$  of  $\varphi$ , satisfying

$$\mathcal{U}_\Theta \pi_\varphi(A) \mathcal{U}_\Theta^* = \pi_\varphi(\Theta(A)), \quad (A \in \mathcal{A}), \quad (7.20)$$

$$\mathcal{U}_\Theta \Omega_\varphi = \Omega_\varphi. \quad (7.21)$$

Since  $H(I)$  is even by assumption, it follows from the commutativity of  $\mathcal{U}_\Theta$  with  $\Delta_\varphi$  [42] and the above equations (7.20), (7.21) that the perturbed vector  $\Omega_\varphi^{\hat{h}}$  is  $\mathcal{U}_\Theta$  invariant. Therefore  $\varphi^{\hat{h}}$  is even, since it is the vector functional by  $\Omega_\varphi^{\hat{h}}$ . Hence  $\varphi^{\hat{h}}$  vanishes on every odd element and (7.19) is satisfied if  $A_1, A_2$  and  $B$  are all odd. Now (7.19) is proved for all the cases.

Since  $\mathcal{A}(\mathbf{I})$  is a  $2^{|\mathbf{I}|} \times 2^{|\mathbf{I}|}$  full matrix algebra, any element  $A \in \mathcal{A}(\mathbf{I})$  can be written as

$$A = \tau(A) + \sum_j [A_{j1}, A_{j2}]$$

for some  $A_{j1}, A_{j2} \in \mathcal{A}(\mathbf{I})$ . Hence (7.19) implies

$$\varphi^{\hat{h}}(AB) = \tau(A)\varphi^{\hat{h}}(B) \quad (7.22)$$

for any  $A \in \mathcal{A}(\mathbf{I})$  and  $B \in \mathcal{A}(\mathbf{I}^c)$ . This means that  $\varphi^{\hat{h}}$  has a form of the product of  $\tau$  of  $\mathcal{A}(\mathbf{I})$  and its restriction to  $\mathcal{A}(\mathbf{I}^c)$ .

Since  $U(\mathbf{I})$  is in the centralizer of  $\varphi^{\hat{h}}$ , we have

$$\varphi^{\hat{w}} = \{\varphi^{\hat{h}}\}^{-\hat{u}} = \varphi^{\hat{h}} \cdot e^{-\hat{u}}.$$

Hence, for any  $A \in \mathcal{A}(\mathbf{I})$  and  $B \in \mathcal{A}(\mathbf{I}^c)$ ,

$$\varphi^{\hat{w}}(AB) = \tau(e^{-\hat{u}})\varphi_1^c(A)\varphi^{\hat{h}}(B).$$

By setting  $A = \mathbf{1}$ , we have

$$\varphi^{\hat{w}}(B) = \tau(e^{-\hat{u}})\varphi^{\hat{h}}(B).$$

Therefore

$$\varphi^{\hat{w}}(AB) = \varphi_1^c(A)\varphi^{\hat{w}}(B). \quad (7.23)$$

Hence we have the desired product property of  $\varphi^{\hat{w}}$ .

We now prove the converse, starting from the assumption that  $\varphi^{\hat{h}}$  has a product form (7.18).

We note that

$$\tau(a_i a_i^*) = \tau(a_i^* a_i) = \tau\left(\frac{1}{2}(a_i a_i^* + a_i^* a_i)\right) = \tau\left(\frac{1}{2}\mathbf{1}\right) = \frac{1}{2}$$

due to CAR. On the other hand,  $a_i$  anticommutes with any odd element  $B$  in  $\mathcal{A}(\mathbf{I}^c)$  and hence

$$\varphi^{\hat{h}}(a_i a_i^* B) = \varphi^{\hat{h}}(a_i^* B a_i) = -\varphi^{\hat{h}}(a_i^* a_i B), \quad (7.24)$$

where the first equality follows because  $a_i$  is in the centralizer of  $\varphi^{\hat{h}}$  due to the Gibbs condition. By the product form assumption,

$$\varphi^{\hat{h}}(AB) = \varphi^{\hat{h}}(A)\varphi^{\hat{h}}(B)/\varphi^{\hat{h}}(\mathbf{1})$$

for  $A \in \mathcal{A}(\mathbf{I})$  and  $B \in \mathcal{A}(\mathbf{I}^c)$ . Since  $A$  is in the centralizer,  $\varphi^{\hat{h}}(A)/\varphi^{\hat{h}}(\mathbf{1}) = \tau(A)$  for the unique tracial state  $\tau$  of  $\mathcal{A}(\mathbf{I})$ . Hence

$$\begin{aligned} \varphi^{\hat{h}}(a_i a_i^* B) &= \tau(a_i a_i^*)\varphi^{\hat{h}}(B) = \frac{1}{2}\varphi^{\hat{h}}(B), \\ \varphi^{\hat{h}}(a_i^* a_i B) &= \tau(a_i^* a_i)\varphi^{\hat{h}}(B) = \frac{1}{2}\varphi^{\hat{h}}(B). \end{aligned} \quad (7.25)$$

From (7.24) and (7.25), we obtain

$$\varphi^{\hat{h}}(B) = 0 \quad (7.26)$$

for any  $B \in \mathcal{A}(\mathbf{I}^c)_-$ . Since  $\mathcal{A}_- = \mathcal{A}(\mathbf{I})_+ \mathcal{A}(\mathbf{I}^c)_- + \mathcal{A}(\mathbf{I})_- \mathcal{A}(\mathbf{I}^c)_+$  for a finite  $\mathbf{I}$ ,  $\varphi^{\hat{h}}$  vanishes on odd elements of  $\mathcal{A}$ . We conclude that  $\varphi^{\hat{h}}$  is even. This implies that  $\varphi$  is also even by the same argument as in the first part of this proof due to  $\varphi = \{\varphi^{\hat{h}}\}^{-\hat{h}}$ .  $\square$

*Remark.* By the above Proposition, we have already shown that if a Gibbs state  $\varphi$  satisfies the condition that  $\varphi^{\pi_{\varphi}(\beta W(\mathbf{I}))}$  has the product property (7.18) for the pair  $(\mathcal{A}(\mathbf{I}), \mathcal{A}(\mathbf{I}^c))$  for one non-empty finite  $\mathbf{I}$ , then  $\varphi$  has this product property for every finite subset  $\mathbf{I}$ .

In connection with Proposition 7.7, if  $\mathcal{A}(\mathbf{I}^c)$  is replaced by the commutant algebra  $\mathcal{A}(\mathbf{I})'$  in the product property (7.18), then  $\varphi^{\hat{w}}$  is a product of the local Gibbs state of  $\mathcal{A}(\mathbf{I})$  and its restriction to  $\mathcal{A}(\mathbf{I})'$  for every finite region  $\mathbf{I}$  irrespective of whether  $\varphi$  is even or not as is shown in the following corollary. This situation is much the same as in quantum spin lattice systems.

**Corollary 7.8.** *Assume the conditions (I) and (II) for the dynamics. Let  $\varphi$  be a modular state. The state  $\varphi$  satisfies the Gibbs condition if and only if the perturbed functional  $\varphi^{\hat{w}}$  is a product of the local Gibbs state  $\varphi_{\mathbf{I}}^c$  of  $\mathcal{A}(\mathbf{I})$  and its restriction to  $\mathcal{A}(\mathbf{I})'$  for every finite  $\mathbf{I}$ .*

*Proof.* For a finite  $\mathbf{I}$ ,  $\mathcal{A}(\mathbf{I})$  is a full matrix algebra and hence  $\mathcal{A}$  is an (algebraic) tensor product of  $\mathcal{A}(\mathbf{I})$  and  $\mathcal{A}(\mathbf{I})'$ .

If  $\varphi^{\hat{w}}$  has the product property described above, then the GNS representation of  $\mathcal{A}$  associated with  $\varphi^{\hat{w}}$  is the tensor product of those for  $(\mathcal{A}(\mathbf{I}), \varphi_{\mathbf{I}}^c)$  and  $(\mathcal{A}(\mathbf{I})', \psi)$  where  $\psi = \varphi^{\hat{w}}|_{\mathcal{A}(\mathbf{I})'}$ . Therefore the product of the modular automorphisms for these two pairs satisfies the KMS condition (with  $\beta = -1$ ) for  $(\mathcal{A}, \varphi^{\hat{w}})$  and must be the modular operator for  $(\mathcal{A}, \varphi^{\hat{w}})$ . In particular, the restriction of the modular automorphisms of  $(\mathcal{A}, \varphi^{\hat{w}})$  to  $\mathcal{A}(\mathbf{I})$  coincides with the modular automorphisms  $\alpha_t^{\mathbf{I}} (= \text{Ad}(e^{-i\beta U(\mathbf{I})t}))$  for  $(\mathcal{A}(\mathbf{I}), \varphi_{\mathbf{I}}^c)$ . Hence the Gibbs condition is satisfied.

Conversely, assume that the Gibbs condition is satisfied for  $\varphi$ . By the elementwise commutativity of  $\mathcal{A}(\mathbf{I})$  and  $\mathcal{A}(\mathbf{I})'$ , we can show directly (7.19) in Proposition 7.7 in this case for any  $A_1, A_2 \in \mathcal{A}(\mathbf{I})$  and  $B \in \mathcal{A}(\mathbf{I})'$  skipping the previous discussion about even and odd elements. The argument showing (7.22) and (7.23) are still valid after we replace  $\mathcal{A}(\mathbf{I}^c)$  by  $\mathcal{A}(\mathbf{I})'$ .  $\square$

## 8 Translation Invariant Dynamics

### 8.1 Translation Invariance and Covariance

From now on, we need the following assumption for the dynamics  $\alpha_t$  for the most part of our theory.

(IV)  $\alpha_t \tau_k = \tau_k \alpha_t$  for all  $t \in \mathbb{R}$  and  $k \in \mathbb{Z}^\nu$ .

If (IV) holds,  $\alpha_t$  is said to be translation invariant. This assumption implies our earlier assumption (I) due to the following Proposition, which we owe to a referee.

**Proposition 8.1.** *Any automorphism  $\alpha_t$  commuting with the lattice translation  $\tau_k$ ,  $k \in \mathbb{Z}^\nu$ , must commute with  $\Theta$ .*

For its proof, we need the following Lemma.

**Lemma 8.2.** *An element  $x \in \mathcal{A}$  is  $\Theta$ -even if and only if the following asymptotically central property holds.*

$$\lim_{k \rightarrow \infty} \|[\tau_k(x), y]\| = 0 \quad (8.1)$$

for all  $y \in \mathcal{A}$ .

*Proof.* If  $x \in (\mathcal{A}_\circ)_+$  and  $y \in \mathcal{A}_\circ$ , then  $[\tau_k(x), y] = 0$  for sufficiently large  $k$ . By the density of  $(\mathcal{A}_\circ)_+$  in  $\mathcal{A}_+$  and  $\mathcal{A}_\circ$  in  $\mathcal{A}$ , we obtain (8.1) for  $x \in \mathcal{A}_+$  and  $y \in \mathcal{A}$ .

In the converse direction, consider a general  $x \in \mathcal{A}$  and define  $x_\pm = 1/2(x \pm \Theta(x)) \in \mathcal{A}_\pm$ . Due to the validity of (8.1) for  $x_+$ , which is just shown, we have

$$\lim_{k \rightarrow \infty} \|[\tau_k(x), y]\| = \lim_{k \rightarrow \infty} \|[\tau_k(x_-), y]\|.$$

Take a unitary  $y \in \mathcal{A}_-$  (e.g.,  $a_i + a_i^*$ ). Then

$$\|[\tau_k(x_-), y]\| = 2\|\tau_k(x_-)y\| = 2\|x_-\|.$$

Hence (8.1) for  $x$  implies  $x_- = 0$ , namely  $x \in \mathcal{A}_+$ . □

*Proof of Proposition 8.1 :*

Due to  $\tau_k \alpha = \alpha \tau_k$ , we have

$$\|[\tau_k(\alpha(x)), \alpha(y)]\| = \|\alpha\{[\tau_k(x), y]\}\| = \|[\tau_k(x), y]\|.$$

Hence  $\alpha(x) \in \mathcal{A}_+$  if and only if  $x \in \mathcal{A}_+$  by Lemma 8.2. Let

$$E_+ \equiv \frac{1}{2}(\text{id} + \Theta). \quad (8.2)$$

It is the conditional expectation from  $\mathcal{A}$  onto  $\mathcal{A}_+$ , characterized by  $E_+(x) \in \mathcal{A}_+$  for all  $x \in \mathcal{A}$  and  $\tau(xy) = \tau(E_+(x)y)$  for all  $x \in \mathcal{A}$  and  $y \in \mathcal{A}_+$ . Then  $\alpha(\alpha^{-1}(y)) = y \in \mathcal{A}_+$  implies  $\alpha^{-1}(y) \in \mathcal{A}_+$  and

$$\begin{aligned} \tau(E_+(\alpha(x))y) &= \tau(\alpha(x)y) = \tau(\alpha(x\alpha^{-1}(y))) = \tau(x\alpha^{-1}(y)) \\ &= \tau(E_+(x)\alpha^{-1}(y)) = \tau(\alpha^{-1}\{\alpha(E_+(x))y\}) = \tau(\alpha(E_+(x))y), \end{aligned}$$

where we have used  $\alpha^{-1}(y) \in \mathcal{A}_+$  in the fourth equality. Since  $E_+(\alpha(x)) \in \mathcal{A}_+$  and  $\alpha(E_+(x)) \in \mathcal{A}_+$  (due to  $E_+(x) \in \mathcal{A}_+$ ), we have  $E_+(\alpha(x)) = \alpha(E_+(x))$ . Therefore  $E_+\alpha = \alpha E_+$  and  $\alpha$  commutes with  $\Theta$ . □

*Remark.* A referee pointed out the following approach (which we have not adopted). Under assumption IV, any  $\alpha_t|_{\mathcal{A}_+}$ -KMS state of  $\mathcal{A}_+$  has a unique even extension to an  $\alpha_t$ -KMS state of  $\mathcal{A}$  (e.g. by [11]). This allows one to reduce the analysis of KMS states to the case of asymptotically abelian system due to (8.1).

The dynamics  $\alpha_t$  is translation invariant if and only if its generator  $\alpha_t$  commutes with every  $\tau_k$  ( $k \in \mathbb{Z}^\nu$ ). (This statement includes the  $\tau_k$ -invariance of the domain of the generator.)

The corresponding standard potential (which exists under the assumptions (I) and (II)) satisfies the following translation covariance condition:

$$(\Phi\text{-f}) \quad \tau_k \Phi(I) = \Phi(I + k), \text{ for all finite subsets } I \text{ of } \mathbb{Z}^\nu \text{ and all } k \in \mathbb{Z}^\nu.$$

Such a potential will be said to be translation covariant.

We consider the set  $\mathcal{P}_\tau$  of all translation covariant potentials in  $\mathcal{P}$ . Namely,  $\mathcal{P}_\tau$  is defined to be the set of all  $\Phi$  satisfying all conditions of Definition 5.10, i.e., ( $\Phi$ -a,b,c,d,e) and the translation covariance ( $\Phi$ -f).

We make  $\mathcal{P}_\tau$  a real vector space as a function space on the set of finite subsets of  $\mathbb{Z}^\nu$  by the linear operation given in (5.24).

In the same way, we define  $\mathbf{H}_\tau$  to be the subspace of  $\mathbf{H}$  such that each element  $H$  satisfies the following translation covariance condition:

$$(H\text{-vi}) \quad \tau_k(H(I)) = H(I + k) \text{ for all } k \in \mathbb{Z}^\nu.$$

We denote the set of all translation invariant derivations in  $\Delta(\mathcal{A}_0)$  by  $\Delta_\tau(\mathcal{A}_0)$ . Namely,  $\Delta_\tau(\mathcal{A}_0)$  is the set of all  $*$ -derivations with  $\mathcal{A}_0$  as their domain, commuting with  $\Theta$  and also with  $\tau$ .

From Theorems 5.7, 5.12 and 5.13, the following corollaries obviously follow.

**Corollary 8.3.** *The relation (H-iii) (as given in § 5.2) between  $H \in \mathbf{H}_\tau$  and  $\delta \in \Delta_\tau(\mathcal{A}_0)$  gives a bijective, real linear map from  $\mathbf{H}_\tau$  to  $\Delta_\tau(\mathcal{A}_0)$ .*

**Corollary 8.4.** *The equations (5.22) and (5.23) for  $\Phi \in \mathcal{P}_\tau$  and  $H \in \mathbf{H}_\tau$  give a bijective, real linear map from  $\mathcal{P}_\tau$  to  $\mathbf{H}_\tau$ .*

**Corollary 8.5.** *The equations (5.27) and (5.28) between  $\Phi \in \mathcal{P}_\tau$  and  $\delta_\Phi \in \Delta_\tau(\mathcal{A}_0)$  gives a bijective, real linear map from  $\mathcal{P}_\tau$  to  $\Delta_\tau(\mathcal{A}_0)$ .*

For  $\Phi \in \mathcal{P}_\tau$ , we define

$$\|\Phi\| \equiv \|H(\{n\})\|$$

which is independent of  $n \in \mathbb{Z}^\nu$  due to the translation covariance of  $\Phi$ . It defines a norm on  $\mathcal{P}_\tau$ . We show that this norm makes  $\mathcal{P}_\tau$  a Banach space, after giving the following energy estimates.

**Lemma 8.6.** *For  $\Phi \in \mathcal{P}_\tau$ , the following estimate hold :*

$$\|U(I)\| \leq \|H(I)\| \leq \|\Phi\| \cdot |I|, \quad (8.3)$$

*In particular, if  $\|\Phi\| = 0$ ,  $H = U = \Phi = 0$  (as functions of finite subsets  $I$  of  $\mathbb{Z}^\nu$ ).*

*Proof.* For  $I = \emptyset$ , both sides of the above inequalities are 0.

For  $I = \{n_1, \dots, n_{|I|}\}$ , we obtain

$$\begin{aligned}
H(I) &= \lim_{J \nearrow \mathbb{Z}^\nu} \sum_K \left\{ \Phi(K); K \cap I \neq \emptyset, K \subset J \right\} \\
&= \lim_{J \nearrow \mathbb{Z}^\nu} \sum_{i=1}^{|I|} \sum_K \left\{ \Phi(K); K \ni n_i, K \not\ni n_1, \dots, n_{i-1}, K \subset J \right\} \\
&= \lim_{J \nearrow \mathbb{Z}^\nu} \sum_{i=1}^{|I|} E_{\{n_1, \dots, n_{i-1}\}^c} \sum_K \left\{ \Phi(K); K \ni n_i, K \subset J \right\} \\
&= \sum_{i=1}^{|I|} E_{\{n_1, \dots, n_{i-1}\}^c} H(\{n_i\}),
\end{aligned}$$

where the third equality comes from the following identities

$$E_{\{n_1, \dots, n_{i-1}\}^c} \Phi(K) = \begin{cases} 0 & \text{if } \{n_1, \dots, n_{i-1}\} \cap K \neq \emptyset, \text{ i.e. } \{n_1, \dots, n_{i-1}\}^c \not\supset K \\ \Phi(K) & \text{if } n_1, \dots, n_{i-1} \notin K, \text{ i.e. } \{n_1, \dots, n_{i-1}\}^c \supset K, \end{cases}$$

and the interchange of  $\lim_{J \nearrow \mathbb{Z}^\nu}$  and  $E_{\{n_1, \dots, n_{i-1}\}^c}$  in the fourth equality is allowed due to  $\|E_{\{n_1, \dots, n_{i-1}\}^c}\| = 1$ .

The following estimate follows:

$$\begin{aligned}
\|H(I)\| &\leq \sum_{i=1}^{|I|} \|E_{\{n_1, \dots, n_{i-1}\}^c} H(\{n_i\})\| \\
&\leq \sum_{i=1}^{|I|} \|H(\{n_i\})\| = |I| \cdot \|\Phi\|.
\end{aligned} \tag{8.4}$$

Since  $U(I) = E_I(H(I))$  and  $\|E_I\| = 1$ , we obtain

$$\|U(I)\| \leq \|H(I)\| \leq \|\Phi\| \cdot |I|.$$

If  $\|\Phi\| = 0$ , then  $H(I) = U(I) = 0$  for all  $I$  by this estimate and hence  $\Phi(I) = 0$  by (5.16).  $\square$

The following estimate will be used later.

**Lemma 8.7.** *For disjoint finite subsets  $I$  and  $J$  of  $\mathbb{Z}^\nu$ ,*

$$\|U(I \cup J) - U(I)\| \leq \|\Phi\| \cdot |J|. \tag{8.5}$$

*Proof.* Due to  $I \cap J = \emptyset$ ,

$$U(I \cup J) - U(I) = \left\{ \Phi(K); K \cap J \neq \emptyset, K \subset I \cup J \right\}.$$

Therefore, we have

$$U(I \cup J) - U(I) = E_{I \cup J} H(J),$$

because  $H(J)$  is the sum of  $\Phi(K)$  for all  $K$  satisfying  $K \cap J \neq \emptyset$ , and  $E_{I \cup J}$  annihilates all  $\Phi(K)$  for which  $K$  is not contained in  $I \cup J$  while it retains  $\Phi(K)$  unchanged if  $K$  is contained in  $I \cup J$ . Hence

$$\|U(I \cup J) - U(I)\| = \|E_{I \cup J}H(J)\| \leq \|H(J)\| \leq \|\Phi\| \cdot |J|.$$

□

**Proposition 8.8.**  $\mathcal{P}_\tau$  is a real Banach space with respect to the norm  $\|\Phi\| = \|H(\{n\})\|$ .

*Proof.*  $\mathcal{P}_\tau$  is a normed space with respect to  $\|\Phi\|$ , because

$$\begin{aligned} \|\Phi_1 + \Phi_2\| &= \|H_{\Phi_1 + \Phi_2}(\{n\})\| \\ &= \|H_{\Phi_1}(\{n\}) + H_{\Phi_2}(\{n\})\| \\ &\leq \|H_{\Phi_1}(\{n\})\| + \|H_{\Phi_2}(\{n\})\| \\ &= \|\Phi_1\| + \|\Phi_2\| \end{aligned}$$

$$\begin{aligned} \|c\Phi\| &= \|cH_\Phi(\{n\})\| \\ &= |c| \|H_\Phi(\{n\})\| = |c| \|\Phi\|, \end{aligned}$$

for  $\Phi_1, \Phi_2, \Phi \in \mathcal{P}_\tau$ , and  $c \in \mathbb{R}$ , due to the linear dependence of  $H_\Phi$  on  $\Phi$  and because  $\|\Phi\| = 0$  implies  $\Phi(I) = 0$  for all  $I$  due to Lemma 8.6 and (5.16).

We now show its completeness. Suppose  $\{\Phi_n\}$  is a Cauchy sequence in  $\mathcal{P}_\tau$  with respect to the norm  $\|\cdot\|$ . Let us denote the corresponding  $H(I)$  and  $U(I)$  for  $\Phi_n$  by  $H_n(I)$  and  $U_n(I)$ , respectively. The linear dependence of  $H(I)$  on  $\Phi$  and Lemma 8.6 imply that  $\{H_n(I)\}$  is a Cauchy sequence in  $\mathcal{A}$  with respect to the  $\mathbf{C}^*$ -norm. Since  $\mathcal{A}$  is a  $\mathbf{C}^*$ -algebra,  $\{H_n(I)\}$  has a unique limit in  $\mathcal{A}$ , which will be denoted by  $H_\infty(I)$ .

Since  $U(I) = E_I(H(I))$  with  $\|E_I\| = 1$ ,  $\{U_n(I)\}$  is also a Cauchy sequence in  $\mathcal{A}$ , has a unique limit  $U_\infty(I)$ , and  $U_\infty(I) = E_I(H_\infty(I))$ .

For each finite subset  $I$  of  $\mathbb{Z}^\nu$ ,  $\{\Phi_n(I)\}$  also converges to the potential  $\Phi_\infty(I)$  for  $U_\infty(I)$  in the  $\mathbf{C}^*$ -norm because  $\Phi(I)$  is a finite linear combination of  $U(J)$ ,  $J \subset I$  due to (5.16), and  $\{U_n(J)\}$  converges to  $U_\infty(J)$  in the  $\mathbf{C}^*$ -norm for every such  $J$ . For any finite subsets  $I, J$  of  $\mathbb{Z}^\nu$ , we obtain

$$\begin{aligned} &\sum_K \{\Phi_\infty(K); K \cap I \neq \emptyset, K \subset J\} \\ &= \sum_K \lim_n \{\Phi_n(K); K \cap I \neq \emptyset, K \subset J\} = \lim_n \sum_K \{\Phi_n(K); K \cap I \neq \emptyset, K \subset J\} \\ &= \lim_n E_J(H_n(I)) = E_J\left(\lim_n H_n(I)\right) \\ &= E_J(H_\infty(I)), \end{aligned}$$

where the third equality is due to (5.20). Hence, by (4.23) we have

$$\lim_{J \nearrow \mathbb{Z}^\nu} \left( \sum_K \{\Phi_\infty(K); K \cap I \neq \emptyset, K \subset J\} \right) = \lim_{J \nearrow \mathbb{Z}^\nu} E_J(H_\infty(I)) = H_\infty(I).$$



Thus  $\Phi_\infty$  satisfies the condition ( $\Phi$ -e) in the definition of  $\mathcal{P}_\tau$ . The other conditions ( $\Phi$ -a), ( $\Phi$ -b), ( $\Phi$ -c), ( $\Phi$ -d), and ( $\Phi$ -f) are satisfied since each  $\Phi_n$  satisfies them and  $\lim_n \Phi_n(I) = \Phi_\infty(I)$  for every finite subset  $I$  of  $\mathbb{Z}^\nu$ . In conclusion, we have  $\Phi_\infty \in \mathcal{P}_\tau$ .

Finally, we have

$$\lim_n \|\Phi_n - \Phi_\infty\| = \lim_n \|H_n(\{0\}) - H_\infty(\{0\})\| = 0.$$

We have now shown the completeness of  $\mathcal{P}_\tau$ .  $\square$

## 8.2 Finite Range Potentials

**Definition 8.9.** (1) A potential  $\Phi \in \mathcal{P}_\tau$  is said to be of a finite range if there exists an  $r \geq 0$  such that  $\Phi(I) = 0$  whenever

$$\text{diam}(I) = \max\{|i - j|; i, j \in I\} > r. \quad (8.6)$$

The infimum of such  $r$  is called the range of  $\Phi$ .

(2) The subspace of  $\mathcal{P}$  consisting of all potentials  $\Phi \in \mathcal{P}$  of a finite range is denoted by  $\mathcal{P}^f$ . Furthermore, we denote

$$\mathcal{P}_\tau^f \equiv \mathcal{P}^f \cap \mathcal{P}_\tau. \quad (8.7)$$

For  $a \in \mathbb{N}$ ,  $\mathcal{C}_a$  denotes the following cube in  $\mathbb{Z}^\nu$

$$\mathcal{C}_a \equiv \{x \in \mathbb{Z}^\nu; 0 \leq x_i \leq a - 1, i = 1, \dots, \nu\}. \quad (8.8)$$

We introduce the following averaged conditional expectation.

$$E_a \equiv \frac{1}{|\mathcal{C}_a|} \sum_{i \in \mathcal{C}_a} E_{\mathcal{C}_a - i}, \quad (8.9)$$

where  $|\mathcal{C}_a| = a^\nu$  is the number of lattice points in  $\mathcal{C}_a$ , called the volume of  $\mathcal{C}_a$ . (The sum in the above equation is over all translates of  $\mathcal{C}_a$  which contain the origin  $0 \in \mathbb{Z}^\nu$ .)

For any finite subset  $I \subset \mathbb{Z}^\nu$ ,  $l(a, I)$  denotes the number of translates of  $\mathcal{C}_a$  containing  $I$ . By definition, for any  $m \in \mathbb{Z}^\nu$ ,

$$l(a, I) = l(a, I + m) \quad (8.10)$$

We need the following lemma in this subsection and later.

**Lemma 8.10.** For a finite  $I$ ,

$$\lim_{a \rightarrow \infty} \frac{l(a, I)}{|\mathcal{C}_a|} = 1 \quad (8.11)$$

*Proof.* Let  $d \in \mathbb{N}$  be fixed such that there exists a translate  $\mathcal{C}_d + k$  ( $k \in \mathbb{Z}^\nu$ ) of  $\mathcal{C}_d$  containing  $I$ . For  $a > d$ , a translate of  $\mathcal{C}_a$  contains  $I$  if it contains  $\mathcal{C}_d + k$ .

Hence  $l(a, \mathbf{I})$  is bigger than the number of translates of  $\mathcal{C}_a$  which contains  $\mathcal{C}_d$ , which is  $(a - d + 1)^\nu$ . Hence

$$1 \geq \frac{l(a, \mathbf{I})}{|\mathcal{C}_a|} \geq \frac{(a - d + 1)^\nu}{|\mathcal{C}_a|} = \left(1 - \frac{(d - 1)}{a}\right)^\nu \rightarrow 1 \quad (a \rightarrow \infty).$$

This shows (8.11).  $\square$

In order to prove that the subspace  $\mathcal{P}_\tau^f$  is dense in  $\mathcal{P}_\tau$ , we need the following Lemma.

**Lemma 8.11.** *For any  $A \in \mathcal{A}$ ,*

$$\lim_{a \rightarrow \infty} E_a(A) = A. \quad (8.12)$$

*Proof.* Since  $\mathcal{A}_0$  is dense in  $\mathcal{A}$ , there exists  $A_\varepsilon \in \mathcal{A}_0$  for any  $\varepsilon > 0$  such that

$$\|A_\varepsilon - A\| < \varepsilon. \quad (8.13)$$

Let  $A_\varepsilon \in \mathcal{A}(\mathbf{I}_\varepsilon)$  for a finite  $\mathbf{I}_\varepsilon$ . Then there exists a sufficiently large positive integer  $b$  such that a translate of  $\mathcal{C}_b$ , say  $\mathcal{C}_b - k$ , contains both 0 (the origin of  $\mathbb{Z}^\nu$ ) and  $\mathbf{I}_\varepsilon$ . If a translate  $\mathcal{C}_a - i$  of  $\mathcal{C}_a$  contains  $\mathcal{C}_b - k$ , then  $E_{\mathcal{C}_a - i}(A_\varepsilon) = A_\varepsilon$  because  $\mathcal{C}_a - i \supset \mathcal{C}_b - k \supset \mathbf{I}_\varepsilon$  and  $A_\varepsilon \in \mathcal{A}(\mathbf{I}_\varepsilon)$ . Such  $i$  belongs to  $\mathcal{C}_a$  due to  $0 \in \mathcal{C}_b - k \subset \mathcal{C}_a - i$ . The number of translates  $\mathcal{C}_a - i$  of  $\mathcal{C}_a$  which contains  $\mathcal{C}_b - k$  is equal to  $l(a, \mathcal{C}_b)$  (the number of translates of  $\mathcal{C}_a$  which contains  $\mathcal{C}_b$ ). Therefore, we obtain

$$\begin{aligned} & \|A_\varepsilon - E_a(A_\varepsilon)\| \\ = & \left\| \left(1 - \frac{l(a, \mathcal{C}_b)}{|\mathcal{C}_a|}\right) A_\varepsilon - \frac{1}{|\mathcal{C}_a|} \sum \{E_{\mathcal{C}_a - i}(A_\varepsilon); i \in \mathcal{C}_a, \mathcal{C}_a - i \not\supset \mathcal{C}_b - k\} \right\|. \end{aligned}$$

Hence, by using  $\|E_{\mathcal{C}_a - i}(A_\varepsilon)\| \leq \|A_\varepsilon\|$  due to  $\|E_{\mathcal{C}_a - i}\| = 1$ , we obtain

$$\begin{aligned} \|A_\varepsilon - E_a(A_\varepsilon)\| & \leq \left( \left\{1 - \frac{l(a, \mathcal{C}_b)}{|\mathcal{C}_a|}\right\} + \frac{1}{|\mathcal{C}_a|} \{|\mathcal{C}_a| - l(a, \mathcal{C}_b)\} \right) \|A_\varepsilon\| \\ & = 2 \left(1 - \frac{l(a, \mathcal{C}_b)}{|\mathcal{C}_a|}\right) \|A_\varepsilon\|. \end{aligned}$$

By Lemma 8.10

$$\lim_{a \rightarrow \infty} \frac{l(a, \mathcal{C}_b)}{|\mathcal{C}_a|} = 1.$$

Hence, there exists  $n_\varepsilon \in \mathbb{N}$  such that for  $a \geq n_\varepsilon$ ,

$$\|A_\varepsilon - E_a(A_\varepsilon)\| < \varepsilon. \quad (8.14)$$

Hence, for  $a \geq n_\varepsilon$ ,

$$\begin{aligned} \|A - E_a(A)\| & \leq \|A - A_\varepsilon\| + \|A_\varepsilon - E_a(A_\varepsilon)\| + \|E_a(A_\varepsilon - A)\| \\ & < 3\varepsilon \end{aligned}$$

by (8.13), (8.14) and  $\|E_a\| = 1$ .  $\square$

**Theorem 8.12.**  $\mathcal{P}_\tau^f$  is dense in  $\mathcal{P}_\tau$ .

*Proof.* Let  $\Phi \in \mathcal{P}_\tau$ . For any finite  $I \subset \mathbb{Z}^\nu$  containing the origin 0 of  $\mathbb{Z}^\nu$ ,

$$E_a(\Phi(I)) = \frac{l(a, I)}{|\mathcal{C}_a|} \Phi(I), \quad (8.15)$$

because  $E_{\mathcal{C}_a-i}(\Phi(I)) = \Phi(I)$  if  $\mathcal{C}_a - i$  contains  $I$  while  $E_{\mathcal{C}_a-i}(\Phi(I)) = 0$  if  $\mathcal{C}_a - i$  does not contain  $I$  due to  $(\Phi-d)$ . Note that all translates of  $\mathcal{C}_a$  which contains  $I$  appear in the sum (8.9) since  $I$  is assumed to contain 0.

We now consider the following potential

$$\Phi_a(I) = \frac{l(a, I)}{|\mathcal{C}_a|} \Phi(I). \quad (8.16)$$

Due to  $\Phi \in \mathcal{P}_\tau$ ,  $(\Phi-a)$ ,  $(\Phi-b)$ ,  $(\Phi-c)$  and  $(\Phi-d)$  for  $\Phi_a$  follow automatically. Since  $\Phi \in \mathcal{P}_\tau$  is translation covariant and  $l(a, I)$  is translation invariant under translation of  $I$  by (8.10),  $\Phi_a$  satisfies the translation covariance  $(\Phi-f)$ .  $\Phi_a$  is of a finite range because there is no translates of  $\mathcal{C}_a$  containing  $I$  if  $\text{diam}(I) > \sqrt{\nu}(a-1)$  and hence  $l(a, I) = 0$  for such  $I$  and  $a \in \mathbb{N}$ . Hence  $(\Phi-e)$  is automatically satisfied. Therefore we conclude that  $\Phi_a \in \mathcal{P}_\tau^f$ . We compute

$$\begin{aligned} E_a(H_\Phi(\{0\})) &= \sum_{J \ni 0} \frac{1}{|\mathcal{C}_a|} \sum_{i \in \mathcal{C}_a} E_{\mathcal{C}_a-i}(\Phi(J)) \\ &= \sum_{J \ni 0} \frac{l(a, J)}{|\mathcal{C}_a|} \Phi(J) = H_{\Phi_a}(\{0\}), \end{aligned}$$

where we have used  $E_{\mathcal{C}_a-i}(\Phi(J)) = \Phi(J)$  for  $\mathcal{C}_a - i \supset J$  and  $E_{\mathcal{C}_a-i}(\Phi(J)) = 0$  for  $\mathcal{C}_a - i \not\supset J$  due to  $(\Phi-d)$ . (Note that if a translate  $\mathcal{C}_a - i$  contains  $J$ , then  $i \in \mathcal{C}_a$  due to  $0 \in J$  and hence the number of  $i \in \mathcal{C}_a$ , for which  $\mathcal{C}_a - i \supset J$ , is  $l(a, J)$ .)

By Lemma 8.11, we obtain

$$\begin{aligned} \lim_{a \rightarrow \infty} \|\Phi - \Phi_a\| &= \lim_{a \rightarrow \infty} \|H_\Phi(\{0\}) - H_{\Phi_a}(\{0\})\| \\ &= \lim_{a \rightarrow \infty} \|H_\Phi(\{0\}) - E_a(H_\Phi(\{0\}))\| = 0. \end{aligned}$$

This completes the proof.  $\square$

**Corollary 8.13.**  $\mathcal{P}_\tau$  is a separable Banach space.

*Proof.* For each  $n \in \mathbb{N}$ , the set of all  $\Phi \in \mathcal{P}_\tau^f$  with its range not exceeding  $n$  is a finite dimensional subspace of  $\mathcal{P}_\tau$ , because such  $\Phi$  is determined by  $\Phi(I)$  for a finite number of  $I$  containing the origin and satisfying  $\text{diam}(I) \leq n$ , and so has a dense countable subset. Taking union over  $n \in \mathbb{N}$ , we have a countable dense subset of  $\mathcal{P}_\tau^f$ . By Theorem 8.12, the same countable subset is dense in  $\mathcal{P}_\tau$ . We have now shown that  $\mathcal{P}_\tau$  is separable.  $\square$

## 9 Thermodynamic Limit

The van Hove limits of the densities (volume average) of extensive quantities are usually called thermodynamic limits. We now provide their existence theorems. The same proof as the case of spin lattice systems (see, e.g., [17], [23] and [40]) is applicable to the present Fermion lattice case. We, however, present slightly simplified proof by using methods different from those of the known proof. First we derive a surface energy estimate which we will find useful and crucial in the argument of the present section.

### 9.1 Surface Energy Estimate

**Lemma 9.1.** *For  $\Phi \in \mathcal{P}_\tau$ ,*

$$\text{v.H.} \lim_{I \rightarrow \infty} \frac{\|W(I)\|}{|I|} = 0 \quad (9.1)$$

*Proof.* Let  $\{I_\alpha\}$  be an arbitrary van Hove net of  $\mathbb{Z}^\nu$ . For  $n \in \mathbb{Z}^\nu$  and a finite subset  $I$  of  $\mathbb{Z}^\nu$ , let

$$\begin{aligned} W_n(I) &\equiv \lim_{J \nearrow \mathbb{Z}^\nu} \sum_K \left\{ \Phi(K); K \ni n, K \cap I^c \neq \emptyset, K \subset J \right\} \\ &= \lim_{J \nearrow \mathbb{Z}^\nu} \left( H_J(\{n\}) - E_I \left\{ H_J(\{n\}) \right\} \right) \\ &= H(\{n\}) - E_I \{ H(\{n\}) \}. \end{aligned}$$

Let  $B_r^{\mathbb{Z}^\nu}(n)$  be the intersection of  $B_r(n)$  (the ball with its center  $n$  and radius  $r$ ) and  $\mathbb{Z}^\nu$ . If  $n \in I$  and  $n \notin \text{surf}_r(I)$ , then  $B_r^{\mathbb{Z}^\nu}(n) \subset I$  and hence

$$E_I \left( H_{B_r^{\mathbb{Z}^\nu}(n)}(\{n\}) \right) = H_{B_r^{\mathbb{Z}^\nu}(n)}(\{n\}).$$

Therefore,

$$W_n(I) = H(\{n\}) - H_{B_r^{\mathbb{Z}^\nu}(n)}(\{n\}) - E_I \left\{ H(\{n\}) - H_{B_r^{\mathbb{Z}^\nu}(n)}(\{n\}) \right\}.$$

From this, we obtain

$$\|W_n(I)\| \leq 2 \|H(\{n\}) - H_{B_r^{\mathbb{Z}^\nu}(n)}(\{n\})\|.$$

By (5.23), for given  $\varepsilon > 0$ , we can take sufficiently large  $r > 0$  (hence sufficiently large  $B_r(0)$ ) satisfying

$$\|H(\{0\}) - H_{B_r^{\mathbb{Z}^\nu}(0)}(\{0\})\| < \frac{\varepsilon}{4}.$$

By the translation covariance assumption on  $\Phi$ , we have

$$\begin{aligned} \|H(\{n\}) - H_{B_r(n)}(\{n\})\| &= \left\| \tau_n \left\{ H(\{0\}) - H_{B_r^{\mathbb{Z}^\nu}(0)}(\{0\}) \right\} \right\| \\ &= \|H(\{0\}) - H_{B_r^{\mathbb{Z}^\nu}(0)}(\{0\})\| \\ &< \frac{\varepsilon}{4}. \end{aligned}$$

Hence

$$\|W_n(\mathbf{I})\| \leq \frac{\varepsilon}{2}, \quad (9.2)$$

if  $n \in \mathbf{I}$  and  $n \notin \text{surf}_r(\mathbf{I})$ .

For  $\mathbf{I} = \{n_1, \dots, n_{|\mathbf{I}|}\}$ , we have

$$W(\mathbf{I}) = \sum_{i=1}^{|\mathbf{I}|} E_{\{n_1, \dots, n_{i-1}\}^c} W_{n_i}(\mathbf{I}) \quad (9.3)$$

and hence

$$\|W(\mathbf{I})\| \leq \sum_{i=1}^{|\mathbf{I}|} \|W_{n_i}(\mathbf{I})\|. \quad (9.4)$$

For  $n = n_i \notin \text{surf}_r(\mathbf{I})$ , we use the estimate (9.2) for  $\|W_n(\mathbf{I})\|$ . For  $n = n_i \in \text{surf}_r(\mathbf{I})$ , we use

$$\|W_n(\mathbf{I})\| = \|H(\{n\}) - E_{\mathbf{I}}(H(\{n\}))\| \leq 2\|H(\{n\})\| = 2\|\Phi\|.$$

Then

$$\|W(\mathbf{I})\| \leq \frac{\varepsilon}{2} \cdot |\mathbf{I}| + 2\|\Phi\| \cdot |\text{surf}_r(\mathbf{I})|. \quad (9.5)$$

Since  $\{\mathbf{I}_\alpha\}$  is a van Hove net, there exists  $\alpha_\varepsilon$  such that, for  $\alpha \geq \alpha_\varepsilon$ ,

$$\frac{|\text{surf}_r(\mathbf{I}_\alpha)|}{|\mathbf{I}_\alpha|} < \frac{\varepsilon}{4\|\Phi\|}.$$

For such  $\alpha$ , we obtain

$$\frac{\|W(\mathbf{I}_\alpha)\|}{|\mathbf{I}_\alpha|} < \varepsilon,$$

which completes the proof.  $\square$

**Lemma 9.2.** *Let  $\{\mathbf{I}_\alpha\}$  be a van Hove net of  $\mathbb{Z}^\nu$ . For each  $\mathbf{I}_\alpha$  and  $a \in \mathbb{N}$ , take a set of mutually disjoint  $\mathbf{n}_a^-(\mathbf{I}_\alpha)$  translates  $D_i^{(a,\alpha)}$  of  $\mathcal{C}_a$  which are all packed in  $\mathbf{I}_\alpha$ . For any  $\varepsilon > 0$ , take an  $a_0 \in \mathbb{N}$  such that  $\|W(\mathcal{C}_a)\| < |\mathcal{C}_a| \varepsilon/2$  for all  $a > a_0$ . For any such  $a$ , there exists an  $\alpha_0(a)$  such that, for  $\alpha > \alpha_0(a)$ ,*

$$\|H(\mathbf{I}_\alpha) - \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)})\| < \mathbf{n}_a^-(\mathbf{I}_\alpha) |\mathcal{C}_a| \varepsilon, \quad (9.6)$$

$$\|U(\mathbf{I}_\alpha) - \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)})\| < \mathbf{n}_a^-(\mathbf{I}_\alpha) |\mathcal{C}_a| \varepsilon, \quad (9.7)$$

and

$$1 \geq \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha) |\mathcal{C}_a|}{|\mathbf{I}_\alpha|} \geq 1 - \frac{\varepsilon}{\|\Phi\|}. \quad (9.8)$$

*Proof.* Before we start the proof, we note that the existence of  $a_0$  is guaranteed by Lemma 9.1. Let us set

$$D^{(a,\alpha)} \equiv \bigcup_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} D_i^{(a,\alpha)}, \quad D'^{(a,\alpha)} \equiv \mathbf{I}_\alpha \setminus D^{(a,\alpha)}.$$

Obviously

$$|D'^{(a,\alpha)}| \leq \left( \mathbf{n}_a^+(\mathbf{I}_\alpha) - \mathbf{n}_a^-(\mathbf{I}_\alpha) \right) |\mathcal{C}_a|,$$

and

$$\mathbf{n}_a^+(\mathbf{I}_\alpha) |\mathcal{C}_a| \geq |\mathbf{I}_\alpha| \geq \mathbf{n}_a^-(\mathbf{I}_\alpha) |\mathcal{C}_a|.$$

From this, we obtain

$$\begin{aligned} 1 &\geq \frac{|\mathbf{I}_\alpha|}{\mathbf{n}_a^+(\mathbf{I}_\alpha) |\mathcal{C}_a|} \geq \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)}{\mathbf{n}_a^+(\mathbf{I}_\alpha)}, \\ 1 &\geq \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha) |\mathcal{C}_a|}{|\mathbf{I}_\alpha|} \geq \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)}{\mathbf{n}_a^+(\mathbf{I}_\alpha)}. \end{aligned} \quad (9.9)$$

On the other hand,

$$\begin{aligned} &H(\mathbf{I}_\alpha) - \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)}) \\ &= \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} E_{\{D_1^{(a,\alpha)} \cup \dots \cup D_{i-1}^{(a,\alpha)}\}^c} \left( W(D_i^{(a,\alpha)}) \right) + E_{\{D^{(a,\alpha)}\}^c} \left( H(D'^{(a,\alpha)}) \right). \end{aligned}$$

Therefore,

$$\begin{aligned} \|H(\mathbf{I}_\alpha) - \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)})\| &\leq \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} \|W(D_i^{(a,\alpha)})\| + \|H(D'^{(a,\alpha)})\| \\ &\leq \mathbf{n}_a^-(\mathbf{I}_\alpha) |\mathcal{C}_a| \cdot \frac{\varepsilon}{2} + \|\Phi\| |D'^{(a,\alpha)}|, \end{aligned} \quad (9.10)$$

where in the second inequality the assumption  $\|W(\mathcal{C}_a)\| < |\mathcal{C}_a| \varepsilon/2$  together with the translation covariance of  $\Phi$  are used for  $\|W(D_i^{(a,\alpha)})\|$ , and Lemma 8.6 is used for  $\|H(D'^{(a,\alpha)})\|$ . Due to condition (I) for the van Hove limit, there exists  $\alpha_0(a)$  for given  $\varepsilon_1 > 0$  such that, for  $\alpha \geq \alpha_0(a)$ ,

$$0 \leq 1 - \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)}{\mathbf{n}_a^+(\mathbf{I}_\alpha)} < \varepsilon_1. \quad (9.11)$$

If  $\varepsilon_1 < 1$ , then

$$\begin{aligned} \mathbf{n}_a^+(\mathbf{I}_\alpha) &< \frac{1}{1 - \varepsilon_1} \mathbf{n}_a^-(\mathbf{I}_\alpha), \\ |D'^{(a,\alpha)}| &\leq \mathbf{n}_a^+(\mathbf{I}_\alpha) \varepsilon_1 |\mathcal{C}_a| < \frac{\varepsilon_1}{1 - \varepsilon_1} \mathbf{n}_a^-(\mathbf{I}_\alpha) |\mathcal{C}_a|. \end{aligned}$$

Now we choose  $\varepsilon_1$  which satisfies

$$\frac{2\varepsilon_1}{1-\varepsilon_1}\|\Phi\| < \varepsilon, \text{ and } (0 <) \varepsilon_1 < 1. \quad (9.12)$$

Then from (9.10) and (9.12), we have

$$\begin{aligned} \|H(\mathbf{I}_\alpha) - \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)})\| &\leq \mathbf{n}_a^-(\mathbf{I}_\alpha)|\mathcal{C}_a| \cdot \frac{\varepsilon}{2} + \|\Phi\| \frac{\varepsilon_1}{1-\varepsilon_1} \mathbf{n}_a^-(\mathbf{I}_\alpha)|\mathcal{C}_a| \\ &= \mathbf{n}_a^-(\mathbf{I}_\alpha)|\mathcal{C}_a| \left( \frac{\varepsilon}{2} + \frac{\varepsilon_1}{1-\varepsilon_1} \|\Phi\| \right) \\ &< \mathbf{n}_a^-(\mathbf{I}_\alpha)|\mathcal{C}_a|\varepsilon. \end{aligned}$$

We also have

$$\begin{aligned} \|U(\mathbf{I}_\alpha) - \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)})\| &= \|E_{\mathbf{I}_\alpha} \left( H(\mathbf{I}_\alpha) - \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)}) \right)\| \\ &< \mathbf{n}_a^-(\mathbf{I}_\alpha)|\mathcal{C}_a|\varepsilon. \end{aligned}$$

Due to (9.12),

$$\varepsilon_1 < \frac{\varepsilon}{\|\Phi\|}$$

By (9.9), (9.11) and this inequality, we obtain

$$1 \geq \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)|\mathcal{C}_a|}{|\mathbf{I}_\alpha|} \geq 1 - \frac{\varepsilon}{\|\Phi\|}.$$

□

## 9.2 Pressure

**Theorem 9.3.** Assume  $\Phi \in \mathcal{P}_\tau$ .

(1) The following limit exists:

$$\begin{aligned} p(\Phi) &\equiv \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{1}{|\mathbf{I}|} \log \tau(e^{-H(\mathbf{I})}) \\ &= \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{1}{|\mathbf{I}|} \log \tau(e^{-U(\mathbf{I})}). \end{aligned} \quad (9.13)$$

(2)  $p(\Phi)$  is a convex functional of  $\Phi$  satisfying the following continuity property:

$$|p(\Phi) - p(\Psi)| \leq \|\Phi - \Psi\|. \quad (9.14)$$

*Proof.* We first prove (1) in four steps.

Step 1. We need the following well-known matrix inequality:

$$|\log \tau(e^{-A}) - \log \tau(e^{-B})| \leq \|A - B\|, \quad (9.15)$$

for  $A, B \in \mathcal{A}_0$ . This follows from the following computation:

$$\begin{aligned} |\log \tau(e^{-A}) - \log \tau(e^{-B})| &= \left| \int_0^1 \frac{d}{d\lambda} \left\{ \log \tau(e^{-\lambda A - (1-\lambda)B}) \right\} d\lambda \right| \\ &= \left| \int_0^1 \frac{\tau(e^{-\lambda A - (1-\lambda)B} \cdot (B - A))}{\tau(e^{-\lambda A - (1-\lambda)B})} d\lambda \right| \leq \|A - B\|, \end{aligned}$$

where we have used the fact that  $\tau(e^c x)/\tau(e^c)$  for  $c = c^* \in \mathcal{A}_0$  is a state function of  $x \in \mathcal{A}$  and hence bounded by  $\|x\|$ . Setting  $B = 0$  and noting  $\log \tau(e^{-B}) = 0$  for  $B = 0$ , we have

$$|\log \tau(e^{-A})| \leq \|A\|. \quad (9.16)$$

Step 2. We use the notation in the preceding Lemma. Because  $U(D_i^{(a,\alpha)})$  with distinct  $i$ 's mutually commute due to the disjointness of  $D_i^{(a,\alpha)}$ , (5.21), (4.8) and  $(\Phi\text{-c})$ , we have

$$\begin{aligned} \log \tau \left( e^{\{-\sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)})\}} \right) &= \log \tau \left( \prod_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} e^{-U(D_i^{(a,\alpha)})} \right) \\ &= \log \prod_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} \tau(e^{-U(D_i^{(a,\alpha)})}) = \sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} \log \tau(e^{-U(D_i^{(a,\alpha)})}) \\ &= \mathbf{n}_a^-(\mathbf{I}_\alpha) \log \tau(e^{-U(\mathcal{C}_a)}), \end{aligned} \quad (9.17)$$

where the second equality is due to the product property (4.13) of the tracial state, and the last equality follows from the translation covariance  $(\Phi\text{-f})$ . By (9.16), (8.3) and (9.8), we have

$$\left| \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)}{|\mathbf{I}_\alpha|} \log \tau(e^{-U(\mathcal{C}_a)}) - \frac{1}{|\mathcal{C}_a|} \log \tau(e^{-U(\mathcal{C}_a)}) \right| < \varepsilon. \quad (9.18)$$

Step 3. By (9.17), (9.18), (9.15), (9.6) and (9.8),

$$\begin{aligned} &\left| \frac{1}{|\mathbf{I}_\alpha|} \log \tau(e^{-H(\mathbf{I}_\alpha)}) - \frac{1}{|\mathcal{C}_a|} \log \tau(e^{-U(\mathcal{C}_a)}) \right| \\ &= \left| \frac{1}{|\mathbf{I}_\alpha|} \log \tau(e^{-H(\mathbf{I}_\alpha)}) - \frac{1}{|\mathbf{I}_\alpha|} \log \tau \left( e^{\{-\sum_{i=1}^{\mathbf{n}_a^-(\mathbf{I}_\alpha)} U(D_i^{(a,\alpha)})\}} \right) + \left( \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)}{|\mathbf{I}_\alpha|} - \frac{1}{|\mathcal{C}_a|} \right) \log \tau(e^{-U(\mathcal{C}_a)}) \right| \\ &< 2\varepsilon \end{aligned} \quad (9.19)$$

for any  $\alpha > \alpha_0(a)$ . Hence for any  $\alpha, \beta > \alpha_0(a)$ , we have

$$\left| \frac{1}{|\mathbf{I}_\alpha|} \log \tau(e^{-H(\mathbf{I}_\alpha)}) - \frac{1}{|\mathbf{I}_\beta|} \log \tau(e^{-H(\mathbf{I}_\beta)}) \right| < 4\varepsilon.$$

Therefore,  $\frac{1}{|\mathbf{I}_\alpha|} \log \tau(e^{-H(\mathbf{I}_\alpha)})$  is a Cauchy net in  $\mathbb{R}$  and has the (van Hove) limit.

Step 4. Due to

$$\text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{\|H(\mathbf{I}) - U(\mathbf{I})\|}{|\mathbf{I}|} = \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{\|W(\mathbf{I})\|}{|\mathbf{I}|} = 0$$



and

$$|\log \tau(e^{-H(I)}) - \log \tau(e^{-U(I)})| \leq \|H(I) - U(I)\|,$$

the convergence of  $\frac{1}{|I_\alpha|} \log \tau(e^{-H(I_\alpha)})$  implies that of  $\frac{1}{|I_\alpha|} \log \tau(e^{-U(I_\alpha)})$  to the same value.

Now we prove (2). Since  $H_\Phi(I)$  is linear in  $\Phi$ , we have the convexity of  $\log \tau(e^{-H_\Phi(I)})$  in  $\Phi$  due to the well-known convexity of the function:

$$\lambda \mapsto \log \tau(e^{(A+\lambda B)}) \quad \text{for } A = A^* \text{ and } B = B^*.$$

Hence the convexity of  $p(\Phi)$  follows. By (9.15), the linearity of  $H_\Phi(I)$  in  $\Phi$  and (8.3), we obtain

$$\begin{aligned} & \left| \frac{1}{|I|} \log \tau(e^{-H_\Phi(I)}) - \frac{1}{|I|} \log \tau(e^{-H_\Psi(I)}) \right| \\ & \leq \frac{1}{|I|} \|H_\Phi(I) - H_\Psi(I)\| = \frac{1}{|I|} \|H_{\Phi-\Psi}(I)\| \\ & \leq \|\Phi - \Psi\| \end{aligned}$$

for any finite  $I$ . Hence (9.14) follows.  $\square$

The pressure functional  $P(\Phi)$  of  $\Phi \in \mathcal{P}_\tau$  is conventionally defined by using the matrix trace in contrast to  $p(\Phi)$  in the preceding theorem defined in terms of the tracial state:

$$P(\Phi) \equiv \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \log \mathbf{Tr}_I(e^{-H(I)}) \left( = \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \log \mathbf{Tr}_I(e^{-U(I)}) \right), \quad (9.20)$$

where  $\mathbf{Tr}_I$  denotes the matrix trace on  $\mathcal{A}(I)$  and hence  $\mathbf{Tr}_I = 2^{|I|} \tau$ . Therefore, for any  $\Phi \in \mathcal{P}_\tau$ ,

$$P(\Phi) = p(\Phi) + \log 2. \quad (9.21)$$

Due to the preceding theorem, we have obviously

**Corollary 9.4.** *Assume  $\Phi \in \mathcal{P}_\tau$ .*

(1) *The following limit exists:*

$$\begin{aligned} P(\Phi) & \equiv \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \log \mathbf{Tr}_I(e^{-H(I)}) \\ & = \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \log \mathbf{Tr}_I(e^{-U(I)}). \end{aligned} \quad (9.22)$$

(2)  *$P(\Phi)$  is a convex functional of  $\Phi$  satisfying the following continuity property:*

$$|P(\Phi) - P(\Psi)| \leq \|\Phi - \Psi\|. \quad (9.23)$$

*Remark.* We have

$$p(0) = 0, \quad |p(\Phi)| \leq \|\Phi\| \quad (9.24)$$

which do not hold for  $P(\Phi)$ .

### 9.3 Mean Energy

**Theorem 9.5.** *For  $\Phi \in \mathcal{P}_\tau$  and a translation invariant state  $\omega$  of  $\mathcal{A}$ , the following limit exists:*

$$\begin{aligned} e_\Phi(\omega) &\equiv \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \omega(H(I)) \\ &= \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \omega(U(I)). \end{aligned} \quad (9.25)$$

The mean energy  $e_\Phi(\omega)$  so obtained is linear in  $\Phi$ , affine in  $\omega$ , bounded by  $\|\Phi\|$ , and weak\* continuous in  $\omega$ :

$$e_{c\Phi+d\Psi}(\omega) = ce_\Phi(\omega) + de_\Psi(\omega) \quad (c, d \in \mathbb{R}), \quad (9.26)$$

$$e_\Phi(\lambda\omega_1 + (1-\lambda)\omega_2) = \lambda e_\Phi(\omega_1) + (1-\lambda)e_\Phi(\omega_2) \quad (0 \leq \lambda \leq 1), \quad (9.27)$$

$$|e_\Phi(\omega)| \leq \|\Phi\|, \quad (9.28)$$

$$\lim_\gamma e_\Phi(\omega_\gamma) = e_\Phi(\omega), \quad (9.29)$$

where  $\Phi$  and  $\Psi$  are in  $\mathcal{P}_\tau$ ,  $\omega, \omega_1, \omega_2$  and  $\omega_\gamma$  are in  $\mathcal{A}_{+,1}^{*\tau}$ , and  $\{\omega_\gamma\}$  is a net converging to  $\omega$  in the weak\* topology.

*Proof.* By the argument leading to (9.19) in Theorem 9.3, there exists  $a \in \mathbb{N}$  and  $\alpha_0(a)$  for any given  $\varepsilon > 0$  such that for all  $\alpha > \alpha_0(a)$

$$\left| \frac{1}{|I_\alpha|} \omega(H(I_\alpha)) - \frac{1}{|\mathcal{C}_a|} \omega(U(\mathcal{C}_a)) \right| < 2\varepsilon, \quad (9.30)$$

where we can take the same  $a \in \mathbb{N}$  and  $\alpha_0(a)$  uniformly in  $\omega \in \mathcal{A}_{+,1}^*$ . This estimate implies that  $\left\{ \frac{1}{|I_\alpha|} \omega(H(I_\alpha)) \right\}_\alpha$  is a Cauchy net in  $\mathbb{R}$  and hence converges.

Since  $\omega(H(I))$  is linear in  $\Phi$  and affine in  $\omega$ , so is  $e_\Phi(\omega)$ . Due to (8.3), we obtain  $|e_\Phi(\omega)| \leq \|\Phi\|$ .

Finally we show the continuity in  $\omega$ . Let  $\{\omega_\gamma\}_\gamma$  be a net of states converging to  $\omega$  in the weak\* topology. For any  $\varepsilon > 0$ , we fix  $a \in \mathbb{N}$  satisfying (9.30) for all  $\alpha > \alpha_0(a)$  and for all states  $\omega$ . From the weak\* convergence of  $\{\omega_\gamma\}_\gamma$  to  $\omega$ , there exists  $\gamma_\varepsilon$  such that for all  $\gamma \geq \gamma_\varepsilon$

$$\frac{1}{|\mathcal{C}_a|} \left| \omega(U(\mathcal{C}_a)) - \omega_\gamma(U(\mathcal{C}_a)) \right| < \varepsilon.$$

Thus we have

$$\left| \frac{1}{|I_\alpha|} \omega(H(I_\alpha)) - \frac{1}{|I_\alpha|} \omega_\gamma(H(I_\alpha)) \right| < 5\varepsilon,$$

for all  $\alpha > \alpha_0(a)$ . By taking the van Hove limit, we obtain

$$|e_\Phi(\omega) - e_\Phi(\omega_\gamma)| < 5\varepsilon$$

for all  $\gamma \geq \gamma_\varepsilon$ . Hence  $e_\Phi(\omega)$  is continuous in  $\omega$  relative to the weak\* topology.  $\square$

## 10 Entropy for Fermion Systems

### 10.1 SSA for Fermion Systems

We first show the SSA property of entropy for the Fermion case, which is a consequence of the results on the conditional expectations in § 3 and § 4.

**Theorem 10.1.** *For finite subsets  $I$  and  $J$  of  $\mathbb{Z}^\nu$ , the strong subadditivity (SSA) of  $\widehat{S}$  holds for any state  $\psi$  of  $\mathcal{A}$ :*

$$\widehat{S}(\psi_{I \cup J}) - \widehat{S}(\psi_I) - \widehat{S}(\psi_J) + \widehat{S}(\psi_{I \cap J}) \leq 0, \quad (10.1)$$

where  $\psi_K$  denotes the restriction of  $\psi$  to  $\mathcal{A}(K)$ .  $\widehat{S}$  in this inequality can be replaced by  $S$ :

$$S(\psi_{I \cup J}) - S(\psi_I) - S(\psi_J) + S(\psi_{I \cap J}) \leq 0. \quad (10.2)$$

*Proof.* The SSA of  $\widehat{S}$  follows from Theorem 3.7 and Theorem 4.13. By (3.1) and

$$\log 2^{|I \cup J|} - \log 2^{|I|} - \log 2^{|J|} + \log 2^{|I \cap J|} = 0,$$

the SSA of  $\widehat{S}$  implies that of  $S$ . □

*Remark 1.* The strong subadditivity can be rewritten as

$$S(\psi_{123}) - S(\psi_{13}) - S(\psi_{23}) + S(\psi_3) \leq 0, \quad (10.3)$$

for any disjoint subsets  $I_1, I_2$  and  $I_3$  of  $\mathbb{Z}^\nu$ , where  $\psi_{123}$  denotes the restriction of  $\psi$  to  $\mathcal{A}(I_1 \cup I_2 \cup I_3)$ , and so on.

*Remark 2.* The SSA for Fermion systems above does not seem to follow from those for the tensor product systems ([27], [28]) in any obvious way.

*Remark 3.* Note that the SSA for Fermion systems holds whether the state  $\psi$  is  $\Theta$ -even or not. For two disjoint finite regions  $I$  and  $J$ , the so-called triangle inequality of entropy

$$|S(\psi_I) - S(\psi_J)| \leq S(\psi_{I \cup J})$$

is known to hold for quantum spin lattice systems [1]. However, it can fail for Fermion lattice systems when  $\psi$  breaks  $\Theta$ -evenness (see a concrete example in [33]).

The following is a special case of Theorem 10.1 when  $I \cap J = \emptyset$ .

**Corollary 10.2.** *For disjoint finite subsets  $I$  and  $J$ , the following subadditivity holds.*

$$\widehat{S}(\psi_{I \cup J}) \leq \widehat{S}(\psi_I) + \widehat{S}(\psi_J), \quad (10.4)$$

$$S(\psi_{I \cup J}) \leq S(\psi_I) + S(\psi_J). \quad (10.5)$$

## 10.2 Mean Entropy

We now show the existence of mean entropy (von Neumann entropy density) for translation invariant states of  $\mathcal{A}$ .

For  $s = (s_1, \dots, s_\nu) \in \mathbb{N}^\nu$ , we define  $R_s$  as the following box region with edges of length  $s_i - 1$  containing  $s_i$  points of  $\mathbb{Z}^\nu$  and with the volume  $|R_s| = \prod_{i=1}^\nu s_i$ .

$$R_s \equiv \{x \in \mathbb{Z}^\nu ; 0 \leq x_i \leq s_i - 1, i = 1, \dots, \nu\}. \quad (10.6)$$

**Theorem 10.3.** *Let  $\omega$  be a translation invariant state. The van Hove limit*

$$s(\omega) \equiv \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} S(\omega_I) \quad (10.7)$$

*exists and is given as the following infimum*

$$s(\omega) = \inf_{s \in \mathbb{N}^\nu} \frac{1}{|R_s|} S(\omega_{R_s}). \quad (10.8)$$

*The mean entropy functional*

$$\omega \mapsto s(\omega) \in [0, \log 2] \quad (10.9)$$

*defined on the set  $\mathcal{A}_{+,1}^{*\tau}$  of translation invariant states is affine and upper semi-continuous with respect to the weak\* topology.*

*Proof.* The SSA property of von Neumann entropy proved in Theorem 10.1 is sufficient for the same proof of this Theorem as in the case of quantum spin lattice systems. (See e.g. Proposition 6.2.38 of [17].)  $\square$

The following results about Lipschitz continuity of bounded affine functions on a state space and, in particular, of entropy density are known.

**Proposition 10.4.** *A bounded affine function  $f$  on  $\mathcal{A}_{+,1}^{*\tau}$  satisfies*

$$|f(\omega_1) - f(\omega_2)| \leq (M/2) \|\omega_1 - \omega_2\| \quad (10.10)$$

*for any  $\omega_1, \omega_2 \in \mathcal{A}_{+,1}^{*\tau}$ , where*

$$M \equiv \sup\{|f(\omega_1) - f(\omega_2)| ; \omega_1, \omega_2 \in \mathcal{A}_{+,1}^{*\tau}\}.$$

**Corollary 10.5.** *The mean entropy  $s(\omega)$  satisfies*

$$|s(\omega_1) - s(\omega_2)| \leq \frac{1}{2} (\log 2) \|\omega_1 - \omega_2\| \quad (10.11)$$

*for any  $\omega_1, \omega_2 \in \mathcal{A}_{+,1}^{*\tau}$ .*

Proposition 10.4 is the first equation on page 108 of [23] and Corollary 10.5 is Corollary IV.4.3 on the same page of [23]. The inequality (10.11) without  $\frac{1}{2}$  factor is obtained in [20]. The coefficient  $\frac{1}{2} \log 2$  is best possible, the equality being reached by  $\omega_1 = \tau$  and any pure translation invariant state  $\omega_2$  with vanishing mean entropy  $s(\omega_2) = 0$ , in which case  $\|\omega_1 - \omega_2\| = 2$  because  $\pi_\tau$  (type II) and  $\pi_{\omega_2}$  (type I) are disjoint. An example of such an  $\omega_2$  is given by Theorem 11.2 as a ‘product state extension’ of  $\Theta$ -even pure states  $\varphi_i$  of  $\mathcal{A}(\{i\})$  ( $i \in \mathbb{Z}^\nu$ ) satisfying the covariance condition  $\tau_k^* \varphi_i = \varphi_{i+k}$  for all  $k \in \mathbb{Z}^\nu$ .

We define mean entropy  $\hat{s}(\omega)$  for  $\omega \in \mathcal{A}_{+,1}^{*\tau}$  by using trace  $\tau$  instead of matrix trace  $\mathbf{Tr}_I$  for each finite  $I$ :

$$\hat{s}(\omega) \equiv \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \hat{S}(\omega_I). \quad (10.12)$$

It is obviously related to  $s(\omega)$  by

$$s(\omega) = \hat{s}(\omega) + \log 2, \quad (10.13)$$

for any  $\omega \in \mathcal{A}_{+,1}^{*\tau}$ .

### 10.3 Entropy Inequalities for Translation Invariant States

In addition to Theorem 10.3, the SSA property of von Neumann entropy plays an essential role in the derivation of some basic entropy inequalities for the present Fermion lattice systems in the same way as for quantum spin lattice systems. The following two consequences are about monotone properties of entropy as a function on the set of box regions of the lattice; the first one is a monotone decreasing property of the finite-volume entropy density and the second one is a monotone increasing property of the entropy.

**Theorem 10.6.** *Let  $\omega$  be a translation invariant state on  $\mathcal{A}$  and let  $R_s$  and  $R_{s'}$  be finite boxes of  $\mathbb{Z}^\nu$  such that  $R_s \subset R_{s'}$ . Then*

$$\frac{1}{|R_s|} S(\omega_{R_s}) \geq \frac{1}{|R_{s'}|} S(\omega_{R_{s'}}), \quad (10.14)$$

$$S(\omega_{R_s}) \leq S(\omega_{R_{s'}}). \quad (10.15)$$

This theorem follows from [24], where (10.14) and (10.15) are derived from the following properties without any other input.

- Positivity and finiteness of the entropy of every local region,
- Strong subadditivity.
- Shift invariance.

In [16], sufficient conditions are given for a sequence of regions of more general shape than boxes which guarantee a monotone decreasing property of the form (10.14) for any translation invariant state  $\omega$ . This result also applies to our Fermion lattice systems.

## 11 Variational Principle

We first prove the existence of a (unique) product state extension of given states in any (finite or infinite) number of mutually disjoint regions under the condition that all given states except for at most one are  $\Theta$ -even.

This result is a crucial tool to overcome possible difficulties which originate in the non-commutativity of Fermion systems in connection with the proof of variational equality in this section and in the equivalence proof of the variational principle with the KMS condition in the next section.

### 11.1 Extension of Even States

For each  $I$ ,  $\mathcal{A}(I)$  is invariant under  $\Theta$  and hence the restriction of  $\Theta$  to  $\mathcal{A}(I)$  is an automorphism of  $\mathcal{A}(I)$  and will be denoted by the same symbol  $\Theta$ . We need the following lemma.

**Lemma 11.1.** *Let  $I$  be a finite subset of  $\mathbb{Z}^\nu$ . Let  $\varphi$  be a state of  $\mathcal{A}(I)$  and  $\varrho \in \mathcal{A}(I)$  be its adjusted density matrix :*

$$\varphi(A) = \tau(\varrho A) = \tau(A\varrho), \quad (A \in \mathcal{A}(I)).$$

*Then  $\varphi$  is an even state if and only if  $\varrho$  is  $\Theta$ -even.*

*Proof.* Since the tracial state  $\tau$  is invariant under any automorphism, we obtain

$$\begin{aligned} \varphi(A) &= \varphi(\Theta(A)) = \tau(\varrho \Theta(A)) = \tau(\Theta\{\varrho \Theta(A)\}) \\ &= \tau(\Theta(\varrho)A) \end{aligned}$$

if  $\varphi$  is even. By the uniqueness of the density matrix, we have  $\Theta(\varrho) = \varrho$ .

By the same computation,  $\varphi(\Theta(A)) = \varphi(A)$  for every  $A \in \mathcal{A}(I)$  if  $\Theta(\varrho) = \varrho$ .

□

**Theorem 11.2.** *Let  $\{I_i\}$  be a (finite or infinite) family of mutually disjoint subsets of  $\mathbb{Z}^\nu$  and  $\varphi_i$  be a state of  $\mathcal{A}(I_i)$  for each  $i$ . Let  $I = \bigcup_i I_i$ . Then there exists a state  $\varphi$  of  $\mathcal{A}(I)$  satisfying*

$$\varphi(A_{i_1} \cdots A_{i_n}) = \prod_{j=1}^n \varphi_{i_j}(A_{i_j}) \quad (11.1)$$

*for any set  $(i_1, \dots, i_n)$  of distinct indices and for any  $A_{i_j} \in \mathcal{A}(I_{i_j})$  if all states  $\varphi_i$  except for at most one are  $\Theta$ -even. When such  $\varphi$  exists, it is unique.*

*Proof.* (Case 1) A finite family of finite subsets  $\{I_i\}$ ,  $i = 1, \dots, n$ .

For each  $i$ , let  $\varrho_i$  be the density matrix of  $\varphi_i$ :

$$\begin{aligned} \varphi_i(A) &= \tau(\varrho_i A) = \tau(A\varrho_i), \quad (A \in \mathcal{A}(I_i)), \\ \varrho_i &\in \mathcal{A}(I_i), \quad \varrho_i \geq 0, \quad \varrho_i(\mathbf{1}) = 1. \end{aligned}$$

If  $\varphi_i$  is  $\Theta$ -even, then  $\varrho_i$  is  $\Theta$ -even, namely,

$$\varrho_i \in \mathcal{A}(I_i)_+.$$

If all states  $\varphi_i$  except for one is even, all  $\varrho_i$  except for one belong to  $\mathcal{A}(I_i)_+$ . Thus each  $\varrho_i$  commutes with any  $\varrho_j$ . The product

$$\varrho = \varrho_n \cdots \varrho_1 \quad (11.2)$$

is a product of mutually commuting non-negative hermitian operators and hence it is positive. Define

$$\varphi(A) \equiv \tau(\varrho A), \quad A \in \mathcal{A}(I). \quad (11.3)$$

By the product property of  $\tau$  (4.13), we have

$$\begin{aligned}\varphi(A_1 \cdots A_n) &= \tau(\varrho A_1 \cdots A_n) = \tau(\varrho_{n-1} \cdots \varrho_1 A_1 \cdots A_{n-1} A_n \varrho_n) \\ &= \tau(\varrho_{n-1} \cdots \varrho_1 A_1 \cdots A_{n-1}) \tau(A_n \varrho_n) \\ &= \tau(\varrho_{n-1} \cdots \varrho_1 A_1 \cdots A_{n-1}) \varphi_n(A_n).\end{aligned}$$

Using this recursively, we obtain

$$\varphi(A_1 \cdots A_n) = \prod_{i=1}^n \varphi_i(A_i).$$

This also shows  $\varphi(\mathbf{1}) = 1$ . Hence the existence is proved for Case 1.

Since the monomials of the form (4.2) with all indices in  $\mathbf{I}$  are total in  $\mathcal{A}(\mathbf{I})$ , the uniqueness of a state  $\varphi$  of  $\mathcal{A}(\mathbf{I})$  satisfying the product property (11.1) follows.

(Case 2) A general family  $\{\mathbf{I}_i\}$ .

Let  $\{\mathbf{L}_k\}$  be an increasing sequence of finite subsets of  $\mathbb{Z}^\nu$  such that their union is  $\mathbb{Z}^\nu$ . Set  $\mathbf{I}_i^k \equiv \mathbf{I}_i \cap \mathbf{L}_k$  and  $\mathbf{I}^k \equiv \mathbf{I} \cap \mathbf{L}_k$  for each  $k$ . For each  $k$ , only a finite number (which will be denoted by  $n(k)$ ) of  $\mathbf{I}_i^k$  are non-empty and all of them are finite subsets of  $\mathbb{Z}^\nu$ . Note that the restriction of an even state  $\varphi_i$  to  $\mathcal{A}(\mathbf{I}_i^k)$  is even. Hence we can apply the result for Case 1 to  $\{\mathbf{I}_i^k\}$ . We obtain a unique product state  $\varphi^k$  of  $\mathcal{A}(\mathbf{I}^k)$  satisfying

$$\varphi^k(A_{i_1} \cdots A_{i_{n(k)}}) = \prod_{j=1}^{n(k)} \varphi_{i_j}^k(A_{i_j}), \quad A_{i_j} \in \mathcal{A}(\mathbf{I}_{i_j}^k). \quad (11.4)$$

By the uniqueness already proved, the restriction of  $\varphi^k$  to  $\mathcal{A}(\mathbf{I}^l)$  for  $l < k$  coincides with  $\varphi^l$ . There exists a state  $\varphi_\circ$  of the  $*$ -algebra  $\cup_k \mathcal{A}(\mathbf{I}^k)$  defined by

$$\varphi_\circ(A) = \varphi^k(A)$$

for  $A \in \mathcal{A}(\mathbf{I}^k)$ . Since  $\cup_k \mathbf{I}^k = \mathbf{I}$ ,  $\cup_k \mathcal{A}(\mathbf{I}^k)$  is dense in  $\mathcal{A}(\mathbf{I})$ . Then there exists a unique continuous extension  $\varphi$  of  $\varphi_\circ$  to  $\mathcal{A}(\mathbf{I})$  and  $\varphi$  is a state of  $\mathcal{A}(\mathbf{I})$ .

Take an arbitrary index  $n$ . Let

$$A = A_1 \cdots A_n, \quad A_i \in \mathcal{A}(\mathbf{I}_i).$$

Set  $A_i^k \equiv E_{\mathbf{L}_k}(A_i) \in \mathcal{A}(\mathbf{I}_i^k)$ . Since  $\mathbf{L}_k \nearrow \mathbb{Z}^\nu$ ,

$$\begin{aligned}A_i &= \lim_k A_i^k, \\ A &= \lim_k (A_1^k \cdots A_n^k).\end{aligned}$$

Hence

$$\begin{aligned}\varphi(A) &= \lim_k \varphi(A_1^k \cdots A_n^k) \\ &= \lim_k \varphi^k(A_1^k \cdots A_n^k) = \lim_k \prod_{i=1}^n \varphi_i(A_i^k) \\ &= \prod_{i=1}^n \varphi_i(A_i).\end{aligned}$$

Thus  $\varphi$  satisfies the product property (11.1).

The uniqueness of  $\varphi$  is proved in the same way as Case 1.  $\square$

*Remark 1.* This result is given in Theorem 5.4. of Power's Thesis [36].

*Remark 2.* The unique product state extension  $\varphi$  is even if and only if all  $\varphi_i$  are even.

*Remark 3.* The condition that all  $\varphi_i$  except for at most one are  $\Theta$ -even can be shown to be necessary for the existence of the product state extension  $\varphi$  satisfying (11.1) [14].

**Lemma 11.3.** *Let  $\{I_i\}$  be a finite family of mutually disjoint finite subsets of  $\mathbb{Z}^\nu$ . Let  $\varphi_i$  be a state of  $\mathcal{A}(I_i)$  for each  $i$  and all  $\varphi_i$  be  $\Theta$ -even with at most one exception. Let  $\varphi$  be their product state extension given by Theorem 11.2. Then*

$$S(\varphi) = \sum_i S(\varphi_i), \quad \widehat{S}(\varphi) = \sum_i \widehat{S}(\varphi_i). \quad (11.5)$$

*Proof.* This follows from the computation using the density matrix (11.2).

$$\widehat{S}(\varphi) = -\varphi(\log \varrho) = -\sum_i \varphi(\log \varrho_i) = -\sum_i \varphi_i(\log \varrho_i) = \sum_i \widehat{S}(\varrho_i). \quad (11.6)$$

Here the mutual commutativity of  $\varrho_i$  is used. Due to  $|I| = \sum_i |I_i|$ , we can replace  $\widehat{S}$  by  $S$ .  $\square$

## 11.2 Variational Inequality

We have already quoted the positivity of relative entropy:

$$S(\psi, \varphi) = \tau(\hat{\rho}_\varphi \log \hat{\rho}_\varphi - \hat{\rho}_\varphi \log \hat{\rho}_\psi) \geq 0, \quad (11.7)$$

where the equality holds if and only if  $\varphi = \psi$ .

Recall our notation (7.14) for the local Gibbs state  $\varphi_I^c$  of  $\mathcal{A}(I)$  with respect to  $(\Phi, \beta)$ . Let  $\omega$  be a state of  $\mathcal{A}$ . Substituting  $\psi = \varphi_I^c$  and  $\varphi = \omega_I$  into (11.7), we obtain

$$S(\varphi_I^c, \omega_I) = -\widehat{S}(\omega_I) + \beta\omega(U(I)) + \log \tau(e^{-\beta U(I)}) \geq 0. \quad (11.8)$$

Now we assume that  $\omega$  is translation invariant. By dividing the above inequality by  $|I|$  and then taking the van Hove limit  $I \rightarrow \infty$ , we obtain the following variational inequality

$$p(\beta\Phi) \geq \hat{s}(\omega) - \beta e_\Phi(\omega), \quad (11.9)$$

where  $\hat{s}(\omega)$  is given by (10.12). Equivalently, we have

$$P(\beta\Phi) \geq s(\omega) - \beta e_\Phi(\omega). \quad (11.10)$$



### 11.3 Variational Equality

The variational inequality in the preceding subsection is now strengthened to the following variational equality.

**Theorem 11.4.** *Let  $\Phi \in \mathcal{P}_\tau$ . Then*

$$P(\beta\Phi) = \sup_{\omega \in \mathcal{A}_{+,1}^{*\tau}} \left\{ s(\omega) - \beta e_\Phi(\omega) \right\}, \quad (11.11)$$

where  $P(\beta\Phi)$ ,  $s(\omega)$  and  $e_\Phi$  denote the pressure, mean entropy and mean energy, respectively, and  $\mathcal{A}_{+,1}^{*\tau}$  denotes the set of all translation invariant states of  $\mathcal{A}$ .

*Proof.* The proof below will be carried out in the same way as for classical or quantum lattice systems ([37] or e.g., Theorem III.4.5 in [40]), with a help of the product state extension provided by Theorem 11.2.

By the variational inequality (11.10), we only have to find a sequence  $\{\rho_n\}$  of translation invariant states of  $\mathcal{A}$  satisfying

$$\{s(\rho_n) - \beta e_\Phi(\rho_n)\} \rightarrow P(\beta\Phi) \quad (n \rightarrow \infty). \quad (11.12)$$

For this purpose, we interrupt the proof and show the following lemma about mean entropy and mean energy of periodic states. It corresponds to Theorem 10.3 and Theorem 9.5 for translation invariant states.

**Lemma 11.5.** *Let  $a \in \mathbb{N}$ ,  $\omega$  be an  $a\mathbb{Z}^\nu$ -invariant state and  $\Phi \in \mathcal{P}_\tau$ .*

(1) The mean entropy

$$s(\omega) = \lim_{n \rightarrow \infty} \left\{ \frac{S(\omega_{\mathcal{A}(\mathcal{C}_{na})})}{|\mathcal{C}_{na}|} \right\} \quad (11.13)$$

exists. It is affine, weak\* upper semicontinuous in  $\omega$  and translation invariant:

$$s(\omega) = s(\tau_k^*(\omega)), \quad (k \in \mathbb{Z}^\nu). \quad (11.14)$$

(2) The mean energy

$$e_\Phi(\omega) = \lim_{n \rightarrow \infty} \left\{ \frac{(\omega(U(\mathcal{C}_{na})))}{|\mathcal{C}_{na}|} \right\} \quad (11.15)$$

exists. It is linear in  $\Phi$ , bounded by  $\|\Phi\|$ , affine and weak\* continuous in  $\omega$ , and translation invariant:

$$e_\Phi(\omega) = e_\Phi(\tau_k^*(\omega)), \quad (k \in \mathbb{Z}^\nu). \quad (11.16)$$

*Proof.* We introduce a new lattice system  $(\mathcal{A}^a, \mathcal{A}^a(\mathbf{I}))$  where the total algebra  $\mathcal{A}^a$  is equal to  $\mathcal{A}$  and its local algebra is  $\mathcal{A}^a(\mathbf{I}) \equiv \mathcal{A}(\cup_{m \in \mathbf{I}}(\mathcal{C}_a + am))$  for each finite subset  $\mathbf{I}$  of  $\mathbb{Z}^\nu$ .

For this new system  $(\mathcal{A}^a, \{\mathcal{A}^a(\mathbf{I})\})$ , we assign its local Hamiltonian

$$H^a(\mathbf{I}) \equiv H(\cup_{m \in \mathbf{I}}(\mathcal{C}_a + am))$$

to each finite  $\mathbf{I}$ , where  $H(\cdot)$  denotes a local Hamiltonian of the original system  $(\mathcal{A}, \{\mathcal{A}(\mathbf{I})\})$ .

If  $\omega$  is an  $a\mathbb{Z}^\nu$ -invariant state of the system  $(\mathcal{A}, \{\mathcal{A}(\mathbf{I})\})$ , then it goes over to a translation invariant state of the new system  $(\mathcal{A}^a, \{\mathcal{A}^a(\mathbf{I})\})$ .

We denote mean entropy and mean energy of  $\omega$  for the system  $(\mathcal{A}^a, \{\mathcal{A}^a(\mathbf{I})\})$  by  $s^a(\omega)$  and  $e_\Phi^a(\omega)$  which are shown to exist by Theorem 10.3 and Theorem 9.5.

Because of the scale change, we have

$$s(\omega) = \lim_{n \rightarrow \infty} \frac{S(\omega_{\mathcal{C}_{na}})}{|\mathcal{C}_{na}|} = |\mathcal{C}_a|^{-1} s^a(\omega), \quad (11.17)$$

$$e_\Phi(\omega) = \lim_{n \rightarrow \infty} \frac{(\omega(U(\mathcal{C}_{na})))}{|\mathcal{C}_{na}|} = |\mathcal{C}_a|^{-1} e_\Phi^a(\omega). \quad (11.18)$$

Hence those properties of mean entropy and mean energy of translation invariant states given in Theorem 10.3 and Theorem 9.5 go over to those for periodic states.

Now we show (11.14) for any  $a\mathbb{Z}^\nu$ -invariant state  $\omega$  and any  $k \in \mathbb{Z}^\nu$ . Due to the  $a\mathbb{Z}^\nu$ -invariance of  $\omega$ , we only have to show the assertion for any  $k \in \mathcal{C}_a$ . For any  $n \in \mathbb{N}$ , we have

$$S(\tau_k^* \omega|_{\mathcal{A}(\mathcal{C}_{na})}) = S(\omega|_{\mathcal{A}(\mathcal{C}_{na+k})}), \quad (11.19)$$

which is to be compared with  $S(\omega|_{\mathcal{A}(\mathcal{C}_{na})})$ .

Since  $k \in \mathcal{C}_a$ , we have

$$\mathcal{C}_{(n-1)a} + a(1, \dots, 1) \subset \mathcal{C}_{na} + k \subset \mathcal{C}_{(n+1)a}. \quad (11.20)$$

By (3.2), (10.5), and the periodicity of  $\omega$ ,

$$\begin{aligned} S(\omega_{\mathcal{A}(\mathcal{C}_{na+k})}) &\leq S(\omega_{\mathcal{A}(\mathcal{C}_{(n-1)a})}) + \{|\mathcal{C}_{na}| - |\mathcal{C}_{(n-1)a}|\} \log 2, \\ S(\omega_{\mathcal{A}(\mathcal{C}_{na+k})}) &\geq S(\omega_{\mathcal{A}(\mathcal{C}_{(n+1)a})}) - \{|\mathcal{C}_{(n+1)a}| - |\mathcal{C}_{na}|\} \log 2. \end{aligned}$$

Due to

$$\lim_{n \rightarrow \infty} \frac{|\mathcal{C}_{na}|}{|\mathcal{C}_{(n-1)a}|} = 1, \quad \lim_{n \rightarrow \infty} \frac{|\mathcal{C}_{na}|}{|\mathcal{C}_{(n+1)a}|} = 1, \quad (11.21)$$

and (11.19), we obtain

$$\begin{aligned} s(\tau_k^* \omega) &= \lim_{n \rightarrow \infty} \frac{S(\omega_{\mathcal{A}(\mathcal{C}_{na+k})})}{|\mathcal{C}_{na}|} \\ &= \lim_{n \rightarrow \infty} \frac{S(\omega_{\mathcal{A}(\mathcal{C}_{na})})}{|\mathcal{C}_{na}|} = s(\omega), \end{aligned}$$

which is the desired equality (11.14).

It remains to show (11.16). Applying the inequality (8.5) to the pair  $\mathbf{I} = (\mathcal{C}_{(n-1)a} + a(1, \dots, 1))$ ,  $\mathbf{J} = (\mathcal{C}_{na} + k) \setminus \{\mathcal{C}_{(n-1)a} + a(1, \dots, 1)\}$  and to the pair  $\mathbf{I} = (\mathcal{C}_{(n-1)a} + a(1, \dots, 1))$ ,  $\mathbf{J} = \mathcal{C}_{na} \setminus \{\mathcal{C}_{(n-1)a} + a(1, \dots, 1)\}$ , we obtain

$$\begin{aligned} \|U(\mathcal{C}_{na}) - U(\mathcal{C}_{na} + k)\| &\leq \|U(\mathcal{C}_{na}) - U(\mathbf{I})\| + \|U(\mathbf{I}) - U(\mathcal{C}_{na} + k)\| \\ &\leq 2\|\Phi\| \{|\mathcal{C}_{na}| - |\mathcal{C}_{(n-1)a}|\}, \end{aligned}$$

where  $I = (\mathcal{C}_{(n-1)a} + a(1, \dots, 1))$ . Hence due to (11.21) and the periodicity of  $\omega$ ,

$$\begin{aligned} e_{\Phi}(\tau_k^* \omega) &= \lim_{n \rightarrow \infty} \frac{\omega(U(\mathcal{C}_{na} + k))}{|\mathcal{C}_{na}|} \\ &= \lim_{n \rightarrow \infty} \frac{\omega(U(\mathcal{C}_{na}))}{|\mathcal{C}_{na}|} = e_{\Phi}(\omega), \end{aligned}$$

which is the desired equality (11.16).  $\square$

Now we resume the proof of Theorem 11.4.

*Proof of Theorem 11.4 (continued).*

Due to  $\Theta$ -evenness of the internal energy  $U(I)$  for every finite  $I \subset \mathbb{Z}^\nu$ , we have

$$\varphi_I^c \in \mathcal{A}(I)_{+,1}^{*\Theta}. \quad (11.22)$$

Let  $a \in \mathbb{N}$ . For distinct  $m \in \mathbb{Z}^\nu$ ,  $\{\mathcal{C}_a + am\}$  are mutually disjoint and their union for all  $m \in \mathbb{Z}^\nu$  is  $\mathbb{Z}^\nu$ .

We apply Theorem 11.2 to the local Gibbs states  $\varphi_{\mathcal{C}_a + am}^c \in \mathcal{A}(\mathcal{C}_a + am)_{+,1}^{*\Theta}$ ,  $m \in \mathbb{Z}^\nu$  and obtain an even product state of  $\mathcal{A}$ , which we denote by  $\varphi_a^c$ .

For any  $k \in \mathbb{Z}^\nu$ ,  $\tau_{ak}^* \varphi_a^c = \varphi_a^c$  by the uniqueness of the product state with the same component states. Thus  $\varphi_a^c$  is an  $a\mathbb{Z}^\nu$ -invariant state.

By using  $\varphi_a^c$  we construct an averaged state  $\widehat{\varphi}_a^c$  which is translation invariant as follows:

$$\widehat{\varphi}_a^c \equiv \sum_{m \in \mathcal{C}_a} \frac{\tau_m^* \varphi_a^c}{|\mathcal{C}_a|} \in \mathcal{A}_{+,1}^{*\tau}. \quad (11.23)$$

We now show (11.12) by taking  $\rho_n = \widehat{\varphi}_n^c$ . By affine dependence of  $s$  and  $e_{\Phi}$  on the space of periodic states in Lemma 11.5,

$$\begin{aligned} s(\widehat{\varphi}_a^c) &= |\mathcal{C}_a|^{-1} \sum_{m \in \mathcal{C}_a} s(\tau_m^* \varphi_a^c), \\ e_{\Phi}(\widehat{\varphi}_a^c) &= |\mathcal{C}_a|^{-1} \sum_{m \in \mathcal{C}_a} e_{\Phi}(\tau_m^* \varphi_a^c). \end{aligned}$$

Due to (11.14) and (11.16), they imply

$$s(\widehat{\varphi}_a^c) = s(\varphi_a^c), \quad (11.24)$$

$$e_{\Phi}(\widehat{\varphi}_a^c) = e_{\Phi}(\varphi_a^c). \quad (11.25)$$

By (11.24), we have

$$\begin{aligned} s(\widehat{\varphi}_a^c) &= s(\varphi_a^c) = \frac{1}{|\mathcal{C}_a|} S(\varphi_{\mathcal{C}_a}^c) \\ &= \frac{1}{|\mathcal{C}_a|} \left\{ \log \text{Tr}_{\mathcal{C}_a} (e^{-\beta U(\mathcal{C}_a)}) + \beta \varphi_a^c(U(\mathcal{C}_a)) \right\}, \end{aligned} \quad (11.26)$$

where the last equality is given by the substitution of an explicit form of the density matrix of the local Gibbs state  $\varphi_{\mathcal{C}_a}^c$  in Definition 7.3.

In order to show (11.12), we now compare  $e_\Phi(\varphi_a^c)$  with  $\frac{1}{|\mathcal{C}_a|}\varphi_a^c(U(\mathcal{C}_a))$  in (11.26). Let  $k \in \mathbb{N}$  and consider the following division of  $\mathcal{C}_{ka}$  as a disjoint union of translates of  $\mathcal{C}_a$ :

$$\mathcal{C}_{ka} = \bigcup_{m \in \mathcal{C}_k} (\mathcal{C}_a + am). \quad (11.27)$$

We give the lexicographic ordering for elements in  $\mathcal{C}_k$  and set

$$\mathcal{C}_{ka}^m \equiv \bigcup_{m' < m} (\mathcal{C}_a + am')$$

for  $m \in \mathcal{C}_k$ . For any  $k \in \mathbb{N}$ ,

$$U(\mathcal{C}_{ka}) - \sum_{m \in \mathcal{C}_k} U(\mathcal{C}_a + am) = \sum_{m \in \mathcal{C}_k} E_{\{\mathcal{C}_{ka} \setminus \mathcal{C}_{ka}^m\}} W(\mathcal{C}_a + am).$$

By  $\|E\| \leq 1$  and the translation covariance ( $\Phi$ -f) of the potential  $\Phi$ , we obtain

$$\begin{aligned} \frac{1}{|\mathcal{C}_{ka}|} \|U(\mathcal{C}_{ka}) - \sum_{m \in \mathcal{C}_k} U(\mathcal{C}_a + am)\| &\leq \frac{1}{|\mathcal{C}_{ka}|} (|\mathcal{C}_k| \cdot \|W(\mathcal{C}_a)\|) \\ &= \frac{\|W(\mathcal{C}_a)\|}{|\mathcal{C}_a|}. \end{aligned} \quad (11.28)$$

Therefore, by (9.1), there exists  $a_0 \in \mathbb{N}$  for any  $\varepsilon > 0$  such that for all  $a > a_0$

$$\left\| \frac{1}{|\mathcal{C}_{ka}|} \left\{ U(\mathcal{C}_{ka}) - \sum_{m \in \mathcal{C}_k} U(\mathcal{C}_a + am) \right\} \right\| < \varepsilon. \quad (11.29)$$

Note that the above  $a_0$  can be taken independent of  $k \in \mathbb{N}$ . For any  $a \in \mathbb{N}$ ,

$$\varphi_a^c(U(\mathcal{C}_a + am)) = \varphi_a^c(U(\mathcal{C}_a)),$$

for any  $m \in \mathbb{Z}^\nu$ , due to the  $a\mathbb{Z}^\nu$ -invariance of  $\varphi_a^c$ . Therefore, we obtain

$$\left| \frac{1}{|\mathcal{C}_{ka}|} \varphi_a^c(U(\mathcal{C}_{ka})) - \frac{1}{|\mathcal{C}_a|} \varphi_a^c(U(\mathcal{C}_a)) \right| < \varepsilon,$$

for  $a > a_0$ . By taking the limit  $k \rightarrow \infty$ , we have

$$\left| e_\Phi(\varphi_a^c) - \frac{1}{|\mathcal{C}_a|} \varphi_a^c(U(\mathcal{C}_a)) \right| < \varepsilon.$$

From this estimate, (11.25) and (11.26), it follows that

$$\left| s(\widehat{\varphi_a^c}) - \beta e_\Phi(\widehat{\varphi_a^c}) - \frac{1}{|\mathcal{C}_a|} \log \mathbf{Tr}_{\mathcal{C}_a}(e^{-\beta U(\mathcal{C}_a)}) \right| < |\beta| \varepsilon,$$

for all  $a \geq a_0$ . This proves (11.12) for  $\rho_n = \widehat{\varphi_n^c}$  in view of (9.22).  $\square$

## 11.4 Variational Principle

**Definition 11.6.** Any translation invariant state  $\varphi$  satisfying

$$P(\beta\Phi) = s(\varphi) - \beta e_\Phi(\varphi) \quad (11.30)$$

(namely, maximizing the functional  $s - \beta e_\Phi$ ) is called a solution of the  $(\Phi, \beta)$ -variational principle (or a translation invariant equilibrium state for  $\Phi$  at the inverse temperature  $\beta$ ). The set of all solutions of the  $(\Phi, \beta)$ -variational principle is denoted by  $\Lambda_{\beta\Phi}$ .

$$\Lambda_{\beta\Phi} \equiv \{\varphi; \varphi \in \mathcal{A}_{+,1}^{*\tau}, P(\beta\Phi) = s(\varphi) - \beta e_\Phi(\varphi)\}. \quad (11.31)$$

*Remark 1.* Since  $\beta e_\Phi(\varphi) = e_{\beta\Phi}(\varphi)$ , the condition  $\varphi \in \Lambda_{\beta\Phi}$  is equivalent to the condition that  $\varphi$  is a solution of the  $(\beta\Phi, 1)$ -variational principle, and hence  $\Lambda_{\beta\Phi}$  is a consistent notation.

*Remark 2.* In the usual physical convention, the functional  $s - \beta e_\Phi$  is  $-\beta$  times the free energy functional.

**Theorem 11.7.** For any  $\Phi \in \mathcal{P}_\tau$  and  $\beta \in \mathbb{R}$ , there exists a solution  $\varphi (\in \mathcal{A}_{+,1}^{*\tau})$  of  $(\Phi, \beta)$ -variational principle, namely,

$$\Lambda_{\beta\Phi} \neq \emptyset.$$

*Proof.*  $\{\widehat{\varphi}_a^c\}$  in the proof of Theorem 11.4 has an accumulation point in  $\mathcal{A}_{+,1}^{*\tau}$  by the weak\*-compactness of  $\mathcal{A}_{+,1}^{*\tau}$ . Let  $\varphi$  be any such accumulation point. By the proof of Theorem 11.4, the weak\* continuity of  $e_\Phi$  and the weak\* upper semicontinuity of  $s$  in  $\omega$ , the state  $\varphi$  satisfies

$$P(\beta\Phi) = \lim_{a \rightarrow \infty} \left( s(\widehat{\varphi}_a^c) - \beta e_\Phi(\widehat{\varphi}_a^c) \right) \leq s(\varphi) - \beta e_\Phi(\varphi). \quad (11.32)$$

By (11.10), we obtain (11.30) □

Our Fermion algebra  $\mathcal{A}$  is not asymptotically abelian with respect to the lattice translations, but if  $\omega$  is translation invariant state of  $\mathcal{A}$ , it is well known that the pair  $(\mathcal{A}, \omega)$  is  $\mathbb{Z}^\nu$ -abelian and that  $\omega$  is automatically even (see, for example, Example 5.2.21 in [17]). From this consideration and Theorem 11.4, we obtain the following result, which corresponds to Theorem 6.2.44 in [17] in the case of quantum spin lattice systems, by the same argument as for that theorem.

For a convex set  $K$ , we denote the set of extremal points of  $K$  by  $\mathcal{E}(K)$ .

**Proposition 11.8.** For  $\Phi \in \mathcal{P}_\tau$  and  $\beta \in \mathbb{R}$ ,  $\Lambda_{\beta\Phi}$  is a simplex with  $\mathcal{E}(\Lambda_{\beta\Phi}) \subset \mathcal{E}(\mathcal{A}_{+,1}^{*\tau})$  and the unique barycentric decomposition of each  $\varphi$  in  $\Lambda_{\beta\Phi}$  coincides with its unique ergodic decomposition.

## 12 Equivalence of Variational Principle and KMS Condition

Among 5 steps for establishing the equivalence stated in the title (which are described in § 1), Step (1) “KMS condition  $\Rightarrow$  Gibbs condition” is obtained in Theorem 7.5 in § 7.4, Step (4) “dKMS condition on  $\mathcal{A}_o \Rightarrow$  dKMS condition on  $D(\delta_\alpha)$ ” is obtained in Corollary 6.7, and Step (5) “dKMS condition on  $D(\delta_\alpha) \Rightarrow$  KMS condition” is stated in Theorem 6.4.

In this section, we complete the remaining two steps of proof by showing Step (2) “Gibbs condition  $\Rightarrow$  Variational principle” in § 12.1 and Step (3) “Variational principle  $\Rightarrow$  dKMS condition on  $\mathcal{A}_o$ ” in § 12.3. As a preparation for the latter, some tools of convex analysis is gathered in § 12.2.

### 12.1 Variational Principle from Gibbs Condition

**Proposition 12.1.** *For  $\Phi \in \mathcal{P}_\tau$ , each translation invariant state  $\varphi$  satisfying  $(\Phi, \beta)$ -Gibbs condition is a solution of the  $(\Phi, \beta)$ -variational principle.*

*Proof.* We follow the method of proof in [6]. The Gibbs condition for  $\varphi$  implies

$$\left[ \varphi^{\beta W(I)} \right] \Big|_{\mathcal{A}(I)} = \varphi_I^c \quad (12.1)$$

for every finite subset  $I$ , where  $\varphi_I^c$  is given by (7.14), and  $[\varphi^{\beta W(I)}]$  denotes the normalization of  $\varphi^{\beta W(I)}$  given by (7.8).

By (11.8) with  $\omega$  replaced by  $\varphi$ , we have

$$\begin{aligned} S(\varphi_I^c, \varphi_I) &= -\widehat{S}(\varphi_I) + \beta\varphi(U(I)) + \log \tau(e^{-\beta U(I)}) \\ &= -S(\varphi_I) + \beta\varphi(U(I)) + \log \mathbf{Tr}_I(e^{-\beta U(I)}). \end{aligned} \quad (12.2)$$

Since relative entropy is non-negative and is monotone non-increasing under restriction of states, it follows that

$$0 \leq S(\varphi_I^c, \varphi_I) \leq S([\varphi^{\beta W(I)}], \varphi).$$

By (7.8), (7.10) and (7.9), we have

$$S([\varphi^{\beta W(I)}], \varphi) = \log(\varphi^{\beta W(I)}(\mathbf{1})) - \varphi(\beta W(I)) \leq 2\|\beta W_I\|.$$

From these estimates and (12.2), it follows that

$$0 \leq S(\varphi_I^c, \varphi_I) = -S(\varphi_I) + \beta\varphi(U(I)) + \log \mathbf{Tr}_I(e^{-\beta U(I)}) \leq 2\|\beta W_I\|.$$

(Up to this point, the assumption of translation invariance of  $\varphi$  is irrelevant.)

We now divide the above inequality by  $|I|$  and take the van Hove limit  $I \rightarrow \infty$ . Then by the translation invariance of  $\varphi$  and (9.1), we obtain

$$s(\varphi) - \beta e_\Phi(\varphi) = P(\beta\Phi),$$

which completes the proof.  $\square$

Combining this proposition with Theorem 7.5, we immediately obtain the following.

**Corollary 12.2.** *Let  $\alpha_t$  be a dynamics of  $\mathcal{A}$  satisfying the Assumptions (II) and (IV) in § 5 and  $\Phi$  be the (translation covariant) standard potential uniquely corresponding to this  $\alpha_t$ . If  $\varphi$  is a translation invariant  $(\alpha_t, \beta)$ -KMS state of  $\mathcal{A}$ , then  $\varphi$  is a solution of the  $(\Phi, \beta)$ -variational principle.*

We have now completed the proof of Theorem A.

## 12.2 Some Tools of Convex Analysis

We use the pressure functional  $\Phi \in \mathcal{P}_\tau \mapsto P(\Phi) \in \mathbb{R}$ , which is a norm continuous convex function on the Banach space  $\mathcal{P}_\tau$  due to Corollary 9.4.

A continuous linear functional  $\alpha \in \mathcal{P}_\tau^*$  (the dual of  $\mathcal{P}_\tau$ ) is called a tangent of the functional  $P$  at  $\Phi \in \mathcal{P}_\tau$  if it satisfies

$$P(\Phi + \Psi) \geq P(\Phi) + \alpha(\Psi) \quad (12.3)$$

for all  $\Psi \in \mathcal{P}_\tau$ .

**Proposition 12.3.** *For any solution  $\varphi$  of the  $(\Phi, 1)$ -variational principle, define*

$$\alpha_\varphi(\Psi) \equiv -e_\Psi(\varphi) \quad (12.4)$$

*for all  $\Psi \in \mathcal{P}_\tau$ . Then  $\alpha_\varphi$  is a tangent of  $P$  at  $\Phi$ .*

*Proof.* By linear dependence (9.26) of  $e_\Psi$  on  $\Psi$ ,  $\alpha_\varphi$  is a linear functional on  $\mathcal{P}_\tau$ . Due to  $|e_\Psi(\varphi)| \leq \|\Psi\|$  given by (9.28), we have  $\alpha_\varphi \in \mathcal{P}_\tau^*$ . Due to the variational inequality (11.10),

$$\begin{aligned} P(\Phi + \Psi) &\geq s(\varphi) - e_{\Phi + \Psi}(\varphi) \\ &= s(\varphi) - e_\Phi(\varphi) - e_\Psi(\varphi) \\ &= P(\Phi) + \alpha_\varphi(\Psi) \end{aligned}$$

for all  $\Psi \in \mathcal{P}_\tau$ , where the last equality is due to the assumption that  $\varphi$  is a solution of the  $(\Phi, 1)$ -variational principle.  $\square$

(We will establish the bijectivity between solutions of the  $(\Phi, \beta)$ -variational principle and tangents of  $P$  at  $\beta\Phi$  through (12.4) in Theorem 12.10.)

Since  $P(\Phi + k\Psi)$  is a convex continuous function of  $k \in \mathbb{R}$  for any fixed  $\Phi, \Psi \in \mathcal{P}_\tau$ , there exist its right and left derivatives at  $k = 0$ ,

$$(D_\Psi^\pm P)(\Phi) = \lim_{k \rightarrow \pm 0} \frac{P(\Phi + k\Psi) - P(\Phi)}{k}.$$

By the convexity of  $P$ ,

$$(D_\Psi^+ P)(\Phi) \geq (D_\Psi^- P)(\Phi).$$

If and only if they coincide,  $P(\Phi + k\Psi)$  is differentiable at  $k = 0$ . Then we define

$$(D_\Psi P)(\Phi) = (D_\Psi^+ P)(\Phi) = (D_\Psi^- P)(\Phi). \quad (12.5)$$

The derivatives  $(D_\Psi^\pm P)(\Phi)$  and hence  $(D_\Psi P)(\Phi)$  (when it exists) satisfy

$$\begin{aligned} |(D_{\Psi_1}^\pm P)(\Phi) - (D_{\Psi_2}^\pm P)(\Phi)| &\leq \|\Psi_1 - \Psi_2\|, \\ |(D_{\Psi_1} P)(\Phi) - (D_{\Psi_2} P)(\Phi)| &\leq \|\Psi_1 - \Psi_2\|, \end{aligned} \quad (12.6)$$

as is shown by the following computation in the limit  $k \rightarrow \pm 0$ .

$$\begin{aligned}
& \left| \frac{\{P(\Phi + k\Psi_1) - P(\Phi)\} - \{P(\Phi + k\Psi_2) - P(\Phi)\}}{k} \right| \\
&= \left| \frac{P(\Phi + k\Psi_1) - P(\Phi + k\Psi_2)}{k} \right| \\
&\leq |k|^{-1} \|k(\Psi_1 - \Psi_2)\| = \|\Psi_1 - \Psi_2\|,
\end{aligned}$$

where (9.23) is used for the inequality. If (12.5) holds for all  $\Psi$ , then  $P$  is said to be differentiable at  $\Phi$ . Let  $\mathcal{P}_\tau^1$  be the set of all  $\Phi \in \mathcal{P}_\tau$  where  $P$  is differentiable.

**Proposition 12.4.** *If  $\Phi \in \mathcal{P}_\tau^1$ ,*

$$\alpha_\Phi(\Psi) = (D_\Psi P)(\Phi), \quad (\Psi \in \mathcal{P}_\tau), \quad (12.7)$$

*defines an  $\alpha_\Phi \in \mathcal{P}_\tau^*$  which is the unique tangent of  $P$  at  $\Phi$ . Then any solution  $\varphi$  of  $(\Phi, 1)$ -variational principle satisfies*

$$\alpha_\Phi(\Psi) = \alpha_\varphi(\Psi), \quad (12.8)$$

*for all  $\Psi \in \mathcal{P}_\tau$ , where  $\alpha_\varphi$  is given by (12.4).*

*Proof.* By Theorem 11.7, there is a solution  $\varphi$  of the  $(\Phi, 1)$ -variational principle and, by Proposition 12.3,  $\alpha_\varphi$  is a tangent of  $P$  at  $\Phi$ .

Let  $\alpha'$  be any tangent of  $P$  at  $\Phi \in \mathcal{P}_\tau^1$ . We have for  $k > 0$

$$\begin{aligned}
P(\Phi + k\Psi) &\geq P(\Phi) + k\alpha'(\Psi), \\
P(\Phi - k\Psi) &\geq P(\Phi) - k\alpha'(\Psi).
\end{aligned}$$

Hence

$$\begin{aligned}
(D_\Psi^+ P)(\Phi) &= \lim_{k \rightarrow +0} \frac{P(\Phi + k\Psi) - P(\Phi)}{k} \geq \alpha'(\Psi), \\
(D_\Psi^- P)(\Phi) &= \lim_{k \rightarrow +0} \frac{P(\Phi - k\Psi) - P(\Phi)}{(-k)} \leq \alpha'(\Psi).
\end{aligned}$$

By (12.5) for  $\Phi \in \mathcal{P}_\tau^1$ , we obtain

$$\alpha'(\Psi) = (D_\Psi P)(\Phi).$$

Then  $\alpha'$  is unique and (12.8) holds.  $\square$

**Lemma 12.5.** *For each  $A \in \mathcal{A}_\circ$  such that  $A = A^* = \Theta(A)$ , there exists  $\Psi_A \in \mathcal{P}_\tau^f$  such that*

$$e_{\Psi_A}(\varphi) = \varphi(A) - \tau(A) \quad (12.9)$$

*for all translation invariant states  $\varphi$ .*



*Proof.* Let  $A = A^* = \Theta(A) \in \mathcal{A}(I)$  for some finite  $I$  and

$$A_1 \equiv A - \tau(A)\mathbf{1} (\in \mathcal{A}(I)).$$

Since  $E_{I^c}(A_1) = \tau(A_1)\mathbf{1} = 0$ , there exists a unique decomposition

$$A_1 = \sum_{\substack{J \subset I \\ J \neq \emptyset}} A(J), \quad A(J) \in \mathcal{A}(J), \quad (12.10)$$

$$E_K(A(J)) = 0 \quad \text{for } K \not\subset J. \quad (12.11)$$

To show these formulae, let

$$A(J) = \sum_{K \subset J} (-1)^{|J|-|K|} E_K(A_1) \quad (12.12)$$

for all non-empty  $J \subset I$ , a formula in parallel with (5.16). Then

$$E_J(A_1) = \sum_{\substack{K \subset J \\ K \neq \emptyset}} A(K) \quad (12.13)$$

for  $J \subset I$  by exactly the same computation as Step 1 of the proof of Lemma 5.9. (When  $J = \emptyset$ , the right-hand side is interpreted as 0 and  $E_\emptyset(A_1) = 0$ .) We have

$$A(J)^* = A(J) = \Theta(A(J)) \in \mathcal{A}(J), \quad (12.14)$$

because  $A(J)$  is a real linear combination of  $E_K(A_1)$ ,  $K \subset J$ , and all  $E_K(A_1)$  satisfy the same equation. We note that Step 4 of Lemma 5.9 uses only the following properties of  $U(K)$

$$U(\emptyset) = 0, \quad \tau(U(K)) = 0, \quad E_K(U(J)) = U(K),$$

for  $K \subset J \subset I$ , and that all of them are satisfied also by  $E_K(A_1)$ . Therefore, (12.11) follows from the same argument as Step 4 of Lemma 5.9.

We now construct  $\Psi_J \in \mathcal{P}_\tau^f$  for each  $A(J)$  in (12.10) such that

$$e_{\Psi_J}(\varphi) = \varphi(A(J)) \quad (12.15)$$

for all translation invariant states  $\varphi$ . Then by linear dependence of  $e_\Psi$  on  $\Phi \in \mathcal{P}_\tau$ , we obtain for  $\Psi = \sum_{J \subset I} \Psi_J$  the desired relation (12.9):

$$e_\Psi(\varphi) = \sum_{J \subset I} e_{\Psi_J}(\varphi) = \sum_{J \subset I} \varphi(A(J)) = \varphi(A_1) = \varphi(A) - \tau(A).$$

We define a potential  $\Psi_J$  for each  $J \subset I$ ,  $J \neq \emptyset$  by

$$\begin{aligned} \Psi_J(J+m) &= \tau_m(A(J)), \quad (m \in \mathbb{Z}^\nu), \\ \Psi_J(K) &= 0 \quad \text{if } K \text{ is not a translate of } J. \end{aligned} \quad (12.16)$$

Due to the property (12.14) and (12.11),  $\Psi_J$  belongs to  $\mathcal{P}_\tau^f$ . We compute

$$\begin{aligned} \frac{1}{|\mathcal{C}_a|} \varphi(U_{\Psi_J}(\mathcal{C}_a)) &= \frac{1}{|\mathcal{C}_a|} \varphi\left(\sum \{\Psi_J(J+m); J+m \subset \mathcal{C}_a\}\right) \\ &= \frac{N_a}{|\mathcal{C}_a|} \varphi(A(J)), \end{aligned}$$

where  $N_a$  is the number of  $m$  such that  $J + m \subset \mathcal{C}_a$ .

We now show that  $\frac{N_a}{|\mathcal{C}_a|} \rightarrow 1$  as  $a \rightarrow \infty$ . Since  $J + m \subset \mathcal{C}_a$  is equivalent to  $J \subset \mathcal{C}_a - m$ ,  $N_a$  is the same as  $l(a, J)$  (the number of translates of  $\mathcal{C}_a$  containing  $J$ ). By (8.11),

$$\lim_{a \rightarrow \infty} \frac{N_a}{|\mathcal{C}_a|} = \lim_{a \rightarrow \infty} \frac{l(a, J)}{|\mathcal{C}_a|} = 1$$

Hence

$$e_{\Psi_J}(\varphi) = \lim_{a \rightarrow \infty} \frac{1}{|\mathcal{C}_a|} \varphi(U_{\Psi_J}(\mathcal{C}_a)) = \varphi(A(J)).$$

□

**Corollary 12.6.** *If  $\varphi_1$  and  $\varphi_2$  are distinct solutions of  $(\Phi, 1)$ -variational principle for  $\Phi \in \mathcal{P}_\tau$ , then the corresponding tangent of  $P$  at  $\Phi$  are distinct, that is,  $\alpha_{\varphi_1} \neq \alpha_{\varphi_2}$ .*

*Proof.* If  $\varphi_1 \neq \varphi_2$ , there exists an  $A \in \mathcal{A}_0$  such that  $\varphi_1(A) \neq \varphi_2(A)$ . Let  $A_\pm = \frac{1}{2}(A \pm \Theta(A))$ . Then  $A = A_+ + A_-$ . Since  $\varphi_1$  and  $\varphi_2$  are translation invariant, both of them are  $\Theta$ -even, and hence  $\varphi_1(A_-) = \varphi_2(A_-) = 0$ . Thus  $\varphi_1(A_+) \neq \varphi_2(A_+)$ . So we may assume that  $\Theta(A) = A$ . Let  $A_1 = \frac{1}{2}(A + A^*)$ ,  $A_2 = \frac{1}{2i}(A - A^*)$ ,  $A = A_1 + iA_2$ . Then either  $\varphi_1(A_1) \neq \varphi_2(A_1)$  or  $\varphi_1(A_2) \neq \varphi_2(A_2)$ . Since  $A_1^* = A_1$  and  $A_2^* = A_2$ , we may assume  $A = A^* = \Theta(A)$ . Let  $\Psi_A \in \mathcal{P}_\tau^f$  be given as in Lemma 12.5 for this  $A \in \mathcal{A}_0$ . Then

$$\begin{aligned} \alpha_{\varphi_1}(\Psi_A) &= -e_{\Psi_A}(\varphi_1) = -\varphi_1(A) + \tau(A) \\ &\neq -\varphi_2(A) + \tau(A) = -e_{\Psi_A}(\varphi_2) = \alpha_{\varphi_2}(\Psi_A). \end{aligned}$$

Hence  $\alpha_{\varphi_1} \neq \alpha_{\varphi_2}$ . □

**Corollary 12.7.** *For  $\Phi \in \mathcal{P}_\tau^1$ , a solution of  $(\Phi, 1)$ -variational principle is unique.*

*Proof.* This follows from Proposition 12.4 and Corollary 12.6. □

We will use the following result in the proof of Theorem 12.11.

**Theorem 12.8.** (1) *The set  $\mathcal{P}_\tau^1$  of points of unique tangent of  $P$  is residual (an intersection of a countable number of dense open sets) and dense in  $\mathcal{P}_\tau$ .*  
(2) *For any  $\Phi \in \mathcal{P}_\tau$ , any tangent of  $P$  at  $\Phi$  is contained in the weak\* closed convex hull of the set  $\Gamma(\Phi)$  which is defined by*

$$\begin{aligned} \Gamma(\Phi) \equiv \{ \alpha \in \mathcal{P}_\tau^*; \text{ there exists a net } \Phi_\gamma \in \mathcal{P}_\tau^1 \text{ such that } \|\Phi_\gamma - \Phi\| \rightarrow 0, \\ \text{and } \alpha_{\Phi_\gamma} \rightarrow \alpha \text{ in the weak* topology of } \mathcal{P}_\tau^* \}, \end{aligned} \quad (12.17)$$

where  $\alpha_{\Phi_\gamma}$  is the unique tangent of  $P$  at  $\Phi_\gamma$ .

*Proof.* (1) is Mazur's theorem [31].

(2) is Theorem 1 of [26] where the function  $f$  is to be set  $f(\Psi) = P(\Phi + \Psi)$  for our purpose. The proof in [26] is by the Hahn-Banach theorem. (Separability of  $\mathcal{P}_\tau$  given by Corollary 8.13 is needed for both (1) and (2).)  $\square$

We now show a bijective correspondence between solutions of the  $(\Phi, \beta)$ -variational principle and tangents of  $P$  at  $\beta\Phi$ . We first prove a lemma about stability of solutions of the variational principle under the limiting procedure in (12.17).

**Lemma 12.9.** *Let  $\{\Phi_\gamma\}$  be a net in  $\mathcal{P}_\tau$  and  $\{\varphi_\gamma\}$  be a net consisting of a solution  $\varphi_\gamma$  of the  $(\Phi_\gamma, \beta_\gamma)$ -variational principle for each index  $\gamma$  such that*

$$\begin{aligned} \|\Phi_\gamma - \Phi\| &\rightarrow 0, \quad (\Phi \in \mathcal{P}_\tau), \quad \beta_\gamma \rightarrow \beta \in \mathbb{R}, \\ \varphi_\gamma &\rightarrow \varphi \in \mathcal{A}_{+,1}^{*\tau} \text{ in the weak* topology of } \mathcal{A}_{+,1}^{*\tau}. \end{aligned}$$

*Then  $\varphi$  is a solution of the  $(\Phi, \beta)$ -variational principle.*

*Proof.* By the norm continuity (9.23) of  $P$ , the weak\* upper semicontinuity of  $s$  (Theorem 10.3) and the continuous dependence of  $e_\Phi(\varphi)$  on  $\Phi$  in the norm topology (uniformly in  $\varphi$ ) and on  $\varphi$  in the weak\* topology (Theorem 9.5), we have

$$\begin{aligned} P(\beta\Phi) &= \lim_\gamma P(\beta_\gamma\Phi_\gamma), \\ s(\varphi) &\geq \limsup_\gamma s(\varphi_\gamma), \\ e_\Phi(\varphi) &= \lim_\gamma e_{\Phi_\gamma}(\varphi_\gamma). \end{aligned}$$

Since,  $\varphi_\gamma$  is a solution of the  $(\Phi_\gamma, \beta_\gamma)$ -variational principle, we have

$$P(\beta_\gamma\Phi_\gamma) = s(\varphi_\gamma) - \beta_\gamma e_{\Phi_\gamma}(\varphi_\gamma).$$

Hence

$$P(\beta\Phi) \leq s(\varphi) - \beta e_\Phi(\varphi).$$

By the variational inequality (11.10), we have

$$P(\beta\Phi) = s(\varphi) - \beta e_\Phi(\varphi)$$

.

$\square$

**Theorem 12.10.** *For any  $\Phi \in \mathcal{P}_\tau$  and  $\beta \in \mathbb{R}$ , there exists a bijective affine map  $\varphi \mapsto \alpha_\varphi$  from the set  $\Lambda_{\beta\Phi}$  to the set of all tangents of the functional  $P$  at  $\beta\Phi$ , given by*

$$\alpha_\varphi(\Psi) = -e_\Psi(\varphi), \quad \Psi \in \mathcal{P}_\tau. \quad (12.18)$$

*Proof.* By *Remark 1* after Definition 11.6, all solutions of the  $(\Phi, \beta)$ - and  $(\beta\Phi, 1)$ -variational principle coincide. Furthermore, if  $\varphi$  is a solution of the  $(\Phi, \beta)$ -variational principle, then

$$\begin{aligned} P(\beta\Phi + \Psi) &\geq s(\varphi) - e_{\beta\Phi + \Psi}(\varphi) \\ &= s(\varphi) - \beta e_{\Phi}(\varphi) - e_{\Psi}(\varphi) \\ &= P(\beta\Phi) + \alpha_{\varphi}(\Psi). \end{aligned}$$

Namely  $\alpha_{\varphi}$  is a tangent of  $P$  at  $\beta\Phi$ , exactly the same statement as for a solution  $\varphi$  of the  $(\beta\Phi, 1)$ -variational principle. Therefore, it is enough to prove the case of  $\beta = 1$ .

The map  $\varphi \mapsto \alpha_{\varphi}$  is an affine map from the set of all solutions of  $(\Phi, 1)$ -variational principle into the set of all tangents of  $P$  at  $\Phi$ . The map is injective by Corollary 12.6. To show the surjectivity of the map, let  $\alpha$  be a tangent of  $P$  at  $\Phi$ . By Theorem 12.8, there exists a net  $\Phi_{\gamma} \in \mathcal{P}_{\tau}^1$  such that  $\|\Phi_{\gamma} - \Phi\| \rightarrow 0$ , and  $\alpha_{\Phi_{\gamma}} \rightarrow \alpha$  in the weak\* topology of  $\mathcal{P}_{\tau}^*$ , where  $\alpha_{\Phi_{\gamma}}$  is the unique tangent of  $P$  at  $\Phi_{\gamma}$ . By Theorem 11.7, there exists a solution  $\varphi_{\gamma}$  of the  $(\Phi_{\gamma}, 1)$ -variational principle. By Proposition 12.3,  $\alpha_{\varphi_{\gamma}}$  is a tangent of  $P$  at  $\Phi_{\gamma}$  and hence must coincide with the unique tangent  $\alpha_{\Phi_{\gamma}}$ . Due to the weak\* compactness of  $\mathcal{A}_{+,1}^{*\tau}$ , there exists a subnet  $\{\varphi_{\gamma(\mu)}\}_{\mu}$  which converges to some  $\varphi \in \mathcal{A}_{+,1}^{*\tau}$ . By Lemma 12.9 and by  $\|\Phi_{\gamma(\mu)} - \Phi\| \rightarrow 0$ ,  $\varphi$  must be a solution of the  $(\Phi, 1)$ -variational principle. Furthermore, for any  $\Psi \in \mathcal{P}_{\tau}$ , we have

$$\begin{aligned} \alpha_{\varphi}(\Psi) &= -e_{\Psi}(\varphi) \\ &= -\lim_{\mu} e_{\Psi}(\varphi_{\gamma(\mu)}) = -\lim_{\mu} \alpha_{\gamma(\mu)}(\Psi) \\ &= \alpha(\Psi). \end{aligned}$$

Hence  $\alpha = \alpha_{\varphi}$  and the map  $\varphi \rightarrow \alpha_{\varphi}$  is surjective.  $\square$

### 12.3 Differential KMS Condition from Variational Principle

In this subsection, we give a proof for Step 3.

**Theorem 12.11.** *Let  $\Phi \in \mathcal{P}_{\tau}$  and  $\varphi$  be a translation invariant state. If  $\varphi$  is a solution of  $(\Phi, \beta)$ -variational principle, then  $\varphi$  is a  $(\delta_{\Phi}, \beta)$ -dKMS state, where  $\delta_{\Phi} \in \Delta(\mathcal{A}_{\circ})$  corresponds to  $\Phi$  by the bijective linear map of Corollary 8.5.*

*Remark.* We note that this theorem holds for any  $\Phi \in \mathcal{P}_{\tau}$  without any further assumption on  $\Phi$  and we do not need  $\alpha_t$ . Note that the domain  $D(\delta_{\Phi})$  is  $\mathcal{A}_{\circ}$  by definition.

First we present some estimate needed in the proof of this theorem in the form of the following lemma.

**Lemma 12.12.** *Let  $I$  and  $J$  be finite subsets of  $\mathbb{Z}^{\nu}$ . If  $A \in \mathcal{A}(J)$ , then*

$$\left\| [U(I), A] \right\| \leq 2\|\Phi\| \cdot \|A\| \cdot |I \cap J|. \quad (12.19)$$

*Proof.* Let  $I_0$  be the complement of  $I \cap J$  in  $I$ . Then  $I_0 \cap J = \emptyset$  and hence  $U(I_0)$  commutes with  $A (\in \mathcal{A}(J))$  due to  $U(I_0) \in \mathcal{A}(I_0)_+ \subset \mathcal{A}(J)'$ . Since  $I_0$  and  $I \cap J$  are disjoint and have the union  $I$ , the following computation proves (12.19).

$$\begin{aligned} \|[U(I), A]\| &= \|[U(I) - U(I_0), A]\| \\ &\leq 2\|U(I) - U(I_0)\| \|A\| \\ &\leq 2\|\Phi\| \cdot \|A\| \cdot |I \cap J|, \end{aligned}$$

where the last inequality is due to (8.5).  $\square$

*Proof of Theorem 12.11 :*

We note that  $(\Phi, \beta)$ -variational principle and  $(\beta\Phi, 1)$ -variational principle are the same and  $(\delta_\Phi, \beta)$ -dKMS condition and  $(\delta_{\beta\Phi}, 1)$ -dKMS condition are the same. By taking  $\beta\Phi$  as a new  $\Phi$ , we only have to prove the case  $\beta = 1$ .

Let  $\widehat{\varphi}_a^c$  be the translation invariant state defined by (11.23) in the proof of Theorem 11.4. Let  $\varphi$  be any accumulation point of  $\{\widehat{\varphi}_a^c\}_{a \in \mathbb{N}}$ . Then this  $\varphi$  is a solution of  $(\Phi, 1)$ -variational principle as shown in Theorem 11.7.

For the moment, let us assume  $\Phi \in \mathcal{P}_\tau^1$  (the set of  $\Phi \in \mathcal{P}_\tau$  where  $P$  is differentiable, defined in § 12.2). Due to the assumption  $\Phi \in \mathcal{P}_\tau^1$ , any accumulation point of  $\{\widehat{\varphi}_a^c\}_{a \in \mathbb{N}}$  coincides with the unique solution  $\varphi$  of  $(\Phi, 1)$ -variational principle, and hence

$$\lim_{a \rightarrow \infty} \widehat{\varphi}_a^c = \varphi. \quad (12.20)$$

We now prove that the above  $\varphi$  satisfies the conditions (C-1) and (C-2) of Definition 6.3 for each  $A \in \mathcal{A}_0$  by using (12.20).

Let  $A \in \mathcal{A}(I)$  for a finite subset  $I$  of  $\mathbb{Z}^\nu$ . Suppose  $\mathcal{C}_a - k \supset I$  ( $a \in \mathbb{N}$ ,  $k \in \mathbb{Z}^\nu$ ). Since  $\tau_k^* \varphi_a^c$  is the  $(\text{Ad } e^{itU(\mathcal{C}_a - k)}, 1)$ -KMS state on  $\mathcal{A}(\mathcal{C}_a - k)$ , we have

$$\text{Re}(\tau_k^* \varphi_a^c)(A^*[iU(\mathcal{C}_a - k), A]) = 0, \quad (12.21)$$

$$\text{Im}(\tau_k^* \varphi_a^c)(A^*[iU(\mathcal{C}_a - k), A]) \geq S(\tau_k^* \varphi_a^c(AA^*), \tau_k^* \varphi_a^c(A^*A)). \quad (12.22)$$

Our strategy of the proof is to replace  $\tau_k^* \varphi_a^c$  and  $[iU(\mathcal{C}_a - k), A]$  by  $\varphi$  and  $\delta_\Phi(A)$ , respectively, by using an approximation argument.

By (4.23) for  $J \nearrow \mathbb{Z}^\nu$ , there exists a finite subset  $J_\varepsilon$  of  $\mathbb{Z}^\nu$  for any given  $\varepsilon > 0$  such that

$$\|H(I) - E_J(H(I))\| < \varepsilon, \quad (12.23)$$

for all  $J \supset J_\varepsilon$ .

Let  $b$  be sufficiently large so that there exists a translate  $\mathcal{C}_b - l_0$  of  $\mathcal{C}_b$  containing both  $I$  and  $J_\varepsilon$ .

We will use the following convenient expression for  $\widehat{\varphi}_a^c (\in \mathcal{A}_{+,1}^{\tau,*})$  which is equivalent to (11.23):

$$\widehat{\varphi}_a^c = \tau_l^* \widehat{\varphi}_a^c = \sum_{m \in \mathcal{C}_a} \frac{\tau_{l+m}^* \varphi_a^c}{|\mathcal{C}_a|} = \sum_{m \in (\mathcal{C}_a + l)} \frac{\tau_m^* \varphi_a^c}{|\mathcal{C}_a|}, \quad (12.24)$$

for any  $l \in \mathbb{Z}^\nu$ . We will take  $l = l_0$ . We divide  $\mathcal{C}_a + l_0$  into the following two disjoint subsets when  $a > b$ :

$$\mathcal{C}_1 \equiv \mathcal{C}_{a-b} + l_0, \quad \mathcal{C}_2 \equiv (\mathcal{C}_a + l_0) \setminus \mathcal{C}_1. \quad (12.25)$$

Then

$$\mathcal{C}_a - k \supset \mathcal{C}_b - l_0 \supset \mathbf{I} \cup \mathbf{J}_\varepsilon \quad (12.26)$$

if  $k \in \mathcal{C}_1$ , while

$$\frac{|\mathcal{C}_2|}{|\mathcal{C}_a|} = \left(1 - \frac{|\mathcal{C}_{a-b}|}{|\mathcal{C}_a|}\right) \rightarrow 0, \quad (12.27)$$

as  $a \rightarrow \infty$ .

For  $k \in \mathcal{C}_1$ ,  $A(\in \mathcal{A}(\mathbf{I}))$  belongs to  $\mathcal{A}(\mathcal{C}_a - k)$  due to  $\mathbf{I} \subset \mathcal{C}_a - k$ . By using the general property of the conditional expectation, we have

$$\begin{aligned} i[U(\mathcal{C}_a - k), A] &= iE_{\mathcal{C}_a - k}([H(\mathcal{C}_a - k), A]) = iE_{\mathcal{C}_a - k}([H(\mathbf{I}), A]) \\ &= i[E_{\mathcal{C}_a - k}(H(\mathbf{I})), A]. \end{aligned}$$

By (12.23) for  $\mathbf{J} = \mathcal{C}_a - k(\supset \mathbf{J}_\varepsilon)$ , this implies

$$\|i[H(\mathbf{I}), A] - i[U(\mathcal{C}_a - k), A]\| < 2\varepsilon\|A\|.$$

Noting that  $\delta_\Phi(A) = i[H(\mathbf{I}), A]$ , we have

$$\|\delta_\Phi(A) - i[U(\mathcal{C}_a - k), A]\| < 2\varepsilon\|A\|. \quad (12.28)$$

It follows from (12.21) and (12.28) that

$$\left| \mathbf{Re}(\tau_k^* \varphi_a^c)(A^* \delta_\Phi(A)) \right| < 2\varepsilon\|A\|^2 \quad (12.29)$$

for  $k \in \mathcal{C}_1$ . For  $k \in \mathcal{C}_2$ , we use the following obvious estimate.

$$\left| \mathbf{Re}(\tau_k^* \varphi_a^c)(A^* \delta_\Phi(A)) \right| < \|A^* \delta_\Phi(A)\|. \quad (12.30)$$

Substituting (12.29) and (12.30) into (12.24), we obtain

$$\begin{aligned} & \left| \mathbf{Re} \widehat{\varphi_a^c}(A^* \delta_\Phi(A)) \right| \\ & \leq \left| \mathbf{Re} \left( \sum_{k \in \mathcal{C}_1} \frac{1}{|\mathcal{C}_a|} \tau_k^* \varphi_a^c \right) (A^* \delta_\Phi(A)) \right| + \left| \mathbf{Re} \left( \sum_{k \in \mathcal{C}_2} \frac{1}{|\mathcal{C}_a|} \tau_k^* \varphi_a^c \right) (A^* \delta_\Phi(A)) \right| \\ & \leq 2\varepsilon\|A\|^2 + \frac{|\mathcal{C}_2|}{|\mathcal{C}_a|} \|A^* \delta_\Phi(A)\|. \end{aligned}$$

Taking the limit  $a \rightarrow \infty$  and using (12.27), we obtain

$$\left| \mathbf{Re} \varphi(A^* \delta_\Phi(A)) \right| \leq 2\varepsilon\|A\|^2.$$

Due to arbitrariness of  $\varepsilon > 0$ , we obtain

$$\left| \mathbf{Re} \varphi(A^* \delta_\Phi(A)) \right| = 0. \quad (12.31)$$

Hence the condition (C-1) holds.

By (12.22) and (12.28), we have the following inequality for  $k \in \mathcal{C}_1$ ,

$$\mathbf{Im}(\tau_k^* \varphi_a^c)(A^* \delta_\Phi(A)) \geq S\left(\tau_k^* \varphi_a^c(AA^*), \tau_k^* \varphi_a^c(A^* A)\right) - 2\varepsilon \|A\|^2.$$

For  $k \in \mathcal{C}_2$ , we use simply the following estimate.

$$\mathbf{Im}(\tau_k^* \varphi_a^c)(A^* \delta_\Phi(A)) \geq -\|A \delta_\Phi(A)\|.$$

From these inequalities, we obtain

$$\begin{aligned} & \mathbf{Im} \widehat{\varphi_a^c}(A^* \delta_\Phi(A)) \\ &= \mathbf{Im} \left( \sum_{k \in \mathcal{C}_1} \frac{1}{|\mathcal{C}_a|} \tau_k^* \varphi_a^c \right) (A^* \delta_\Phi(A)) + \mathbf{Im} \left( \sum_{k \in \mathcal{C}_2} \frac{1}{|\mathcal{C}_a|} \tau_k^* \varphi_a^c \right) (A^* \delta_\Phi(A)) \\ &\geq \frac{1}{|\mathcal{C}_a|} \sum_{k \in \mathcal{C}_1} S\left(\tau_k^* \varphi_a^c(AA^*), \tau_k^* \varphi_a^c(A^* A)\right) \\ &\quad - 2 \frac{|\mathcal{C}_1|}{|\mathcal{C}_a|} \varepsilon \|A\|^2 - \frac{|\mathcal{C}_2|}{|\mathcal{C}_a|} \|A \delta_\Phi(A)\|. \end{aligned} \tag{12.32}$$

Due to the estimate (12.27), the last term tends to 0 as  $a \rightarrow \infty$ , while the second last term tends to  $-2\varepsilon \|A\|^2$  as  $a \rightarrow \infty$ . Due to the convexity of  $S(\cdot, \cdot)$  in two variables, the first term on the right-hand side has the following lower bound:

$$\frac{1}{|\mathcal{C}_a|} \sum_{k \in \mathcal{C}_1} S\left(\tau_k^* \varphi_a^c(AA^*), \tau_k^* \varphi_a^c(A^* A)\right) \geq \frac{|\mathcal{C}_1|}{|\mathcal{C}_a|} S\left(\widehat{\varphi_a^c}'(AA^*), \widehat{\varphi_a^c}'(A^* A)\right), \tag{12.33}$$

where  $\widehat{\varphi_a^c}'$  is a state of  $\mathcal{A}$  defined by

$$\widehat{\varphi_a^c}'(B) \equiv \frac{1}{|\mathcal{C}_1|} \sum_{k \in \mathcal{C}_1} \tau_k^* \varphi_a^c(B), \quad B \in \mathcal{A}.$$

The difference of the states  $\widehat{\varphi_a^c}$  and  $\widehat{\varphi_a^c}'$  can be estimated as

$$\begin{aligned} \widehat{\varphi_n^c}' - \widehat{\varphi_n^c} &= \left( \frac{1}{|\mathcal{C}_1|} - \frac{1}{|\mathcal{C}_a|} \right) \sum_{k \in \mathcal{C}_1} \tau_k^* \varphi_a^c - \frac{1}{|\mathcal{C}_a|} \sum_{k \in \mathcal{C}_2} \tau_k^* \varphi_a^c \\ &= \frac{|\mathcal{C}_2|}{|\mathcal{C}_a|} \widehat{\varphi_n^c}' - \frac{1}{|\mathcal{C}_a|} \sum_{k \in \mathcal{C}_2} \tau_k^* \varphi_a^c. \end{aligned}$$

Hence

$$\|\widehat{\varphi_n^c}' - \widehat{\varphi_n^c}\| \leq 2 \frac{|\mathcal{C}_2|}{|\mathcal{C}_a|},$$

which tends to 0 as  $a \rightarrow \infty$  by (12.27). We note

$$\begin{aligned} \lim_a \widehat{\varphi_a^c}'(AA^*) &= \lim_a \widehat{\varphi_a^c}(AA^*) = \varphi(AA^*) \\ \lim_a \widehat{\varphi_a^c}'(A^* A) &= \lim_a \widehat{\varphi_a^c}(A^* A) = \varphi(A^* A). \end{aligned}$$

By the lower semi-continuity of  $S(\cdot, \cdot)$ , we obtain

$$\liminf_a S\left(\widehat{\varphi}_a^c(AA^*), \widehat{\varphi}_a^c(A^*A)\right) \geq S\left(\varphi(AA^*), \varphi(A^*A)\right). \quad (12.34)$$

Combining the estimates (12.32), (12.33), (12.34) as well as (12.27), we obtain the following inequality in the limit  $a \rightarrow \infty$ .

$$\mathbf{Im}\varphi(A^*\delta_\Phi(A)) \geq S\left(\varphi(AA^*), \varphi(A^*A)\right) - 2\varepsilon\|A\|^2.$$

Due to arbitrariness of  $\varepsilon$ , we have

$$\mathbf{Im}\varphi(A^*\delta_\Phi(A)) \geq S\left(\varphi(AA^*), \varphi(A^*A)\right),$$

for  $A \in \mathcal{A}_\circ$ . Hence the condition (C-2) holds.

Thus, we have shown that  $\varphi$  satisfies the  $(\delta_\Phi, 1)$ -dKMS condition if  $\varphi$  is the (unique) solution of  $(\Phi, 1)$ -variational principle when  $\Phi \in \mathcal{P}_\tau^1$ .

For general  $\Phi \in \mathcal{P}_\tau$ , we will use the standard argument of the convex analysis in the same way as [26], or Theorem 6.2.42 in [17].

By Theorem 12.8, any solution of the  $(\Phi, 1)$ -variational principle can be obtained by successive use of the following procedures, starting with the unique solution of  $\varphi_\alpha$  of  $(\Phi_\alpha, 1)$ -variational principle for  $\Phi_\alpha \in \mathcal{P}_\tau^1$ .

- (1) Weak\* limits of any converging nets  $\varphi_\alpha$  such that  $\|\Phi_\alpha - \Phi\| \rightarrow 0$ .
- (2) Convex combinations of limits obtained in (1).
- (3) Weak\* limits of a converging net of states obtained in (2).

By Lemma 6.6, the conditions (C-1) and (C-2) are stable under these procedures. As we have already shown these conditions for  $\varphi_\alpha$  when  $\Phi_\alpha$  belongs to  $\mathcal{P}_\tau^1$ , the same holds for any  $\Phi \in \mathcal{P}_\tau$ .  $\square$

We have now shown Theorem B.

## 13 Use of Other Entropy in the Variational Equality

We now consider the possibility to replace the mean entropy  $s(\omega)$  in Theorem 11.4 by other entropy. We take up the CNT entropy  $h_\omega(\tau)$  with respect to the lattice translation automorphism group  $\tau$  as one example. But readers will find that any other entropy will do if it has those basic properties of CNT entropy which are used in the proof of Theorem 13.2. Note that it is not known whether CNT entropy is equal to the mean entropy or not so far, either in some general context or in the present case.

### 13.1 CNT-Entropy

The CNT-entropy is introduced by Connes-Narnhofer-Thirring [19] for a single automorphism and its invariant state, and is extended by Hudetz [22] to the multi-dimensional case of the group  $\mathbb{Z}^\nu$  generated by a finite number ( $=\nu$ ) of commuting automorphisms. We will use the latter extended version for the group of lattice translation automorphisms  $\tau_m$  ( $m \in \mathbb{Z}^\nu$ ).



For a positive integer  $k$ , we consider a finite decomposition of a state  $\omega$  in the state space  $\mathcal{A}_{+,1}^*$ :

$$\omega = \sum_{i(1), i(2), \dots, i(k)} \omega_{i(1)i(2)\dots i(k)}, \quad (13.1)$$

where each  $i(l)$  runs over a finite subset of  $\mathbb{N}$ ,  $l = 1, \dots, k$ , and  $\omega_{i(1)i(2)\dots i(k)}$  is a nonzero positive linear functional of  $\mathcal{A}$ . For each fixed  $l$  and  $i(l)$ , let

$$\omega_{i(l)}^l \equiv \sum_{\substack{i(1), i(2), \dots, i(k) \\ i(l): \text{fixed}}} \omega_{i(1)i(2)\dots i(k)}, \quad \hat{\omega}_{i(l)}^l \equiv \frac{\omega_{i(l)}^l}{\omega_{i(l)}^l(\mathbf{1})}. \quad (13.2)$$

Let  $\eta(x) \equiv -x \log x$  for  $x > 0$  and  $\eta(0) = 0$ . For finite dimensional subalgebras  $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_k$  of  $\mathcal{A}$ , the so-called algebraic entropy  $H_\omega(\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_k)$  is defined by

$$\begin{aligned} & H_\omega(\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_k) \\ & \equiv \sup \left[ \sum_{i(1), i(2), \dots, i(k)} \eta(\omega_{i(1)i(2)\dots i(k)}(\mathbf{1})) - \sum_{l=1}^k \sum_{i(l)} \eta(\omega_{i(l)}^l(\mathbf{1})) \right. \\ & \quad \left. + \sum_{l=1}^k S(\omega|_{\mathcal{A}_l}) - \sum_{l=1}^k \sum_{i(l)} \omega_{i(l)}^l(\mathbf{1}) S(\hat{\omega}_{i(l)}^l|_{\mathcal{A}_l}) \right], \end{aligned} \quad (13.3)$$

where the supremum is taken over all finite decompositions (13.1) of  $\omega$  with a fixed  $k$ .

If  $\omega$  is  $\tau$ -invariant, the following limit (denoted by  $h_{\omega, \tau}(N)$ ) is known to exist (as the infimum over  $a$ ) for any finite dimensional subalgebra  $N \subset \mathcal{A}$ ,

$$h_{\omega, \tau}(N) \equiv \lim_{a \rightarrow \infty} \frac{1}{|\mathcal{C}_a|} H_\omega(N, \dots, \tau^k(N), \dots, \tau^{a-1}(N)),$$

where there are  $|\mathcal{C}_a|$  arguments for  $H_\omega(\dots)$  and each of them is  $\tau^k(N)$ ,  $k \in \mathcal{C}_a$ . Let  $N_1 \subset N_2 \subset \dots \subset N_n \subset \dots$  be an increasing sequence of finite algebras such that the norm closure  $\overline{\bigcup_n N_n}$  is equal to  $\mathcal{A}$ . By a Kolmogorov-Sinai type theorem (Corollary V.4 in [19]), the CNT-entropy  $h_\omega(\tau)$  is given by

$$h_\omega(\tau) = \lim_{n \rightarrow \infty} h_{\omega, \tau}(N_n). \quad (13.4)$$

### 13.2 Variational Equality in Terms of CNT-Entropy

Let  $J_1, J_2, \dots, J_k$  be disjoint finite subsets of  $\mathbb{Z}^\nu$  with their union  $J$ . From Lemma VIII.1 in [19] it follows that

$$H_\omega(\mathcal{A}(J_1), \mathcal{A}(J_2), \dots, \mathcal{A}(J_k)) \leq S(\omega_J). \quad (13.5)$$

When  $\omega$  is an even ‘product state’, the equality holds as follows (the following simple proof is due to a referee).

**Lemma 13.1.** *Let  $J_1, J_2, \dots, J_k$  be disjoint finite subsets with their union  $J$ . Let  $\omega$  be a  $\Theta$ -even state of  $\mathcal{A}$ . Assume that  $\omega$  has the following product property:*

$$\omega(A_1 A_2 \dots A_k B) = \omega(A_1) \omega(A_2) \dots \omega(A_k) \omega(B), \quad (13.6)$$

where  $A_j$  is an arbitrary element in  $\mathcal{A}(J_j)$  ( $j = 1, \dots, k$ ) and  $B$  is an arbitrary element in  $\mathcal{A}(J^c)$ . Then

$$H_\omega(\mathcal{A}(J_1), \mathcal{A}(J_2), \dots, \mathcal{A}(J_k)) = S(\omega_J) = \sum_{l=1}^k S(\omega_{J_l}), \quad (13.7)$$

and

$$H_\omega(\mathcal{A}(J_1), \mathcal{A}(J_2), \dots, \mathcal{A}(J_k)) = \sum_{l=1}^k H_\omega(\mathcal{A}(J_l)). \quad (13.8)$$

*Proof.* We define

$$E_{J_i(+)} \equiv \frac{1}{2}(\text{id} + \Theta^{J_i}).$$

Then  $E_{J_1, \dots, J_k(+)} \equiv E_{J_1(+)} \cdots E_{J_k(+)}$  is the conditional expectation from  $\mathcal{A}$  onto  $\mathcal{A}(J_1)_+ \otimes \cdots \otimes \mathcal{A}(J_k)_+ \otimes \mathcal{A}(J^c)$ . Since  $\omega$  is a product state for the tensor product  $(\mathcal{A}(J_1)_+ \otimes \cdots \otimes \mathcal{A}(J_k)_+) \otimes \mathcal{A}(J^c)$ , there exists an  $\omega$ -preserving conditional expectation  $E'_\omega$  from  $(\mathcal{A}(J_1)_+ \otimes \cdots \otimes \mathcal{A}(J_k)_+) \otimes \mathcal{A}(J^c)$  onto  $\mathcal{A}(J_1)_+ \otimes \cdots \otimes \mathcal{A}(J_k)_+$ . Hence

$$E_{J_1, \dots, J_k(+)}^\omega \equiv E'_\omega E_{J_1, \dots, J_k(+)}$$

is an  $\omega$ -preserving conditional expectation from  $\mathcal{A}$  onto  $\mathcal{A}(J_1)_+ \otimes \cdots \otimes \mathcal{A}(J_k)_+$ . Hence

$$\begin{aligned} & H_\omega(\mathcal{A}(J_1)_+, \mathcal{A}(J_2)_+, \dots, \mathcal{A}(J_k)_+) \\ &= H_{\omega|_{\mathcal{A}(J_1)_+ \otimes \cdots \otimes \mathcal{A}(J_k)_+}}(\mathcal{A}(J_1)_+, \mathcal{A}(J_2)_+, \dots, \mathcal{A}(J_k)_+) \\ &= \sum_{l=1}^k S(\omega|_{\mathcal{A}(J_l)_+}) = S(\omega_J). \end{aligned}$$

On the other hand,

$$\begin{aligned} H_\omega(\mathcal{A}(J_1)_+, \mathcal{A}(J_2)_+, \dots, \mathcal{A}(J_k)_+) &\leq H_\omega(\mathcal{A}(J_1), \mathcal{A}(J_2), \dots, \mathcal{A}(J_k)) \\ &\leq S(\omega_J). \end{aligned}$$

□

We are now in a position to give the main theorem of this subsection.

**Theorem 13.2.** *Assume the same conditions on  $\Phi$  as Theorem 11.4. Then*

$$P(\beta\Phi) = \sup_{\omega \in \mathcal{A}_{+,1}^{\ast,\tau}} [h_\omega(\tau) - \beta e_\Phi(\omega)], \quad (13.9)$$

where  $h_\omega(\tau)$  is the CNT-entropy of  $\omega$  with respect to the lattice translation  $\tau$ .

*Proof.* Based on Lemma 13.1, the proof will go in the same as the case of quantum lattice systems [32]. Basic properties of the CNT-entropy to which we use in the proof are as follows.

(i) Covariance under an automorphism of  $\mathcal{A}$  (the adjoint action on states and conjugacy action on the shift).

- (ii) Scaling property under the scaling of the automorphism group.
- (iii) Concave dependence on states.

Due to (13.5), we have

$$h_\omega(\tau) \leq s(\omega), \quad (13.10)$$

for any translation invariant state  $\omega$ . Hence the variational inequality (11.10) obviously holds when  $s(\omega)$  is replaced by  $h_\omega(\tau)$ .

Due to Lemma 13.1 and the product property of  $\varphi_a^c$ , the translation invariant state  $\widehat{\varphi_a^c}$  defined in (11.23) will play an identical role as in the proof of Theorem 11.4. Therefore the sequence

$$\{h_{\widehat{\varphi_a^c}}(\tau) - e_\Phi(\widehat{\varphi_a^c})\}$$

tends to the supremum value  $P(\Phi)$  of the variational inequality as  $a \rightarrow \infty$ . Hence the theorem follows.  $\square$

*Remark.* (iii) is a general property of CNT-entropy (see e.g. [41]) and is enough for the proof. But in the situation of the above proof, the affinity holds due to the specific nature of the states to be considered.

The preceding result is the variational equality. We are then interested in the variational principle.

**Proposition 13.3.** *Suppose that a translation invariant state  $\varphi$  satisfies*

$$P(\beta\Phi) = h_\varphi(\tau) - \beta e_\Phi(\varphi). \quad (13.11)$$

*Then  $\varphi$  is a solution of the  $(\Phi, \beta)$ -variational principle and*

$$h_\varphi(\tau) = s(\varphi). \quad (13.12)$$

*Proof.* By (13.5), we have

$$s(\varphi) - \beta e_\Phi(\varphi) \geq h_\varphi(\tau) - \beta e_\Phi(\varphi) = P(\beta\Phi).$$

By the variational inequality (11.10), we have

$$s(\varphi) - \beta e_\Phi(\varphi) = P(\beta\Phi). \quad (13.13)$$

Therefore  $\varphi$  is a solution of the  $(\Phi, \beta)$ -variational principle. From (13.11) and (13.13), we obtain (13.12).

*Remark 1.* We have no result about the existence theorem for a solution of the variational principle (13.11) in terms of the CNT-entropy for a general  $\Phi \in \mathcal{P}_\tau$  (like Theorem 11.7) nor the stability of solutions of such a variational principle (like Lemma 12.9), the obstacle in applying the usual method being absence of any result about weak\* upper semicontinuity of  $h_\omega(\tau)$  in  $\omega$ .

In this sense, Proposition 13.3 is a superficial result, and Theorem 11.4 is short of ‘the variational principle’ in terms of the CNT-entropy. See also the discussion in Section 4 of [32].

*Remark 2.* Although we have used CNT entropy throughout this section, other entropy such as  $ht_\omega(\sigma)$  defined by Choda [18] can be substituted into  $h_\omega(\tau)$ , yielding similar results.

## 14 Discussion

The following are some of remaining problems about equilibrium statistical mechanics of Fermion lattice systems which are not covered in this paper.

### 1. Dynamics which does not commute with $\Theta$

Obviously, there is an inner one-parameter group of  $*$ -automorphisms which does not commute with  $\Theta$ . Examples of outer dynamics not commuting with  $\Theta$  can be constructed in the following way (suggested by one of referees). Let  $\{I_i\}_{i=1,2,\dots}$  be a partition of the lattice  $\mathbb{Z}^\nu$  into mutually disjoint finite subsets  $I_i$  and let  $J_j \equiv \cup_{i \leq j} I_i$ . Choose a self-adjoint  $b_i$  in  $\mathcal{A}(I_i)_-$  for each  $i$  and set  $\Phi(J_i) \equiv v_{J_{i-1}} b_i$  where  $v_J$  is given by (4.30). By Theorem 4.17(1), they mutually commute and  $\Phi(J_i) \in \mathcal{A}(J_{i-1})'$  for each  $i$ . Hence  $\alpha_t^{(i)} \equiv \text{Ade}^{it\Phi(J_i)}$ ,  $i = 1, 2, \dots$ , are mutually commuting dynamics of  $\mathcal{A}$ ,  $\alpha_t^{(i)}$  leaving elements of  $\mathcal{A}(J_{i-1})$  invariant. Hence  $\alpha_t \equiv \prod_{i=1}^{\infty} \alpha_t^{(i)}$  gives a dynamics of  $\mathcal{A}$  satisfying  $\Theta\alpha_t = \alpha_{-t}\Theta$ . (Namely, its generator anticommutes with  $\Theta$ .) The corresponding potential is given by  $\Phi(I) = 0$  if  $I \neq J_i$  for any  $i$  and  $\Phi(I) = \Phi(J_i)$  if  $I = J_i$ . This potential satisfies the standardness condition ( $\Phi$ -d) if each  $b_i$  satisfies it for the set  $I_i$ . By looking at the behavior of  $U_{n,N}^* \alpha_t(U_{n,N}) = e^{-2it\sum_{i=0}^N \Phi(J_{n+i})}$  for  $U_{n,N} \equiv \prod_{i=0}^N v_{I_{n+i}}$  as  $n \rightarrow \infty$ , the dynamics is seen to be outer unless  $\sum_i \Phi(J_i)$  is convergent.

### 2. Broken $\Theta$ -invariance of equilibrium states

In connection with the Gibbs condition, we have shown in §7.7 that the perturbed state either by surface energy or by the local interaction energy satisfies the product property if and only if the equilibrium state is  $\Theta$ -invariant. However, we do not know an example of an equilibrium state which is not  $\Theta$ -invariant. Existence or non-existence of such a state seems to be an important question. It seems to be closely related to the next problem 3.

Note that any translation invariant state is  $\Theta$ -invariant. So we need broken translation invariance of an equilibrium state for its broken  $\Theta$ -invariance.

### 3. Local Thermodynamical Stability (LTS)

In parallel with the case of quantum spin lattice system, one can formulate the local stability condition ([10], [39]) for our Fermion lattice system. However, there seems to be two choices of the outside system for a local algebra  $\mathcal{A}(I)$  ( $I$  finite). (1) The commutant  $\mathcal{A}(I)'$ . (2)  $\mathcal{A}(I^c)$ . For the choice (1), all arguments in the case of quantum spin lattice systems seem to go through for the Fermion lattice system leading to equivalence of LTS with the KMS condition under our basic Assumptions (I), (II) and (III).

On the other hand, (2) seems to be physically correct choice, although we do not have an equivalence proof for (2) so far.

In this connection, the problem 2 is crucial. If all equilibrium state is  $\Theta$ -invariant, then the choice (2) also seems to give the LTS which is equivalent to the KMS under our basic assumptions. A paper on this problem is forthcoming [15].

### 4. Downstairs Equivalence

We may say that the dynamics  $\alpha_t$  is working upstairs while its generator is working downstairs. In particular, our arena for the downstairs activity is

$\mathcal{A}_\circ$ . The stair going upstairs seems to be not wide open. On the other hand, there seems to be a lot more room downstairs. There, we have established the one-to-one correspondence between ( $\Theta$ -invariant) derivations on  $\mathcal{A}_\circ$  and standard potentials. We have shown that the solution of the variational principle (described in terms of a translation covariant potential) satisfies the dKMS condition on  $\mathcal{A}_\circ$  (described in terms of the corresponding derivation). How about the converse.

There is also the problem of equivalence of LTS condition (in terms of a potential) and the dKMS condition on  $\mathcal{A}_\circ$  (in terms of the corresponding derivation) where the translation invariance is not needed. Some aspects of this problem will also be included in the forthcoming paper [15].

## 5. Equivalent Potentials

We have introduced the notion of general potentials and equivalence among them in § 5.5. Our theory is developed only for the unique standard potential among each equivalence class. Natural questions about general potentials arise.

Does the existence of the limits defining the pressure  $P(\beta\Phi)$  and the mean energy  $e_\Phi(\varphi)$  hold also for translation covariant general potentials  $\Phi$ ? Assuming the existence, are the  $P(\beta\Phi)$  and  $e_\Phi(\varphi)$  the same as those for the unique standard potential  $\Phi_s$  equivalent to  $\Phi$ ? If they are different, how about the solution of their variational principle?

We give a partial answer to these questions.

**Proposition 14.1.** *Let  $\Phi$  be a translation covariant potential (which satisfies ( $\Phi$ -a,b,c,e,f) by definition) fulfilling the following additional condition: the surface energy*

$$W_\Phi(I) = \lim_{J \nearrow \mathbb{Z}^\nu} \sum_K \{\Phi(K); K \cap I \neq \emptyset, K \cap I^c \neq \emptyset, K \subset J\}. \quad (14.1)$$

*satisfies*

$$\text{v.H.} \lim_{I \rightarrow \infty} \frac{\|W_\Phi(I)\|}{|I|} = 0. \quad (14.2)$$

*Let  $\Phi_s$  be the standard potential (in  $\mathcal{P}_\tau$ ) which is equivalent to  $\Phi$ . Then both van Hove limits defining  $P(\beta\Phi)$  and  $e_\Phi(\omega)$  for all  $\omega \in \mathcal{A}_{+,1}^{\tau,*}$  exist if and only if*

$$C_\Phi \equiv \text{v.H.} \lim_{I \rightarrow \infty} \frac{\tau(H_\Phi(I))}{|I|} \quad (14.3)$$

*exists.*

*If this is the case, then the following relations hold*

$$\begin{aligned} P(\beta\Phi) &= \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \log \text{Tr}_I(e^{-\beta H(I)}) \\ &= \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \log \text{Tr}_I(e^{-\beta U(I)}) \\ &= P(\beta\Phi_s) - \beta C_\Phi, \end{aligned} \quad (14.4)$$

$$\begin{aligned}
e_{\Phi}(\omega) &= \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \omega(H(I)) \\
&= \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \omega(U(I)) \\
&= e_{\Phi_s}(\omega) + C_{\Phi}.
\end{aligned} \tag{14.5}$$

Furthermore,  $(\Phi, \beta)$ - and  $(\Phi_s, \beta)$ - variational principle give the same set of solutions.

*Remark.* If  $\tau(\Phi(I)) = 0$  for all  $I$ , then (14.3) exists and  $C_{\Phi} = 0$ . Hence  $P(\beta\Phi) = P(\beta\Phi_s)$  and  $e_{\Phi}(\omega) = e_{\Phi_s}(\omega)$ . This can be achieved for any general potential  $\Phi$  by changing it to  $\Phi_1 = \Phi - \Phi_0$  where  $\Phi_0$  is a scalar-valued potential given by

$$\Phi_0(I) = \tau(\Phi(I))\mathbf{1}.$$

*Proof.* Since  $\Phi$  and  $\Phi_s$  are equivalent, we have

$$H_{\Phi}(I) - H_{\Phi_s}(I) \in \mathcal{A}(I)'. \tag{14.6}$$

Since  $H_{\Phi}(I) - H_{\Phi_s}(I)$  is  $\Theta$ -even by  $(\Phi\text{-c})$  for  $\Phi$  and  $\Phi_s$ , we have

$$H_{\Phi}(I) - H_{\Phi_s}(I) \in \mathcal{A}(I^c)_+. \tag{14.6}$$

Hence,

$$\begin{aligned}
U_{\Phi}(I) - U_{\Phi_s}(I) &= E_I(U_{\Phi}(I) - U_{\Phi_s}(I)) \\
&= E_I(H_{\Phi}(I) - H_{\Phi_s}(I)) - E_I(W_{\Phi}(I) - W_{\Phi_s}(I)) \\
&= \tau(H_{\Phi}(I) - H_{\Phi_s}(I)) - E_I(W_{\Phi}(I) - W_{\Phi_s}(I)),
\end{aligned}$$

due to (14.6). By  $\tau(H_{\Phi_s}(I)) = 0$  and  $E_I(W_{\Phi_s}(I)) = 0$  due to  $(\Phi\text{-d})$ , we have

$$U_{\Phi}(I) - U_{\Phi_s}(I) = \tau(H_{\Phi}(I)) - E_I(W_{\Phi}(I)).$$

By (14.2), we have

$$\text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \|U_{\Phi}(I) - U_{\Phi_s}(I) - \tau(H_{\Phi}(I))\| = 0.$$

Also by (14.2),

$$\text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \|H_{\Phi}(I) - U_{\Phi}(I)\| = 0.$$

Hence (14.5) follows:

$$\begin{aligned}
\text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \omega(H_{\Phi}(I)) &= \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \omega(U_{\Phi}(I)) \\
&= \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \omega(U_{\Phi_s}(I)) + \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \tau(H_{\Phi}(I)) \\
&= e_{\Phi_s} + \text{v.H.} \lim_{I \rightarrow \infty} \frac{1}{|I|} \tau(H_{\Phi}(I)).
\end{aligned}$$

We also have

$$\begin{aligned} \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{1}{|\mathbf{I}|} \log \mathbf{Tr}_{\mathbf{I}}(e^{-H(\mathbf{I})}) &= \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{1}{|\mathbf{I}|} \log \mathbf{Tr}_{\mathbf{I}}(e^{-U(\mathbf{I})}) \\ &= P(\beta\Phi_s) - \beta \left\{ \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{1}{|\mathbf{I}|} \tau(H_{\Phi}(\mathbf{I})) \right\}, \end{aligned}$$

which shows (14.4).  $\square$

*Remark.* Suppose that  $\Phi$  satisfies  $(\Phi\text{-a})$ ,  $(\Phi\text{-b})$ ,  $(\Phi\text{-c})$ ,  $(\Phi\text{-f})$  and

$$\sum_{\mathbf{I} \ni 0} \|\Phi(\mathbf{I})\| < \infty. \quad (14.7)$$

Then it satisfies  $(\Phi\text{-e})$  automatically and is a general potential. Furthermore, (14.2) is known to be satisfied (the same proof as Lemma 9.1 holds except for estimates (9.2) (9.3), (9.4) and (9.5) which follow from the absolute convergence of (14.7) due to (7.12)) and

$$C_{\Phi} = \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{\tau(H_{\Phi}(\mathbf{I}))}{|\mathbf{I}|} = \text{v.H.} \lim_{\mathbf{I} \rightarrow \infty} \frac{\tau(U_{\Phi}(\mathbf{I}))}{|\mathbf{I}|} = e_{\Phi}(\tau) \quad (14.8)$$

is known to converge. (The same proof as Theorem 9.5 holds except for a modification of proof of some estimates for Lemma 9.2 on the basis of the absolute convergence of (14.7). See also e.g. Proposition 6.2.39 of [17].)

Therefore (14.4) and (14.5) hold and the solutions of  $(\Phi, \beta)$ - and  $(\Phi_s, \beta)$ -variational principle coincide.

## References

- [1] H.Araki and E.H.Lieb, Entropy inequalities, *Commun. Math. Phys.* **18**(1970), 160-170.
- [2] H.Araki, Relative hamiltonian for faithful normal states of a von Neumann algebra, *Publ. RIMS, Kyoto Univ.* **7**(1973), 165-209.
- [3] H.Araki, Expansional in Banach algebra, *Ann. Sci. École Norm Sup. Sér.* **46**(1973), 67-84.
- [4] H.Araki, Golden-Thompson and Peierls-Bogoliubov inequalities for a general von Neumann algebra, *Comm. Math. Phys.* **34**(1973), 167-178.
- [5] H.Araki and P.D.F.Ion, On the equivalence of KMS and Gibbs conditions for states of quantum lattice systems, *Commun. Math. Phys.* **35**(1974), 1-12.
- [6] H.Araki, On the equivalence of the KMS condition and the variational principle for quantum lattice systems, *Commun. Math. Phys.* **38**(1974), 1-10.
- [7] H.Araki, Relative entropy and its application, in Colloques Internationaux du C.N.R.S. No.248 *Les Methodes Mathematiques de la Theorie Quantique des Champs*, eds. F.Guerra, D.W.Robinson, and R.Stora, (CNRS, Paris, 1976), 61-79.

- [8] H.Araki, Relative entropy of states of von Neumann algebras, *Publ. RIMS, Kyoto Univ.* **11**(1976), 809-833.
- [9] H.Araki, Relative entropy of states of von Neumann algebras II, *Publ. RIMS, Kyoto Univ.* **13**(1977), 173-192.
- [10] H.Araki and G.L.Sewell, KMS conditions and local thermodynamical stability of quantum lattice systems, *Commun. Math. Phys.* **52**(1977), 103-109.
- [11] H.Araki, D.Kastler, M.Takesaki and R.Haag, Extension of KMS states and chemical potentials, *Commun. Math. Phys.* **53**(1977), 97-134.
- [12] H.Araki, On KMS states of a  $C^*$ -dynamical system, Lecture Notes in Math.**650**, (Springer-Verlag, 1978), 66-84.
- [13] H.Araki, *Toukeirikigaku no suuri*, Iwanami(Japanese), 1994.
- [14] H.Araki and H.Moriya, Joint extension of states of subsystems for a CAR system, to appear in *Commun. Math. Phys.*
- [15] H.Araki and H.Moriya, Local thermodynamical stability of Fermion lattice systems, *Lett. Math. Phys.* **62**(2002), 33-45.
- [16] B.Baumgartner, A partial ordering of sets, making mean entropy monotone, *J. Phys. A: Math. Gen.* **35** (2002), 3163-3182.
- [17] O.Bratteli and D.W.Robinson, *Operator Algebras and Quantum Statistical Mechanics 2, 2nd edition*, Springer-Verlag, 1996.
- [18] M.Choda, A  $C^*$ -Dynamical Entropy and Applications to Canonical Endomorphisms *J. Funct. Anal.* **173**(2000), 453-480.
- [19] A.Connes, H.Narnhofer and W.Thirring, Dynamical Entropy of  $C^*$ -Algebras and von Neumann Algebras, *Commun. Math. Phys.* **112**(1987), 691-719.
- [20] M.Fannes, A continuity property of the entropy density for spin lattice systems, *Commun. Math. Phys.* **31**(1973), 291-294.
- [21] F.M.Goodman, P. de la Harpe, and V.F.R. Jones, *Coxeter Graphs and Towers of Algebras*, Springer-Verlag, 1989.
- [22] T.Hudetz, Spacetime Dynamical Entropy of Quantum Systems, *Lett. Math. Phys.* **16**(1988), 151-161.
- [23] R.B.Israel, *Convexity in the Theory of Lattice Gases*, Princeton University Press, 1979.
- [24] A.R.Kay and B.S.Kay, Monotonicity with volume of entropy and of mean entropy for translationally invariant systems as consequences of strong subadditivity, *J.Phys.A. Math.Gen.* **34**(2001)365-382.
- [25] H.Kosaki, Relative entropy for states: a variational expressions, *J.Operator.Theory***16**(1986), 335-348.



- [26] O.E. Lanford III and D.W. Robinson, Statistical mechanics of quantum spin systems III, *Commun. Math. Phys.* **9**(1968), 327-338.
- [27] E.H.Lieb and M.B.Ruskai, Proof of the strong subadditivity of quantum-mechanical entropy, *J. Math. Phys.* **14**(1973), 1938-1941.
- [28] E.H.Lieb and M.B.Ruskai, A fundamental property of quantum-mechanical entropy, *Phys.Rev.Lett.* **30**(1973), 434-436.
- [29] T.Matsui, Ground states of fermions on lattices, *Commun. Math. Phys.* **182**(1996), 723-751.
- [30] T.Matsui, Quantum statistical mechanics and Feller semigroup, *Quantum Probability Communication* **10**(1998), 101-124.
- [31] S.Mazur, Über konvexe Menge in linearen normierten Raumen, *Studia. Math.* **4**(1933), 70-84.
- [32] H.Moriya, Variational principle and the dynamical entropy of space translation, *Rev. Math. Phys.* **11**(1999), 1315-1328.
- [33] H.Moriya, Some aspects of quantum entanglement for CAR systems, *Lett. Math. Phys.* **60**(2002), 109-121.
- [34] S.Neshveyev and E.Størmer, The variational principle for a class of asymptotically abelian  $C^*$ -algebras, *Commun. Math. Phys.* **215**(2000), 177-196.
- [35] D.Petz, On certain properties of the relative entropy of states of operator algebras, *Math. Z.* **206**(1991), 351-361.
- [36] R.T.Powers, Representations of the canonical anticommutation relations, Thesis, Princeton University(1967).
- [37] D.Ruelle, A variational formulation of equilibrium statistical mechanics and the Gibbs phase rule, *Commun. Math. Phys.* **5**(1967), 324-329.
- [38] S.Sakai, On one-parameter subgroups of  $*$ -automorphisms on operator algebras and the corresponding unbounded derivations, *Amer. J. Math.* **98**(1976), 427-440.
- [39] G.L.Sewell, KMS conditions and local thermodynamical stability of quantum lattice systems II, *Commun. Math. Phys.* **55**(1977), 53-61.
- [40] B.Simon, *The Statistical Mechanics of Lattice Gases*, Princeton University Press, 1993.
- [41] E.Størmer, A survey of noncommutative dynamical entropy, *Oslo preprint, Dep. of Mathematics* **18**(2000).
- [42] M.Takesaki, Tomita's Theory of Modular Hilbert-Algebras and its Application, Lecture Notes in Math.**128**, Springer-Verlag(1970).
- [43] M.Takesaki, *Theory of Operator Algebras I*, Springer-Verlag, 1979.
- [44] J.Tomiyama, On the projection of norm one in  $W^*$ -algebras, *Proc. Japan. Acad.* **33**(1957), 609-612.

- [45] H.Umegaki, Conditional expectation in an operator algebra IV, (entropy and information), *Kodai. Math. Sem. Rep.* **14**(1962), 59-85.

## A Appendix: Van Hove Limit

For the sake of mathematical precision, we present some digression about Van Hove limit.

### A.1 Van Hove Net

We introduce mutually equivalent two types of conditions for the van Hove limit. First we start with our notation about the shapes of regions of  $\mathbb{Z}^\nu$ , which will be used hereafter. Recall that  $\mathcal{C}_a$  is a cube of size  $a$  given by (8.8). For a finite subset  $I$  of  $\mathbb{Z}^\nu$  and  $a \in \mathbb{N}$ , let  $\mathbf{n}_a^+(I)$  be the smallest number of translates of  $\mathcal{C}_a$  whose union covers  $I$ , while  $\mathbf{n}_a^-(I)$  be the largest number of mutually disjoint translates of  $\mathcal{C}_a$  that can be packed in  $I$ .

Let  $B_r(n)$  be a closed ball in  $\mathbb{R}^\nu(\supset \mathbb{Z}^\nu)$  with the center  $n \in \mathbb{Z}^\nu$  and the radius  $r \in \mathbb{R}$ . Denote the surface of  $I$  with a thickness  $r(> 0)$  by

$$\text{surf}_r(I) \equiv \left\{ n \in I; B_r(\{n\}) \cap I^c \neq \emptyset \right\}. \quad (\text{A.1})$$

In what follows, we consider a net of finite subsets  $I_\alpha$  of  $\mathbb{Z}^\nu$  where the set of indices  $\alpha$  is a directed set. Its partial ordering need not have any relation with the set inclusion partial ordering of  $I_\alpha$ .

**Lemma A.1.** *For a net of finite subsets  $I_\alpha$  of  $\mathbb{Z}^\nu$ , the following two conditions are equivalent :*

(1) *For any  $a \in \mathbb{N}$ ,*

$$\lim_{\alpha} \frac{\mathbf{n}_a^-(I_\alpha)}{\mathbf{n}_a^+(I_\alpha)} = 1. \quad (\text{A.2})$$

(2) *For any  $r > 0$ ,*

$$\lim_{\alpha} \frac{1}{|I_\alpha|} \left| \text{surf}_r(I_\alpha) \right| = 0 \quad (\text{A.3})$$

*Proof.* (1)  $\rightarrow$  (2):

Let  $\varepsilon > 0$  and  $r > 0$  be given. Let  $a \in \mathbb{N}$  be sufficiently large so that  $a \geq 2r + 1$  and

$$\varepsilon_1 \equiv 1 - \left( \frac{[a - 2r]^\nu}{a^\nu} \right) < \frac{\varepsilon}{2},$$

where  $[b]$  indicates the maximal integer not exceeding  $b$ .

By the condition (1), there exists an index  $\alpha_0$  of the net  $\{I_\alpha\}$  such that, for  $\alpha \geq \alpha_0$ ,

$$\varepsilon_2 \equiv 1 - \frac{\mathbf{n}_a^-(I_\alpha)}{\mathbf{n}_a^+(I_\alpha)} < \frac{\varepsilon}{2}.$$

Let  $D_1, \dots, D_N$ , with  $N = \mathbf{n}_a^-(I_\alpha)$ , be mutually disjoint translates of  $\mathcal{C}_a$  contained in  $I_\alpha$ .

Let  $D'_i$  be a translate of  $\mathcal{C}_{[a-2r]}$  placed in  $D_i$  with a distance larger than  $r$  from the complement of  $D_i$  in  $\mathbb{Z}^\nu$  for each  $i = 1, \dots, N$  which exists. Then

$$\frac{|D'_i|}{|D_i|} = \left( \frac{[a - 2r]^\nu}{a^\nu} \right) = 1 - \varepsilon_1.$$

Let  $D$  be the union of  $D_1, \dots, D_N$  and  $D'$  be the union of  $D'_1, \dots, D'_N$ . Then

$$\frac{|D \setminus D'|}{|D|} = 1 - \frac{|D'|}{|D|} = 1 - (1 - \varepsilon_1) = \varepsilon_1.$$

Since  $\mathbf{n}_a^+(\mathbf{I}_\alpha)$  translates of  $\mathcal{C}_a$  covers  $\mathbf{I}_\alpha$ , we have

$$|\mathbf{I}_\alpha| \leq \mathbf{n}_a^+(\mathbf{I}_\alpha)|\mathcal{C}_a| = \mathbf{n}_a^+(\mathbf{I}_\alpha)a^\nu.$$

Hence

$$\frac{|\mathbf{I}_\alpha \setminus D|}{|\mathbf{I}_\alpha|} = 1 - \frac{|D|}{|\mathbf{I}_\alpha|} \leq 1 - \frac{|D|}{\mathbf{n}_a^+(\mathbf{I}_\alpha)a^\nu} = 1 - \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)}{\mathbf{n}_a^+(\mathbf{I}_\alpha)} = \varepsilon_2.$$

Due to  $\mathbf{I}_\alpha \supset D$ ,

$$\frac{|D \setminus D'|}{|\mathbf{I}_\alpha|} \leq \frac{|D \setminus D'|}{|D|} = \varepsilon_1.$$

By construction, the distance between  $D'_i$  and the complement of  $D_i$  (in  $\mathbb{Z}^\nu$ ) is larger than  $r$ , and hence the distance between  $D'_i$  and the complement of  $\mathbf{I}_\alpha$  is larger than  $r$ . Thus,

$$\text{surf}_r(\mathbf{I}_\alpha) \subset \mathbf{I}_\alpha \setminus D' = (D \setminus D') \cup (\mathbf{I}_\alpha \setminus D).$$

For  $\alpha \geq \alpha_0$ , we obtain

$$\frac{|\text{surf}_r(\mathbf{I}_\alpha)|}{|\mathbf{I}_\alpha|} \leq \varepsilon_1 + \varepsilon_2 < \varepsilon.$$

Now (1)  $\rightarrow$  (2) is proved.

(2)  $\rightarrow$  (1):

Let  $\varepsilon > 0$  and  $a \in \mathbb{N}$  be given. Take  $r > \sqrt{\nu}a$ . Let  $\alpha_0$  be an index of the net  $\mathbf{I}_a$  such that, for  $\alpha \geq \alpha_0$ ,

$$\frac{|\text{surf}_r(\mathbf{I}_\alpha)|}{|\mathbf{I}_\alpha|} < a^{-\nu}\varepsilon.$$

The translates  $\mathcal{C}_a + an$  of  $\mathcal{C}_a$  are disjoint for distinct  $n \in \mathbb{Z}^\nu$  and their union over  $n \in \mathbb{Z}^\nu$  is  $\mathbb{Z}^\nu$ . Let  $O_\alpha$  be the union of all those  $\mathcal{C}_a + an$  contained in  $\mathbf{I}_\alpha$  and  $N_1$  be their number. Let  $O'_\alpha$  be the union of all those  $\mathcal{C}_a + an$  which have non-empty intersections with both  $\mathbf{I}_\alpha$  and  $(\mathbf{I}_\alpha)^c$ , and  $N_2$  be their number. From the construction, the following estimates follow

$$N_1 \leq \mathbf{n}_a^-(\mathbf{I}_\alpha) \leq \mathbf{n}_a^+(\mathbf{I}_\alpha) \leq N_1 + N_2.$$

Furthermore, since  $\mathcal{C}_a + an$  in  $O'_\alpha$  contains a point in  $\mathbf{I}_\alpha$  as well as a point in  $(\mathbf{I}_\alpha)^c$ , and the distance of any two points in it is at most  $\sqrt{\nu}a < r$ , it has a non-empty intersection with  $\mathbf{I}_\alpha$ , which is contained in  $\text{surf}_r(\mathbf{I}_\alpha)$ . Therefore,

$$\begin{aligned} |\text{surf}_r(\mathbf{I}_\alpha)| &\geq N_2 = (N_1 + N_2) - N_1 \\ &\geq \mathbf{n}_a^+(\mathbf{I}_\alpha) - \mathbf{n}_a^-(\mathbf{I}_\alpha). \end{aligned}$$

We have also

$$|\mathbf{I}_\alpha| \leq \mathbf{n}_a^+(\mathbf{I}_\alpha) |\mathcal{C}_a| = \mathbf{n}_a^+(\mathbf{I}_\alpha) a^\nu.$$

Combining above estimates, we obtain for  $\alpha \geq \alpha_0$

$$\begin{aligned} 0 &\leq 1 - \frac{\mathbf{n}_a^-(\mathbf{I}_\alpha)}{\mathbf{n}_a^+(\mathbf{I}_\alpha)} = \frac{\mathbf{n}_a^+(\mathbf{I}_\alpha) - \mathbf{n}_a^-(\mathbf{I}_\alpha)}{\mathbf{n}_a^+(\mathbf{I}_\alpha)} \\ &\leq \frac{|\text{surf}_r(\mathbf{I}_\alpha)| a^\nu}{|\mathbf{I}_\alpha|} \\ &< \varepsilon. \end{aligned}$$

Hence, (2)  $\rightarrow$  (1) is now proved.  $\square$

**Definition A.2.** *If a net of finite subsets  $\{\mathbf{I}_\alpha\}$  satisfies the above condition (1) (or equivalently (2)), then it is said to be a van Hove net (in  $\mathbb{Z}^\nu$ ).*

We introduce the third condition on a net of finite subsets  $\mathbf{I}_\alpha$  of  $\mathbb{Z}^\nu$ :

(3) For any finite subset  $\mathbf{I}$  of  $\mathbb{Z}^\nu$ , there exists an index  $\alpha_o$  such that  $\mathbf{I}_\alpha \supset \mathbf{I}$  for all  $\alpha \geq \alpha_o$ .

**Definition A.3.** *If a net  $\{\mathbf{I}_\alpha\}$  (in  $\mathbb{Z}^\nu$ ) satisfies the conditions (1) (or equivalently (2)) and (3), then it is said to be a van Hove net tending to  $\mathbb{Z}^\nu$ .*

*Remark.* The condition (1) (or equivalently (2)) does not imply the condition (3).  $\{\mathcal{C}_n\}_{n \in \mathbb{N}}$  of (8.8) is obviously a van Hove sequence. But it does not cover the whole  $\mathbb{Z}^\nu$ . Hence it is not a van Hove sequence tending to  $\mathbb{Z}^\nu$ .

**Lemma A.4.** *For any van Hove net and for any van Hove net tending to  $\mathbb{Z}^\nu$ , the directed set can not have a maximal element.*

*Proof.* Let  $\{\mathbf{I}_\alpha\}_{\alpha \in A}$  be a van Hove net where  $A$  is a directed set of indices. We show that for any  $\alpha_o \in A$ , there exists  $\alpha' \in A$  satisfying  $\alpha' \geq \alpha_o$ ,  $\alpha' \neq \alpha_o$ .

In fact, for a given  $\alpha_o$ , there exist  $a(\alpha_o) \in \mathbb{N}$  and  $n \in \mathbb{Z}^\nu$  such that

$$\mathbf{I}_{\alpha_o} \subset \mathcal{C}_{a(\alpha_o)-n},$$

and hence

$$\mathbf{n}_{a(\alpha_o)}^-(\mathbf{I}_{\alpha_o}) = 0.$$

On the other hand, for the above  $a(\alpha_o) \in \mathbb{N}$  there exists  $\alpha_1$  such that

$$1 - \frac{\mathbf{n}_{a(\alpha_o)}^-(\mathbf{I}_\alpha)}{\mathbf{n}_{a(\alpha_o)}^+(\mathbf{I}_\alpha)} < \frac{1}{2}$$

for all  $\alpha \geq \alpha_1$ , since  $\{\mathbf{I}_\alpha\}_{\alpha \in A}$  is a van Hove net.

For any  $\alpha' \in A$  satisfying both  $\alpha' \geq \alpha_1$  and  $\alpha' \geq \alpha_o$ , we have  $\mathbf{n}_{a(\alpha_o)}^-(\mathbf{I}_{\alpha'}) \neq 0$  due to  $\alpha' \geq \alpha_1$ , and hence  $\alpha' \neq \alpha_o$ . We have shown the existence of a desired  $\alpha'$ .

A van Hove net tending to  $\mathbb{Z}^\nu$  is a special case of a van Hove net. Hence the assertion for this case obviously follows.  $\square$

## A.2 Van Hove Limit

Let  $f(I)$  be an  $\mathbb{R}$ -valued function of finite subsets  $I$  of  $\mathbb{Z}^\nu$ . We first show the following lemma which asserts the independence of the limit on the choice of van Hove net (van Hove net tending to  $\mathbb{Z}^\nu$ ) when  $f(I_\alpha)$  has a limit for any van Hove net (van Hove net tending to  $\mathbb{Z}^\nu$ )  $\{I_\alpha\}$ .

**Lemma A.5.** *If  $f(I_\alpha)$  has a limit for any van Hove net  $\{I_\alpha\}$ , then its limit is independent of such a net.*

*If  $f(I_\alpha)$  has a limit for any van Hove net  $\{I_\alpha\}$  tending to  $\mathbb{Z}^\nu$ , then its limit is independent of such a net.*

*Proof.* Let  $\{I_\alpha^1\}_{\alpha \in A}$  and  $\{I_\beta^2\}_{\beta \in B}$  be two van Hove nets where  $A$  and  $B$  are directed sets of indices. We introduce a new index set

$$C \equiv \{(\alpha, \beta, i); \alpha \in A, \beta \in B, i = 1, 2\}$$

with the partial ordering

$$(\alpha, \beta, i) \geq (\alpha', \beta', i')$$

either if  $\alpha > \alpha'$  and  $\beta > \beta'$  or if  $\alpha = \alpha'$ ,  $\beta = \beta'$  and  $i \geq i'$ .

For any  $(\alpha_1, \beta_1, i_1) \in C$  and  $(\alpha_2, \beta_2, i_2) \in C$ , there exist  $\alpha \in A$  and  $\beta \in B$  such that  $\alpha > \alpha_1$ ,  $\alpha > \alpha_2$ ,  $\beta > \beta_1$ ,  $\beta > \beta_2$ , because  $A$  and  $B$  are directed sets without maximal elements due to Lemma A.4. Hence  $(\alpha, \beta, 2) \in C$  obviously satisfies

$$(\alpha, \beta, 2) > (\alpha_1, \beta_1, i_1), \quad (\alpha, \beta, 2) > (\alpha_2, \beta_2, i_2).$$

So  $C$  is a directed set.

Let

$$I_{(\alpha, \beta, i)} = \begin{cases} I_\alpha^1 & \text{if } i = 1, \\ I_\beta^2 & \text{if } i = 2. \end{cases}$$

Since  $\{I_\alpha^1\}$  and  $\{I_\beta^2\}$  are van Hove nets, there exists  $\alpha_o \in A$  and  $\beta_o \in B$  for any  $d > 0$  and  $\epsilon > 0$  such that

$$\begin{aligned} \frac{|\text{surf}_d(I_\alpha^1)|}{|I_\alpha^1|} &< \epsilon \quad \text{if } \alpha \geq \alpha_o \\ \frac{|\text{surf}_d(I_\beta^2)|}{|I_\beta^2|} &< \epsilon \quad \text{if } \beta \geq \beta_o. \end{aligned}$$

Set  $\gamma_o \equiv (\alpha_o, \beta_o, 1)$ . For any  $\gamma = (\alpha, \beta, i) \geq \gamma_o$ , we have obviously  $\alpha \geq \alpha_o$  and  $\beta \geq \beta_o$  by the definition of the ordering. Hence,

$$\frac{|\text{surf}_d(I_\gamma)|}{|I_\gamma|} \leq \max \left\{ \frac{|\text{surf}_d(I_\alpha^1)|}{|I_\alpha^1|}, \frac{|\text{surf}_d(I_\beta^2)|}{|I_\beta^2|} \right\} < \epsilon.$$

Thus  $\{I_\gamma\}_{\gamma \in C}$  is also a van Hove net. If  $\{I_\alpha^1\}$  and  $\{I_\beta^2\}$  are van Hove nets tending to  $\mathbb{Z}^\nu$ , then  $\{I_\gamma\}$  is also a van Hove net tending to  $\mathbb{Z}^\nu$  by its definition.

Since  $\{I_\gamma\}_{\gamma \in C}$  is a van Hove net (van Hove net tending to  $\mathbb{Z}^\nu$ ),  $f$  has the following limit by the assumption on  $f$

$$f_\infty = \lim_{\gamma} \{f(I_\gamma), \gamma \in C\}.$$

Thus for any  $\varepsilon$ , there exists a  $\gamma_o = (\alpha_o, \beta_o, 1)$  or  $\gamma_o = (\alpha_o, \beta_o, 2)$  such that

$$|f_\infty - f(I_\gamma)| < \varepsilon$$

for  $\gamma \geq \gamma_o$ . This inequality holds especially for  $\gamma = (\alpha, \beta, 1) \geq \gamma_o$  with  $\alpha > \alpha_o$  and  $\beta > \beta_o$ . For this  $\gamma$ ,  $I_\gamma = I_\alpha^1$ , and hence  $f(I_\gamma) = f(I_\alpha^1)$ . Thus we have

$$|f_\infty - f(I_\alpha^1)| < \varepsilon$$

for  $\alpha > \alpha_o$ . Therefore, we obtain

$$f_\infty = \lim_{\alpha} f(I_\alpha^1).$$

Similarly,

$$f_\infty = \lim_{\beta} f(I_\beta^2).$$

Now we have shown that the limit is the same for  $\{I_\alpha^1\}_{\alpha \in A}$  and  $\{I_\beta^2\}_{\beta \in B}$ . Hence the independence of the limit on the choice of the net follows.  $\square$

**Definition A.6.** If  $f(I_\alpha)$  has a limit for any van Hove net  $\{I_\alpha\}$ , then  $f(I)$  is said to have the van Hove limit for large  $I$ , and its limit is denoted by

$$\text{v.H.} \lim_{I \rightarrow \infty} f(I). \quad (\text{A.4})$$

If  $f(I_\alpha)$  has a limit for any van Hove net  $\{I_\alpha\}$  tending to  $\mathbb{Z}^\nu$ , then  $f(I)$  is said to have the van Hove limit for  $I$  tending to  $\mathbb{Z}^\nu$ , and its limit is denoted by

$$\text{v.H.} \lim_{I \rightarrow \mathbb{Z}^\nu} f(I). \quad (\text{A.5})$$

In general, the first condition is stronger than the second. If  $f(I)$  is translation invariant, however, the existence of the two limits are equivalent as shown below.

**Lemma A.7.** If  $f(I)$  is translation invariant in the sense that

$$f(I + n) = f(I)$$

for any finite subset  $I$  of  $\mathbb{Z}^\nu$  and  $n \in \mathbb{Z}^\nu$ , then  $f(I)$  has the van Hove limit for large  $I$  if and only if  $f$  has the van Hove limit for  $I$  tending to  $\mathbb{Z}^\nu$ .

*Proof.* The only if part is obvious. Let  $\{I_\alpha\}_{\alpha \in A}$  be an arbitrary van Hove net. Let  $a(\alpha)$  be the largest integer  $a$  such that a translate of  $\mathcal{C}_a$  is contained in  $I_\alpha$ . Let  $\mathcal{C}_{a(\alpha)} + n \subset I_\alpha$  and hence  $\mathcal{C}_{a(\alpha)} \subset I_\alpha - n$ . Now we shift an approximate center of  $\mathcal{C}_{a(\alpha)}$  to the origin of  $\mathbb{Z}^\nu$  and simultaneously shift  $I_\alpha - n$  by the same amount. More precisely,  $I_\alpha - n$  is shifted to

$$I'_\alpha \equiv I_\alpha - n - \left\lfloor \frac{a(\alpha) - 1}{2} \right\rfloor (1, \dots, 1).$$

Obviously,

$$\frac{|\text{surf}_d(I'_\alpha)|}{|I'_\alpha|} = \frac{|\text{surf}_d(I_\alpha)|}{|I_\alpha|}$$

for all  $d > 0$  and  $\alpha \in A$ .

We show that this  $\{I'_\alpha\}(\alpha \in A)$  is tending to  $\mathbb{Z}^\nu$ . Let  $I$  be a finite subset of  $\mathbb{Z}^\nu$ . For sufficiently large integer  $a$ ,  $I \subset \mathcal{C}_{a-\lfloor \frac{a-1}{2} \rfloor}$ . For this  $a$ , there exists  $\alpha_1$  such that  $\mathbf{n}_a^-(I_\alpha) > 0$  for  $\alpha \geq \alpha_1$ . Then  $a(\alpha) \geq a$  and

$$I'_\alpha \supset \mathcal{C}_{a(\alpha)-\lfloor \frac{a(\alpha)-1}{2} \rfloor} \supset \mathcal{C}_{a-\lfloor \frac{a-1}{2} \rfloor} \supset I$$

for  $\alpha \geq \alpha_1$ . Thus  $\{I'_\alpha\}(\alpha \in A)$  is a van Hove net tending to  $\mathbb{Z}^\nu$ . Since  $f$  is translation invariant,

$$f(I_\alpha) = f(I'_\alpha).$$

By the assumption that  $f$  has the van Hove limit tending to  $\mathbb{Z}^\nu$ ,  $\lim_\alpha f(I'_\alpha)$  exists, and hence  $\lim_\alpha f(I_\alpha)$  exists.  $\square$